

**SEARCH REQUEST FORM**

Scientific and Technical Information Center

Requester's Full Name: Margaret B. Madley Examiner #: 60850 Date: 7-17-01  
 Art Unit: 1714 Phone Number 30 8-2518 Serial Number: 091 549428 April 2000  
 Mail Box and Bldg/Room Location: 3-3 Results Format Preferred (circle): PAPER DISK E-MAIL

3-4 DO9  
 If more than one search is submitted, please prioritize searches in order finished.

\*\*\*\*\*  
 Please provide a detailed statement of the search topic, and describe as specifically as possible the subject matter to be searched. Include the elected species or structures, keywords, synonyms, acronyms, and registry numbers, and combine with the concept or utility of the invention. Define any terms that may have a special meaning. Give examples or relevant citations, authors, etc, if known. Please attach a copy of the cover sheet, pertinent claims, and abstract.

Title of Invention: Process for the Separation of Enantiomers and  
Enantiopure Reagent  
 Inventors (please provide full names): Delpplanché, Thierry et al

Earliest Priority Filing Date: \_\_\_\_\_

Search in the compels of ds. 10-Band 18  
 \*For Sequence Searches Only\* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.

Search 6

\*\*\*\*\*  
**STAFF USE ONLY**


	Type of Search	Vendors and cost where applicable
Searcher: <u>206</u>	NA Sequence (#) _____	STN <u>✓</u>
Searcher Phone #: <u>8-4139</u>	AA Sequence (#) _____	Dialog _____
Searcher Location: <u>ELC 1700</u>	Structure (#) <u>(27)</u>	Questel/Orbit _____
Date Searcher Picked Up: _____	Bibliographic _____	Dr.Link _____
Date Completed: <u>7-19-01</u>	Litigation _____	Lexis/Nexis _____
Searcher Prep & Review Time: <u>55</u>	Fulltext _____	Sequence Systems _____
Clerical Prep Time: <u>15</u>	Patent Family _____	WWW/Internet _____
Online Time: <u>65</u>	Other _____	Other (specify) _____

L32 80299 SEA FILE=REGISTRY ABB=ON PLU=ON ?PHENYLALANIN?/CNS  
 L33 4672 SEA FILE=REGISTRY ABB=ON PLU=ON ?NITRO?/CNS (L) L32  
 L34 757 SEA FILE=REGISTRY ABB=ON PLU=ON N-4-NITRO? (L) L33  
 L35 1 SEA FILE=REGISTRY ABB=ON PLU=ON 2-METHOXY? (L) L34

=> d 135

L35 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2001 ACS  
 RN 328406-65-1 REGISTRY  
 CN L-Phenylalanine, N-[(4-nitrophenoxy)carbonyl]-, 2-methoxyethyl ester  
 (9CI) (CA INDEX NAME)  
 FS STEREOSEARCH  
 MF C19 H20 N2 O7  
 SR CA  
 LC STN Files: CA, CAPLUS, TOXLIT

Absolute stereochemistry.



1 REFERENCES IN FILE CA (1967 TO DATE)  
 1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

=>

L36 1 SEA FILE=REGISTRY ABB=ON PLU=ON 328406-65-1  
L37 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L36

=> d all hitstr 137

L37 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2001 ACS  
AN 2001:57848 HCAPLUS  
DN 134:212851  
TI Application of (S)-N-(4-nitrophenoxy carbonyl) phenylalanine methoxyethyl ester as a new chiral derivatizing agent for proteinogenic amino acid analysis by high-performance liquid chromatography  
AU Peter, A.; Vekes, E.; Torok, G.  
CS Department of Inorganic and Analytical Chemistry, University of Szeged, Szeged, 6720, Hung.  
SO Chromatographia (2000), 52(11/12), 821-826  
CODEN: CHRGB7; ISSN: 0009-5893  
PB Friedrich Vieweg & Sohn Verlagsgesellschaft mbH  
DT Journal  
LA English  
CC 64-3 (Pharmaceutical Analysis)  
AB The application of (S)-N-(4-nitrophenoxy carbonyl) phenylalanine methoxyethyl ester, (S)-NIFE, as a new chiral derivatizing agent for the resolu. of compds. possessing an amino group is described. Its applicability is demonstrated by the resolu. of proteinogenic amino acid enantiomers. The diastereomeric derivs. produced were sepd. by reversed-phase high-performance liq. chromatog. The effects of pH, excess reagent and reaction time on the derivatization kinetics, and the effects of pH and the org. modifier on the sepn., were investigated.  
ST nitrophenoxy carbonyl phenylalanine deriv. chiral agent HPLC; liq chromatog detn amino acid detn  
IT HPLC  
Reversed phase HPLC  
(sepn. of amino acids by reversed phase HPLC using (S)-N-(4-nitrophenoxy carbonyl) phenylalanine methoxyethyl ester as chiral derivatizing agent)  
IT 52-90-4, L-Cysteine, analysis 56-41-7, L-Alanine, analysis 56-45-1, L-Serine, analysis 56-84-8, L-Aspartic acid, analysis 56-85-9, L-Glutamine, analysis 56-86-0, L-Glutamic acid, analysis 56-87-1, L-Lysine, analysis 60-18-4, L-Tyrosine, analysis 61-90-5, L-Leucine, analysis 63-68-3, L-Methionine, analysis 63-91-2, L-Phenylalanine, analysis 70-47-3, L-Asparagine, analysis 71-00-1, L-Histidine, analysis 72-18-4, L-Valine, analysis 72-19-5, L-Threonine, analysis 73-22-3, L-Tryptophan, analysis 73-32-5, L-Isoleucine, analysis 74-79-3, L-Arginine, analysis 147-85-3, L-Proline, analysis 302837-19-0 302837-20-3 302837-22-5 302837-24-7 302837-25-8 302837-27-0 302837-29-2 302837-30-5 302837-31-6 302837-34-9 302837-35-0 302837-36-1 328406-66-2 328406-67-3 328406-68-4 328406-69-5 328406-70-8 328406-71-9 328406-72-0  
RL: ANT (Analyte); ANST (Analytical study)  
(sepn. of amino acids by reversed phase HPLC using (S)-N-(4-nitrophenoxy carbonyl) phenylalanine methoxyethyl ester as chiral derivatizing agent)  
IT 328406-65-1  
RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)  
(sepn. of amino acids by reversed phase HPLC using (S)-N-(4-nitrophenoxy carbonyl) phenylalanine methoxyethyl ester as chiral derivatizing agent)  
RE.CNT 11  
RE

- (1) Anon; J Biol Chem 1989, V264, P668
- (2) Beesley, T; Chiral Chromatography 1998
- (3) Bojarski, J; Chem Anal 1997, V42, P139 HCAPLUS
- (4) Delplanche, T; Peptidomimetics Symposium 1999, 11
- (5) Gorog, S; Chromatogr B 1994, V659, P51 MEDLINE
- (6) Kleidernigg, O; Chromatographia 1997, V44, P465 HCAPLUS
- (7) Lunn, G; Handbook of Derivatization Reactions for HPLC 1998
- (8) Nimura, N; J Chromatogr 1980, V202, P375 HCAPLUS
- (9) Okamoto, Y; Chromatographic Enantiomer Separation on Chiral Polymers 1997
- (10) Solvay; patent pending
- (11) Toyo'oka, T; Modern Derivatization Methods for Separation Science 1999


IT 328406-65-1

RL: ARG (Analytical reagent use); ANST (Analytical study); USES (Uses)  
 (sepn. of amino acids by reversed phase HPLC using (S)-N-(4-nitrophenoxycarbonyl) phenylalanine methoxyethyl ester as chiral derivatizing agent)

RN 328406-65-1 HCAPLUS


CN L-Phenylalanine, N-[(4-nitrophenoxy) carbonyl]-, 2-methoxyethyl ester (9CI)  
 (CA INDEX NAME)

Absolute stereochemistry.



L1

STR



## NODE ATTRIBUTES:

HCOUNT	IS M1	AT	10
NSPEC	IS R	AT	1
NSPEC	IS R	AT	2
NSPEC	IS R	AT	3
NSPEC	IS R	AT	4
NSPEC	IS R	AT	5
NSPEC	IS R	AT	6
NSPEC	IS C	AT	7
NSPEC	IS C	AT	8
NSPEC	IS C	AT	9
NSPEC	IS C	AT	10
NSPEC	IS C	AT	11
NSPEC	IS C	AT	12
NSPEC	IS C	AT	13
NSPEC	IS C	AT	14

DEFAULT MLEVEL IS ATOM

MLEVEL IS CLASS AT 7 8 9 10 11 12 13 14

DEFAULT ECLEVEL IS LIMITED

## GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 14

## STEREO ATTRIBUTES: NONE

L3	501	SEA FILE=REGISTRY SSS FUL L1	
L4	213	SEA FILE=HCAPLUS ABB=ON PLU=ON	L3
L6	1	SEA FILE=HCAPLUS ABB=ON PLU=ON	2000:755247/AN
L14	75483	SEA FILE=HCAPLUS ABB=ON PLU=ON	(AMINO ACIDS OR PEPTIDES OR PROTEINS)/CC
L15	83	SEA FILE=HCAPLUS ABB=ON PLU=ON	L4 AND L14
L16	82	SEA FILE=HCAPLUS ABB=ON PLU=ON	L15 NOT L6
L18	4	SEA FILE=HCAPLUS ABB=ON PLU=ON	(HPLC OR CHROMATOGARPH?) AND L16
L19	9	SEA FILE=HCAPLUS ABB=ON PLU=ON	(ENANTIO? OR SEPARAT? OR CHROMOPHO? OR REAGENT) AND L16
L20	12	SEA FILE=HCAPLUS ABB=ON PLU=ON	L18 OR L19

=&gt;

L20 ANSWER 1 OF 12 HCAPLUS COPYRIGHT 2001 ACS

AN 2001:220249 HCAPLUS

DN 134:237834

TI Method for preparation of 2-[[N-(hydrocarbyloxycarbonyl)-L-alanyl]amino]thiazole-4-acetic acid ester derivatives

IN Hirota, Yoshihiro; Matsunaga, Tomonori; Iwasaki, Fumiaki

PA Tokuyama Corp., Japan

SO Jpn. Kokai Tokkyo Koho, 12 pp.

CODEN: JKXXAF

DT Patent

LA Japanese


IC ICM C07D277-44

CC 34-2 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 1, 26

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 2001081083	A2	20010327	JP 1999-259786	19990914
OS	CASREACT 134:237834; MARPAT 134:237834				
GI					



AB The title compds. (I; R1 = C1-6 alkyl, C6-8 aryl, CH2Ph; R2 = C1-6 alkyl, Ph, CH2Ph) are prepd. by condensation of N-(hydrocarbyloxycarbonyl)-L-alanine (II; R1 = same as above) with 2-(2-amino-5-thiazolyl)-2-methoxyiminoacetic acid ester (III; R2 = same as above) using a condensing agent, more specifically N,N'-carbonyldiimidazole and converted into 2-[[N-(hydrocarbyloxycarbonyl)-L-alanyl]amino]thiazol-4-yl]-2-methoxyiminoacetic acid I (R1 = same as above; R2 = H) by hydrolysis in the presence of base and neutralization. The latter 2-(2-aminothiazol-4-yl)-2-methoxyiminoacetic acid derivs. are useful as intermediates for drugs, and in particular used as side chains for cephalosporin antibiotics. N,N'-carbonyldiimidazole is superior to other condensing agents such as dicyclohexylcarbodiimide and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride and gives I in high yields. This process also gives III with high purity, enables the skipping of purifn. step for III prior to sapon., and simplifies and significantly improves the prodn. efficiency for the sapond. product, i.e. free acid I (R1 = same as above; R2 = H). Thus, 47.3 g N-tert-butoxycarbonyl-L-

alanine and 57.3 g 2-(2-aminothiazol-4-yl)-2-(Z)-(methoxyimino)acetic acid Et ester were added to 250 mL EtOAc, cooled to 0.degree., treated slowly with 40.5 g N,N'-carbonyldiimidazole over a period of 10 min at .ltoreq.5.degree., and stirred at 4.degree. for 1 h. The reaction mixt. was washed with 2 N HCl twice and 2 N NaOH and the org. layer was **sepd.** to give, after evapn. of the solvent and silica gel chromatog., 90.4 g I (R1 = tert-Bu, R2 = Et) (90.0% yield). In a sapon. step, the org. layer obtained above was distd. in vacuo to the wt. of 113.2 g, treated with 67.2 g MeOH, and then slowly with 125 mL 3 N NaOH at .ltoreq.25.degree. over a period of 15 min, and allowed to react at 25.degree. for 5 h. The solvent was distd. in vacuo from the reaction mixt. until the mixt. weighed at 187.2 g, followed by adding 130 g H2O, and the resulting mixt. was cooled 7.degree., treated slowly with 260 mL 2 N HCl over a period of 1 h, and stirred at .ltoreq.5.degree. for 2 h to give, after centrifugation and washing the **sepd.** solid with water and drying, 81.1 g I (R1 = tert-Bu, R2 = H) (87.0% yield).

ST alanylaminothiazolylmethoxyiminoacetic acid ester prepn intermediate cephalosporin antibiotic; hydrocarbyloxycarbonylalanine condensation aminothiazolylmethoxyiminoacetic acid ester; methoxyiminoacetic acid ester alanyl aminothiazolyl prepn intermediate cephalosporin antibiotic

IT Condensation reaction  
(prepn. of [[N-(hydrocarbyloxycarbonyl)-L-alanyl]amino]thiazoleacetic acid ester derivs. by condensation of N-(hydrocarbyloxycarbonyl)-L-alanine with (aminothiazolyl)methoxyiminoacetic acid ester using N,N'-carbonyldiimidazole)

IT Lactams  
RL: PNU (Preparation, unclassified); PREP (Preparation)  
(.beta.-, antibiotics; prepn. of [[N-(hydrocarbyloxycarbonyl)-L-alanyl]amino]thiazoleacetic acid ester derivs. by condensation of N-(hydrocarbyloxycarbonyl)-L-alanine with (aminothiazolyl)methoxyiminoacetic acid ester using N,N'-carbonyldiimidazole)

IT 530-62-1, N,N'-Carbonyldiimidazole 1142-20-7 15761-38-3 16639-86-4, N-Ethoxycarbonyl-L-alanine 33294-53-0 64485-88-7 65243-09-6  
**141695-27-4**, N-Pheoxycarbonyl-L-alanine 162281-00-7  
RL: RCT (Reactant)  
(prepn. of [[N-(hydrocarbyloxycarbonyl)-L-alanyl]amino]thiazoleacetic acid ester derivs. by condensation of N-(hydrocarbyloxycarbonyl)-L-alanine with (aminothiazolyl)methoxyiminoacetic acid ester using N,N'-carbonyldiimidazole)


IT 330566-03-5P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of [[N-(hydrocarbyloxycarbonyl)-L-alanyl]amino]thiazoleacetic acid ester derivs. by condensation of N-(hydrocarbyloxycarbonyl)-L-alanine with (aminothiazolyl)methoxyiminoacetic acid ester using N,N'-carbonyldiimidazole)

IT 88970-81-4P 330566-04-6P 330566-05-7P 330566-06-8P 330566-07-9P 330566-08-0P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of [[N-(hydrocarbyloxycarbonyl)-L-alanyl]amino]thiazoleacetic acid ester derivs. by condensation of N-(hydrocarbyloxycarbonyl)-L-alanine with (aminothiazolyl)methoxyiminoacetic acid ester using N,N'-carbonyldiimidazole)


IT **141695-27-4**, N-Pheoxycarbonyl-L-alanine  
RL: RCT (Reactant)  
(prepn. of [[N-(hydrocarbyloxycarbonyl)-L-alanyl]amino]thiazoleacetic acid ester derivs. by condensation of N-(hydrocarbyloxycarbonyl)-L-alanine with (aminothiazolyl)methoxyiminoacetic acid ester using N,N'-carbonyldiimidazole)

RN 141695-27-4 HCAPLUS  
CN L-Alanine, N-(phenoxycarbonyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L20 ANSWER 2 OF 12 HCAPLUS COPYRIGHT 2001 ACS  
AN 2000:853644 HCAPLUS  
DN 134:193704  
TI A solid-phase approach to analogues of the antibiotic mureidomycin  
AU Bozzoli, Andrea; Kazmierski, Wieslaw; Kennedy, Gordon; Pasquarello, Alessandra; Pecunioso, Angelo  
CS GlaxoWellcome SpA, Medicines Research Centre, Verona, 37135, Italy  
SO Bioorg. Med. Chem. Lett. (2000), 10(24), 2759-2763  
CODEN: BMCLE8; ISSN: 0960-894X  
PB Elsevier Science Ltd.  
DT Journal  
LA English  
CC 34-3 (Amino Acids, Peptides, and Proteins)  
Section cross-reference(s): 26, 33  
GI



AB A library of 80 analogs of the antibacterial family of mureidomycins [(I); R = H, mureidomycin A; R = Gly, mureidomycin C], was prepd. using solid-phase chem. techniques. Analog fragments (2,3-diaminopropionic acid, methionine, p-tyrosine, m-tyrosine, as enantiomers, and tert-Bu 2-Ph malonate, as a racemate), were used in the synthesis. A selection of ten of the compds. with best a/a values (by LC-MS anal.) was presented. No inhibitory activity on the growth of *S. aureus* 853, *E. coli* 1952, *Saccharomyces cerevisiae* NCY81, *P. aeruginosa* and *P. aeruginosa* 2033 were detected for any library member.  
ST mureidomycin analog combinatorial library prepn solid phase MSBAR  
IT Combinatorial library  
Solid phase synthesis  
Structure-activity relationship  
(prepn. of a combinatorial library of mureidomycin analogs using



solid-phase synthesis)

IT Natural products  
 RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (prepn. of a combinatorial library of mureidomycin analogs using solid-phase synthesis)

IT 114797-04-5DP, Mureidomycin A, analogs 327184-82-7P 327184-83-8P  
 327184-84-9P 327184-85-0P 327184-86-1P 327184-87-2P 327184-88-3P  
 327184-89-4P 327184-90-7P 327184-91-8P  
 RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (prepn. of a combinatorial library of mureidomycin analogs using solid-phase synthesis)

IT 348-67-4, D-Methionine 362-43-6 673-06-3, D-Phenylalanine 7693-46-1  
 78342-42-4 109838-85-9 198544-42-2 **327184-80-5** 327184-92-9  
 327184-93-0  
 RL: RCT (Reactant)  
 (prepn. of a combinatorial library of mureidomycin analogs using solid-phase synthesis)

IT 15083-05-3P 15083-09-7P 68691-77-0P 265321-20-8P 327184-50-9P  
 327184-51-0P 327184-52-1P 327184-53-2DP, resin-bound 327184-53-2P  
 327184-54-3P 327184-55-4P 327184-56-5P 327184-57-6P 327184-58-7P  
 327184-59-8P 327184-60-1P 327184-61-2P 327184-62-3P 327184-63-4P  
 327184-64-5P 327184-65-6P 327184-66-7P 327184-67-8P 327184-68-9P  
**327184-69-0P 327184-70-3P** 327184-71-4DP, resin-bound  
 327184-72-5DP, resin-bound 327184-73-6DP, resin-bound 327184-76-9DP,  
 resin-bound 327184-78-1DP, resin-bound 327184-79-2DP, resin-bound  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of a combinatorial library of mureidomycin analogs using solid-phase synthesis)

IT 327184-74-7P 327184-75-8P 327184-77-0P 327184-81-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of a combinatorial library of mureidomycin analogs using solid-phase synthesis)

RE.CNT 20


RE

- (1) Armstrong, A; Tetrahedron Lett 1988, V29, P2483 HCAPLUS
- (2) Brandish, P; Antimicrob Agents Chemother 1996, V40, P1640 HCAPLUS
- (3) Brandish, P; J Biol Chem 1996, V271, P7609 HCAPLUS
- (4) Bugg, T; J Chem Soc, Perkin Trans 1 1999, P1279
- (5) Bugg, T; J Chem Soc, Perkin Trans 1 1999, P1285
- (6) Chang, C; Int J Pept Protein Res 1980, V15, P59 HCAPLUS
- (7) Inukai, M; Antimicrob Agents and Chemother 1993, V37, P980 HCAPLUS
- (8) Isono, F; Antimicrob Agents Chemother 1991, V35, P234 HCAPLUS
- (9) Isono, F; J Antibiot 1989, V42, P667 HCAPLUS
- (10) Isono, F; J Antibiot 1989, V62, P674
- (11) Kaiser, E; Anal Biochem 1970, V34, P595 HCAPLUS
- (12) Lee, V; Med Res Rev 1999, V19, P521 HCAPLUS
- (13) Merrifield, R; J Org Chem 1993, V58, P5167
- (14) Niccolai, D; Chem Commun 1997, P2333 HCAPLUS
- (15) Pedroso, E; Tetrahedron Lett 1993, V34, P2195
- (16) Raju, B; Bioorg Med Chem Lett 1998, V8, P3043 HCAPLUS
- (17) Roth, V; Emerging Therapeutic Targets 1999, V3, P73 HCAPLUS
- (18) Schollkopf, U; Angew Chem, Int Ed Engl 1981, V20, P798
- (19) Setti, E; Drugs Future 1997, V22, P271 HCAPLUS
- (20) Williams, R; Synthesis of Optically Active .alpha.-Amino Acids 1989

IT **327184-80-5**  
 RL: RCT (Reactant)  
 (prepn. of a combinatorial library of mureidomycin analogs using solid-phase synthesis)

RN 327184-80-5 HCAPLUS  
CN D-Phenylalanine, N-[(4-nitrophenoxy)carbonyl]-, 1,1-dimethylethyl ester  
(9CI) (CA INDEX NAME)


Absolute stereochemistry.



IT 327184-69-0P 327184-70-3P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of a combinatorial library of mureidomycin analogs using  
solid-phase synthesis)


RN 327184-69-0 HCAPLUS  
CN D-Phenylalanine, 3-(1,1-dimethylethoxy)-N-[(4-nitrophenoxy)carbonyl]-,  
1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 327184-70-3 HCAPLUS  
CN L-Phenylalanine, 3-(1,1-dimethylethoxy)-N-[(4-nitrophenoxy)carbonyl]-,  
1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L20 ANSWER 3 OF 12 HCAPLUS COPYRIGHT 2001 ACS  
AN 2000:421093 HCAPLUS  
DN 133:43809  
TI Preparation of new biphenyl and biphenyl-analogous compounds as integrin  
antagonists  
IN Albers, Markus; Urbahns, Klaus; Vaupel, Andrea; Harter, Michael; Schmidt,

Delf; Stelte-ludwig, Beatrix; Gerdes, Christoph; Stahl, Elke; Keldenich, Jorg; Bruggemeier, Ulf; Lustig, Klemens  
 PA Bayer Aktiengesellschaft, Germany; et al.  
 SO PCT Int. Appl., 360 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM C07C311-19  
 ICS C07C275-42; C07C311-10; C07C311-47; C07D213-40; C07D213-75;  
 C07D235-30; A61K031-18; A61K031-44; A61K031-4184; A61P035-00;  
 A61P019-10; A61P027-02  
 CC 34-2 (Amino Acids, Peptides, and  
 Proteins)  
 Section cross-reference(s): 1, 25

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000035864	A1	20000622	WO 1999-EP9843	19991213
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRAI US 1998-213381	A	19981216		
OS MARPAT 133:43809				
AB Biphenyl compds. R1O2CCHR2-U-V-A-B-W-NR3-C-R4 [R1 = H, (un)substituted alkyl, cycloalkyl, aryl, or (un)satd. heterocyclyl; R2 = H, (un)substituted alkyl, cycloalkyl, aryl, or (un)satd. heterocyclyl, alkenyl, alkynyl, -NR2'SO2R2'', -NR2'CO2R2'', -NR2'COR2'', -NR2'CONR2'2, -NR2'CSNR2'2 (R2' has same definition as R1 and R2'' has same definition as R1 except it is not H); U or W is a direct bond or (un)substituted alkylene; V = (un)substituted alkylene, -NR2'CO- or NR2'SO2-; A and B = (un)substituted 1,3- or 1,4-bridging phenylene group or a 2,4- or 2,5-bridging thienylene group, each of which may have substituents; C is a direct bond, CMe(:X-R5)-Y-N(R6)- (R5 is absent, H, (un)substituted alkyl or cycloalkyl, NO2, acyl, carboxylic or carboxylate group or is connected to R3, Y, R4 or R6, if present; R6 is H, (un)substituted alkyl, cycloalkyl, aryl, or (un)satd. heterocyclyl, an alkylamine or alkylamide residue, or is connected to one of R3, R4, Y, or R5, if present, to form a heterocyclic ring system; X = CHNO2, CHCN, O, N or S; Y is a direct bond or (un)substituted alkylene or alkyne group) or 3,4-dioxo-1,2- cyclobutenediyl-NR6-; R3, R4 = H, (un)substituted alkyl, cycloalkyl, aryl, or (un)satd. heterocyclyl, an alkylamine or alkylamide residue, or is connected to one of R4 (or R3), Y, R5 or R6, if present, to form a heterocyclic ring system] were prepd. as integrin antagonists. Thus, (2R,S)-3-[3-(pyridin-3-ylmethylureido)biphenyl-4-yl]-2-[2,4,6- trimethylbenzenesulfonylamino]propanoic acid, prepd. by reactions of resin-bound (2R,S)-3-(4-bromophenyl)-2-(9-fluorenylmethoxycarbonylamino)pr opanoic acid with sulfonylating, boronic acid, and amine reagents (mesitylenesulfonyl chloride, 3-nitrobenzeneboronic acid, and 2-aminomethylpyridine), showed IC50 = 5 nM for binding to the .alpha.v.beta.3 receptor and IC50 = 480 nM in the smooth muscle cell migration test. ST amino acid biphenyl deriv prepn antagonist integrin IT Angiogenesis Antitumor agents				

Arteriosclerosis

Eye, disease

Osteoporosis

Rheumatoid arthritis

(prepn. of new biphenyl and biphenyl-analogous compds. as integrin antagonists)

IT Amino acids, preparation

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of new biphenyl and biphenyl-analogous compds. as integrin antagonists)

IT Artery, disease

(restenosis; prepn. of new biphenyl and biphenyl-analogous compds. as integrin antagonists)

IT Integrins

RL: BPR (Biological process); BIOL (Biological study); PROC (Process) (.alpha.v.beta.3; prepn. of new biphenyl and biphenyl-analogous compds. as integrin antagonists)

IT	276257-75-1P	276257-76-2P	276257-77-3P	276257-78-4P	276257-79-5P
	276257-80-8P	276257-81-9P	276257-82-0P	276257-83-1P	276257-84-2P
	276257-85-3P	276257-86-4P	276257-87-5P	276257-88-6P	276257-89-7P
	276257-90-0P	276257-91-1P	276257-92-2P	276257-93-3P	276257-94-4P
	276257-95-5P	276257-96-6P	276257-97-7P	276257-98-8P	276257-99-9P
	276258-00-5P	276258-01-6P	276258-02-7P	276258-03-8P	276258-04-9P
	276258-05-0P	276258-06-1P	276258-07-2P	276258-08-3P	276258-09-4P
	276258-10-7P	276258-11-8P	276258-12-9P	276258-13-0P	276258-14-1P
	276258-15-2P	276258-16-3P	276258-17-4P	276258-18-5P	276258-19-6P
	276258-20-9P	276258-21-0P	276258-22-1P	276258-23-2P	276258-24-3P
	276258-25-4P	276258-26-5P	276258-27-6P	276258-28-7P	276258-29-8P
	276258-30-1P	276258-31-2P	276258-32-3P	276258-33-4P	276258-34-5P
	276258-35-6P	276258-36-7P	276258-37-8P	276258-38-9P	276258-39-0P
	276258-40-3P	276258-41-4P	276258-42-5P	276258-43-6P	276258-44-7P
	276258-45-8P	276258-46-9P	276258-47-0P	276258-48-1P	276258-49-2P
	276258-50-5P	276258-51-6P	276258-52-7P	276258-53-8P	276258-54-9P
	276258-55-0P	276258-56-1P	276258-57-2P	276258-58-3P	276258-59-4P
	276258-60-7P	276258-61-8P	276258-62-9P	276258-63-0P	276258-64-1P
	276258-65-2P	276258-66-3P	276258-67-4P	276258-68-5P	276258-69-6P
	276258-70-9P	276258-71-0P	276258-72-1P	276258-73-2P	276258-74-3P
	276258-75-4P	276258-76-5P	276258-79-8P	276258-80-1P	276258-82-3P
	276258-83-4P	276258-84-5P	276258-85-6P	276258-88-9P	276258-94-7P
	276258-97-0P	276259-01-9P	276259-02-0P	276259-04-2P	276259-07-5P
	276259-10-0P	276259-13-3P	276259-15-5P	276259-17-7P	276259-22-4P
	276259-26-8P	276259-27-9P	276259-28-0P	276259-29-1P	276259-30-4P
	276259-31-5P	276259-32-6P	276259-33-7P	276259-34-8P	276259-35-9P
	276259-36-0P	276259-37-1P	276259-38-2P	276259-39-3P	276259-40-6P
	276259-41-7P	276259-42-8P	276259-43-9P	276259-44-0P	276259-45-1P
	276259-46-2P	276259-47-3P	276259-48-4P	276259-49-5P	276259-50-8P
	276259-51-9P	276259-52-0P	276259-53-1P	276259-54-2P	276259-55-3P
	276259-56-4P	276259-57-5P	276259-58-6P	276259-59-7P	276259-60-0P
	276259-61-1P	276259-62-2P	276259-63-3P	276259-64-4P	276259-65-5P
	276259-66-6P	276259-67-7P	276259-68-8P	276259-69-9P	276259-70-2P
	276259-71-3P	276259-72-4P	276259-73-5P	276259-74-6P	276259-75-7P
	276259-76-8P	276259-77-9P	276259-78-0P	276259-79-1P	276259-80-4P
	276259-81-5P	276259-82-6P	276259-83-7P	276259-84-8P	276259-85-9P
	276259-86-0P	276259-87-1P	276259-88-2P	276259-89-3P	276259-90-6P
	276259-91-7P	276259-92-8P	276259-93-9P	276259-94-0P	276259-95-1P
	276259-96-2P	276259-97-3P	276259-98-4P	276259-99-5P	276260-00-5P
	276260-01-6P	276260-02-7P	276260-03-8P	276260-04-9P	276260-05-0P
	276260-06-1P	276260-07-2P	276260-08-3P	276260-09-4P	276260-10-7P

276260-11-8P	276260-12-9P	276260-13-0P	276260-14-1P	276260-15-2P
276260-16-3P	276260-17-4P	276260-18-5P	276260-19-6P	276260-20-9P
276260-21-0P	276260-22-1P	276260-23-2P	276260-24-3P	276260-25-4P
276260-26-5P	<b>276260-27-6P</b>	276260-28-7P	276260-29-8P	
276260-30-1P	276260-31-2P	276260-32-3P	276260-33-4P	276260-34-5P
276260-35-6P	276260-36-7P	276260-37-8P	276260-38-9P	276260-39-0P
276260-40-3P				

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of new biphenyl and biphenyl-analogous compds. as integrin antagonists)

IT	276260-41-4P	276260-42-5P	276260-43-6P	276260-44-7P	276260-45-8P
	276260-46-9P	276260-47-0P	276260-48-1P	276260-49-2P	276260-50-5P
	276260-51-6P	276260-52-7P	276260-53-8P	276260-54-9P	276260-55-0P
	276260-56-1P	276260-57-2P	276260-58-3P	276260-59-4P	276260-60-7P
	276260-61-8P	276260-62-9P	276260-63-0P	276260-64-1P	276260-65-2P
	276260-66-3P	276260-67-4P	276260-68-5P	276260-69-6P	276260-70-9P
	276260-71-0P	276260-72-1P	276260-73-2P	276260-74-3P	276260-75-4P
	276260-76-5P	276260-77-6P	276260-78-7P	276260-79-8P	276260-80-1P
	276260-81-2P	276260-82-3P	276260-83-4P	276260-84-5P	276260-85-6P
	276260-86-7P	276260-87-8P	276260-88-9P	276260-89-0P	276260-91-4P
	276260-93-6P	276260-95-8P	276260-96-9P	276260-97-0P	276260-98-1P
	276260-99-2P	276261-00-8P	276261-01-9P	276261-02-0P	276261-03-1P
	276261-04-2P	276261-05-3P	276261-06-4P	276261-07-5P	276261-08-6P
	276261-09-7P	276261-10-0P	276261-12-2P	276261-14-4P	276261-15-5P
	276261-17-7P	276261-19-9P	276261-21-3P	276261-22-4P	276261-23-5P
	276261-24-6P	276261-25-7P	276261-26-8P	276261-27-9P	276261-28-0P
	276261-29-1P	276261-30-4P	276261-31-5P	276261-32-6P	276261-33-7P
	276261-34-8P	276261-35-9P	276261-36-0P	276261-37-1P	276261-38-2P
	276261-39-3P	276261-41-7P	276261-43-9P	276261-45-1P	276261-47-3P
	276261-48-4P	276261-50-8P	276261-51-9P	276261-53-1P	276261-54-2P
	276261-56-4P	276261-57-5P	276261-58-6P	276261-60-0P	276261-62-2P
	276261-64-4P	276261-65-5P	276261-67-7P	276261-69-9P	276261-71-3P
	276261-72-4P	276261-74-6P	276261-75-7P	276261-76-8P	276261-77-9P
	276261-78-0P	276261-79-1P	276261-80-4P	276261-81-5P	276261-82-6P
	276261-83-7P	276261-84-8P	276261-85-9P	276261-86-0P	276261-87-1P
	276261-88-2P	276261-89-3P	276261-90-6P	276261-91-7P	276261-92-8P
	276261-93-9P	276261-94-0P	276261-95-1P	276261-96-2P	276261-97-3P
	276261-98-4P	276261-99-5P	276262-00-1P	276262-01-2P	276262-02-3P
	276262-03-4P	276262-04-5P	276262-05-6P	276262-06-7P	276262-07-8P
	276262-08-9P	276262-09-0P	276262-10-3P	276262-11-4P	276262-12-5P
	276262-13-6P	276262-14-7P	276262-15-8P	276262-16-9P	276262-17-0P
	276262-18-1P	276262-19-2P	276262-20-5P	276262-21-6P	276262-22-7P
	276262-23-8P	276262-24-9P	276262-25-0P	276262-26-1P	276262-27-2P
	276262-28-3P	276262-29-4P	276262-30-7P	276262-31-8P	276262-32-9P
	276262-33-0P	276262-34-1P	276262-35-2P	276262-36-3P	276262-37-4P
	276262-38-5P	276262-39-6P	276262-40-9P	276262-41-0P	276262-42-1P
	276262-43-2P	276262-44-3P	276262-45-4P	276262-46-5P	276262-47-6P
	276262-48-7P	276262-49-8P	276262-50-1P	276262-51-2P	276262-52-3P
	276262-53-4P	276262-54-5P	276262-55-6P	276262-56-7P	276262-57-8P
	276262-58-9P	276262-59-0P	276262-60-3P	276262-61-4P	276262-62-5P
	276262-63-6P	276262-64-7P	276262-65-8P	276262-66-9P	276262-67-0P
	276262-68-1P	276262-69-2P			

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of new biphenyl and biphenyl-analogous compds. as integrin antagonists)

IT 51-45-6, 2-(Imidazol-4-yl)ethylamine, reactions 78-81-9, Isobutylamine

88-14-2, 2-Furancarboxylic acid 96-15-1, 2-Methylbutylamine 96-50-4,  
 2-Aminothiazole 98-58-8, 4-Bromobenzenesulfonyl chloride 98-60-2,  
 4-Chlorobenzenesulfonyl chloride 107-15-3, 1,2-Ethanediamine, reactions  
 108-00-9, n,n-Dimethylethylenediamine 108-91-8, Cyclohexylamine,  
 reactions 462-08-8, 3-Aminopyridine 504-24-5, 4-Aminopyridine  
 504-29-0, 2-Aminopyridine 645-36-3, Aminoacetaldehyde diethyl acetal  
 765-30-0, Cyclopropylamine 773-64-8, 2,4,6-Trimethylbenzenesulfonyl  
 chloride 934-32-7, 2-Aminobenzimidazole 1001-53-2,  
 n-Acetyleneethylenediamine 1003-03-8, Cyclopentylamine 2905-23-9,  
 2-Chlorobenzenesulfonyl chloride 2905-24-0, 3-Bromobenzenesulfonyl  
 chloride 2991-42-6, 4-Trifluoromethylbenzenesulfonyl chloride  
 3731-51-9, 2-Aminomethylpyridine 3731-52-0, 3-Aminomethylpyridine  
 3731-53-1, 4-Aminomethylpyridine 4548-45-2, 2-Chloro-5-nitropyridine  
 4659-45-4, 2,6-Dichlorobenzoyl chloride 4795-29-3, 2-  
 Aminomethyltetrahydrofuran 5231-87-8 5402-73-3, 2,5-  
 Dichlorobenzenesulfonyl chloride 6335-76-8 7154-73-6,  
 2-Pyrrolidin-1-ylethylamine 7693-46-1, 4-Nitrophenyl chloroformate  
 10191-60-3, Dimethyl cyanimidodithiocarbonate 13258-63-4,  
 4-Pyridineethanamine 13331-27-6 13623-94-4 13952-84-6,  
 sec-Butylamine 20781-20-8, 2,4-Dimethoxybenzylamine 21286-54-4, +  
 Camphor 10 sulfonyl chloride 22374-89-6 23095-05-8,  
 5-Bromo-2-methoxybenzenesulfonyl chloride 29022-11-5, Fmoc gly oh  
 50998-05-5 66472-86-4 79711-73-2 79844-65-8 80500-27-2  
 87199-16-4, 3-Formylbenzeneboronic acid 87199-17-5, 4-  
 Formylbenzeneboronic acid 88831-43-0 99359-32-7 107819-90-9  
 126727-04-6 180181-93-5 276262-70-5 276262-71-6

RL: RCT (Reactant)

(prepn. of new biphenyl and biphenyl-analogous compds. as integrin  
 antagonists)

IT	276258-77-6P	276258-78-7P	276258-81-2P	276258-86-7P	276258-87-8P
	276258-89-0P	276258-90-3P	276258-91-4P	276258-92-5P	276258-93-6P
	276258-95-8P	276258-96-9P	276258-98-1P	276258-99-2P	276259-00-8P
	276259-03-1P	276259-05-3P	276259-06-4P	276259-08-6P	276259-09-7P
	276259-11-1P	276259-12-2P	276259-14-4P	276259-16-6P	276259-18-8P
	276259-19-9P	276259-20-2P	276259-21-3P	276259-23-5P	276259-24-6P
	276259-25-7P	276262-72-7P	276262-73-8P		

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(prepn. of new biphenyl and biphenyl-analogous compds. as integrin  
 antagonists)

RE.CNT 4

RE

- (1) Anon; WO 9736859 A 1997 HCAPLUS
- (2) Merck, P; DE 19548709 A 1997 HCAPLUS
- (3) Merck, P; WO 9800395 A 1998 HCAPLUS
- (4) Tanabe, S; WO 9936393 A 1999 HCAPLUS

IT 276260-27-6P


RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic  
 preparation); THU (Therapeutic use); BIOL (Biological study); PREP  
 (Preparation); USES (Uses)

(prepn. of new biphenyl and biphenyl-analogous compds. as integrin  
 antagonists)

RN 276260-27-6 HCAPLUS

CN [1,1'-Biphenyl]-4-propanoic acid, alpha.-[[[4-  
 methoxyphenoxy)carbonyl]amino]-3'-[[[propylamino)carbonyl]amino]-,  
 (.alpha.S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry:



L20 ANSWER 4 OF 12 HCAPLUS COPYRIGHT 2001 ACS

AN 1999:584481 HCAPLUS

DN 132:12483

TI **Enantiomeric Separation of (.+.)-Amino Acid Esters on (-)-Phenylurea Chiral Stationary Phase**

AU Lee, Kwang-Pill; Lee, Hyun-Bong; Lee, Young Cheol; Choi, Seong-Ho; Ryoo, Jae Jeong; Park, Jung Hag

CS Department of Chemistry, Graduate School, Kyungpook National University, Sangeok-dong, Taegu, 702-701, S. Korea

SO Microchem. J. (1999), 63(1), 18-23

CODEN: MICJAN; ISSN: 0026-265X

PB Academic Press

DT Journal

LA English

CC 34-2 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 9

AB The 3,5-dinitrobenzoyl (.+.)-amino acid esters were successfully resolved on (-)-phenylurea chiral stationary phases (CSPs) in a normal phase mode by high-performance liq. chromatog. (HPLC). The alcs. used for esterification were methanol, ethanol, and n-propanol. The effects of esterification were studied via retention and optical resolu. The solvent and its concn. effect on **enantioselectivity** have been investigated based on the binary or ternary solvent system. The alc. used in the binary or ternary solvent system was crit. to the **enantiomeric** resolu. of 3,5-dinitrobenzoyl amino acid esters while the nonalcoholic solvent was not suitable. The optical condition of the **enantiomeric** resolu. is discussed in terms of the solvent compn. and structure of the amino acid esters. The main chiral recognition mechanism based on the .pi.-.pi. interaction of the nitrobenzoyl group of the amino acid derivs. with the .pi.-basic Ph group of CSPs is described. (c) 1999 Academic Press.

ST amino acid ester **enantiomeric sepn** phenylurea chiral stationary phase; chromatog chiral stationary phase **sepn** amino acid ester **enantiomeric**

IT Chiral recognition

Chromatography

Esterification

Resolution (**separation**)

(**enantiomeric sepn** of (.+.) amino acid esters on (-)phenylurea chiral stationary phase)

IT Amino acids, preparation

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(**enantiomeric sepn** of (.+.) amino acid esters on (-)phenylurea chiral stationary phase)

IT 64-10-8, Phenylurea

RL: RCT (Reactant)

(chiral stationary phase for the resolu. of (.+.)-amino acid esters)

IT 35519-07-4 74928-18-0 74928-20-4 74928-21-5 74928-22-6  
 74928-24-8 92915-46-3 104336-95-0 106145-09-9 135088-75-4  
 135088-79-8 135449-37-5 135449-39-7 138088-19-4 138088-20-7  
 138088-21-8 138088-22-9 148346-54-7 187836-09-5 214782-91-9  
 251538-48-4 251538-49-5 251538-50-8 251538-51-9 251538-52-0  
 251538-54-2 251538-55-3 251538-56-4 251538-57-5 251538-58-6  
 251538-59-7 251538-60-0 251538-61-1 251538-62-2 251538-63-3  
 251538-64-4 **251538-65-5** 251538-66-6 251538-67-7  
**251538-68-8** 251538-69-9 251538-70-2 **251538-71-3**  
 251538-72-4 251538-73-5

RL: RCT (Reactant)

(**enantiomeric sepn.** of on (-)-phenylurea chiral  
 stationary phase)

RE.CNT 13

RE

- (1) Armstrong, D; J Chromatogr 1991, V539, P83 HCAPLUS
- (2) Gisch, D; The 9th International Symposium on Column Liquid Chromatography 1985
- (3) Keith, J; Chirality 1993, V5, P201
- (4) Lam, S; J Chromatogr Sci 1984, V22, P416 HCAPLUS
- (5) Okamoto, Y; Bull Chem Soc Jpn 1990, V63, P955 HCAPLUS
- (6) Okamoto, Y; Chem Lett 1987, P1857 HCAPLUS
- (7) Okamoto, Y; Chemtech 1988, P176 HCAPLUS
- (8) Okamoto, Y; J Chromatogr 1986, V363, P173 HCAPLUS
- (9) Pirkle, W; J Am Chem Soc 1987, V109, P5975 HCAPLUS
- (10) Pirkle, W; J Chromatogr 1984, V316, P585 HCAPLUS
- (11) Pirkle, W; J Org Chem 1992, V57, P3854 HCAPLUS
- (12) Ryu, J; Anal Sci 1997, V13, P217 HCAPLUS
- (13) Ryu, J; J Chromatogr A 1998, V14, P247


IT **251538-65-5 251538-68-8 251538-71-3**

RL: RCT (Reactant)

(**enantiomeric sepn.** of on (-)-phenylurea chiral  
 stationary phase)

RN 251538-65-5 HCAPLUS


CN Valine, N-[(1-naphthalenyloxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)




RN 251538-68-8 HCAPLUS

CN Phenylalanine, N-[(1-naphthalenyloxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)





RN 251538-71-3 HCAPLUS  
 CN Tryptophan, N-[(1-naphthalenyloxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)



L20 ANSWER 5 OF 12 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1999:35812 HCAPLUS  
 DN 130:153966  
 TI Solid-phase synthesis of tyrosine peptide aldehydes. Analogs of (S)-MAPI  
 AU Page, Patrick; Bradley, Mark; Walters, Iain; Teague, Simon  
 CS Department of Chemistry, University of Southampton, Southampton, SO17 1BJ, UK  
 SO J. Org. Chem. (1999), 64(3), 794-799  
 CODEN: JOCEAH; ISSN: 0022-3263  
 PB American Chemical Society  
 DT Journal  
 LA English  
 CC 34-3 (Amino Acids, Peptides, and Proteins)  
 Section cross-reference(s): 1  
 AB We report an efficient solid-phase synthesis of C-terminal tyrosine peptide aldehydes based on the HIV protease inhibitors (S)-MAPI and GE 20372 A. Our strategy consisted of anchoring the side chain of Dde-Tyrosinol onto the brominated Wang linker deriv. ((4-bromomethyl)-phenoxy-allyl acetate) to give after ester hydrolysis the N.alpha.-(Dde)-O-(4-methylphenoxyacetic acid)-L-Tyrosinol template. This was attached to aminomethyl resin and elongated using std. Fmoc protocols. Importantly there was no evidence of esterification side reactions. The unsym. substituted urea linkage of the (S)-MAPI family was incorporated using the N.alpha.-(4-nitrophenyloxycarbonyl)amino acid tert-Bu esters following which the protected tetrapeptide alc. immobilized on the solid support was oxidized to its corresponding aldehyde using sulfur trioxide-pyridine. The efficiency and reliability of the oxidn. step was dramatically improved by the incorporation of a small PEG-spacer between the linker and the solid support. The tetrapeptides were cleaved by acidolysis, purified by RP HPLC, and isolated in high yield and purity, demonstrating the success of the whole synthetic process.  
 ST solid phase synthesis tyrosine peptide aldehyde oxidn  
 IT Peptides, preparation  
 RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic

preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (aldehydes; solid-phase synthesis of tyrosine peptide aldehydes)

IT Oxidation  
 Solid phase synthesis  
 (solid-phase synthesis of tyrosine peptide aldehydes)

IT 70857-49-7P 163565-75-1P, GE 20372a  
 RL: BAC (Biological activity or effector, except adverse); PNU (Preparation, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (solid-phase synthesis of tyrosine peptide aldehydes)

IT 220237-27-4P 220237-28-5P  
 RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (solid-phase synthesis of tyrosine peptide aldehydes)

IT 14221-01-3  
 RL: CAT (Catalyst use); USES (Uses)  
 (solid-phase synthesis of tyrosine peptide aldehydes)

IT 693-13-0, Diisopropyl carbodiimide 7087-68-5, Diisopropyl ethylamine  
 7446-11-9, Sulfur trioxide, reactions 13887-98-4 57260-73-8  
 155505-56-9 187526-99-4 191425-55-5 191426-93-4  
 RL: RCT (Reactant)  
 (solid-phase synthesis of tyrosine peptide aldehydes)

IT 220237-29-6DP, resin-bound 220237-30-9P 220237-31-0P 220237-32-1P  
 220237-33-2P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (solid-phase synthesis of tyrosine peptide aldehydes)

RE.CNT 38

RE

- (1) Aoyagi, T; J Antibiot 1969, V22, P283 HCAPLUS
- (2) Aoyagi, T; Proteases and Biological Control 1975, P429 HCAPLUS
- (3) Bendal, M; Eur J Biochem 1977, V79, P201
- (4) Bloomberg, G; Tetrahedron Lett 1993, V34, P4709 HCAPLUS
- (5) Bullock, T; J Mol Biol 1996, V255, P714 HCAPLUS
- (6) Bycroft, B; J Chem Soc Chem Commun 1993, P778 HCAPLUS
- (7) Chen, C; J Am Chem Soc 1994, V116, P2661 HCAPLUS
- (8) Dinh, T; Tetrahedron Lett 1996, V37, P1161 HCAPLUS
- (9) Ede, N; Tetrahedron Lett 1997, V38, P7119 HCAPLUS
- (10) Fehrentz, J; Tetrahedron Lett 1995, V36, P7871 HCAPLUS
- (11) Frankfater, A; Biochemistry 1981, V20, P5517 HCAPLUS
- (12) Galeotti, N; Lett Pept Sci 1997, V4, P437 HCAPLUS
- (13) Kaneto, R; J Antibiot 1993, V46, P1622 HCAPLUS
- (14) Kennedy, W; Biochemistry 1979, V18, P349 HCAPLUS
- (15) Kunz, H; Angew Chem Int Ed Engl 1984, V23, P436
- (16) Mackenzie, N; Biochemistry 1986, V25, P2293 HCAPLUS
- (17) Malcolm, B; Biochemistry 1995, V34, P8172 HCAPLUS
- (18) Marsh, I; J Org Chem 1997, V62, P6199 HCAPLUS
- (19) Murphy, A; J Am Chem Soc 1992, V114, P3156 HCAPLUS
- (20) Nahm, S; Tetrahedron Lett 1981, V22, P3815 HCAPLUS
- (21) Ngu, K; Tetrahedron Lett 1997, V38, P973 HCAPLUS
- (22) Ogilvie, W; J Med Chem 1997, V40, P4113 HCAPLUS
- (23) Rano, T; Tetrahedron Lett 1995, V36, P3789 HCAPLUS
- (24) Richter, L; Tetrahedron Lett 1994, V35, P4705 HCAPLUS
- (25) Rotella, D; J Am Chem Soc 1996, V118, P12246 HCAPLUS
- (26) Sarin, V; Anal Biochem 1981, V20, P147
- (27) Sarubbi, E; FEBS Lett 1993, V319, P253 HCAPLUS
- (28) Schacht, A; Bioorg Med Chem Lett 1995, V5, P2529 HCAPLUS
- (29) Schroder, E; FEBS Lett 1993, V315, P38 HCAPLUS
- (30) Schultz, R; J Biol Chem 1989, V264, P1497 HCAPLUS

- (31) Stefanelli, S; J Antibiot 1995, V48, P332 HCAPLUS  
 (32) Stella, S; J Antibiot 1991, V44, P1019 HCAPLUS  
 (33) Thompson, R; Biochemistry 1973, V12, P47 HCAPLUS  
 (34) Watanbe, T; Agric Biol Chem 1979, V43, P243  
 (35) Weinshenker, N; Tetrahedron Lett 1972, V32, P3281  
 (36) Westerik, J; J Biol Chem 1972, V247, P8195 HCAPLUS  
 (37) Woo, J; Bioorg Med Chem Lett 1995, V5, P1501 HCAPLUS  
 (38) Zhang, X; J Org Chem 1997, V62, P6420 HCAPLUS

IT 191425-55-5 191426-93-4


RL: RCT (Reactant)

(solid-phase synthesis of tyrosine peptide aldehydes)

RN 191425-55-5 HCAPLUS

CN L-Phenylalanine, N-[(4-nitrophenoxy)carbonyl]-, 1,1-dimethylethyl ester  
 (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 191426-93-4 HCAPLUS

CN L-Tyrosine, O-(1,1-dimethylethyl)-N-[(4-nitrophenoxy)carbonyl]-,  
 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L20 ANSWER 6 OF 12 HCAPLUS COPYRIGHT 2001 ACS

AN 1997:598126 HCAPLUS

DN 127:176703

TI Solid-Phase Total Synthesis of Oscillamide Y and Analogs

AU Marsh, Ian R.; Bradley, Mark; Teague, Simon J.

CS Department of Chemistry, University of Southampton, Southampton, SO17 1BJ,  
 UK

SO J. Org. Chem. (1997), 62(18), 6199-6203

CODEN: JOCEAH; ISSN: 0022-3263


PB American Chemical Society

DT Journal

LA English

CC 34-3 (Amino Acids, Peptides, and  
 Proteins)


GI



I

- AB An efficient solid phase synthesis of oscillamide Y (I) and three analogs is reported. The cyclic peptide was prepd. using a combination of 9-fluorenylmethoxycarbonyl (Fmoc) and allyl chemistries and an acid labile Wang type linker. The urea functionality was smoothly incorporated using D-lysine building block 4-O<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>O<sub>2</sub>C-D-Lys(Fmoc)-OCH<sub>2</sub>CH:CH<sub>2</sub>. Coupling to the N-Me amino acid was readily achieved using HATU, monitoring the reaction using bromophenol blue. Allyl deprotection was accomplished using Pd(PPh<sub>3</sub>)<sub>4</sub> and dimedone, and cyclization was smoothly accomplished using PyBroP. All reactions were monitored using mass spectrometry methodol. The cyclized materials were cleaved by acidolysis and purified by RP HPLC. In all cases the material isolated was the major product and gave the expected mol. ion information. HPLC comparison with an authentic sample of oscillamide Y showed that the isomer contg. N-methyl-L-alanine and L-homotyrosine was the natural product. 1H NMR and 1H-1H COSY NMR expts. further confirmed this identification. The four compds. were tested as competitive and slow-tight binding inhibitors of chymotrypsin but showed, contrary to literature expectations, no inhibitory activity.
- ST oscillamide Y isomer solid phase synthesis
- IT Solid phase peptide synthesis  
(solid-phase total synthesis of oscillamide Y and analogs)
- IT 68858-21-9, 4-(Hydroxymethyl)phenoxyacetic acid  
RL: RCT (Reactant)  
(handle; solid-phase total synthesis of oscillamide Y and analogs)
- IT 34404-32-5, D-Lysine, N6-[(phenylmethoxy)carbonyl]-  
RL: RCT (Reactant)  
(solid-phase total synthesis of oscillamide Y and analogs)
- IT 55878-47-2P 115186-31-7P, Boc-D-Lys(Fmoc)-OH 193948-47-9P  
**193948-49-1P**  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(solid-phase total synthesis of oscillamide Y and analogs)
- IT 168482-80-2P, Oscillamide Y 194038-14-7P 194038-15-8P 194038-16-9P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(solid-phase total synthesis of oscillamide Y and analogs)
- IT **193948-49-1P**  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(solid-phase total synthesis of oscillamide Y and analogs)
- RN 193948-49-1 HCAPLUS
- CN L-Lysine, N6-[(9H-fluoren-9-ylmethoxy)carbonyl]-N2-[(4-nitrophenoxy)carbonyl]-, 2-propenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L20 ANSWER 7 OF 12 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1996:610346 HCAPLUS  
 DN 125:301598  
 TI A solid pharmaceutical composition providing improved oral bioavailability  
 for HIV protease inhibitors  
 IN Al-razzak, Laman A.; Marsh, Kennan C.; Pyter, Richard A.  
 PA Abbott Laboratories, USA  
 SO U.S., 18 pp. Cont.-in-part of U.S. Ser. No. 267,273, abandoned.  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 IC ICM A61K031-16  
 ICS A61K031-695; A61K031-425; A61K031-42  
 NCL 514616000  
 CC 34-2 (Amino Acids, Peptides, and  
 Proteins)  
 Section cross-reference(s): 63

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 5559158	A	19960924	US 1994-297004	19940831
	CA 2167413	AA	19950413	CA 1994-2167413	19940909
	WO 9509614	A1	19950413	WO 1994-US10096	19940909
	W: AU, CA, JP, KR				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	AU 9477229	A1	19950501	AU 1994-77229	19940909
	AU 685509	B2	19980122		
	EP 721330	A1	19960717	EP 1994-928043	19940909
	EP 721330	B1	20010328		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
	JP 09503501	T2	19970408	JP 1994-510810	19940909
	AT 200023	E	20010415	AT 1994-928043	19940909
	IL 110915	A1	19991222	IL 1994-110915	19940911
	US 5610193	A	19970311	US 1996-650261	19960522
PRAI	US 1993-130409	B2	19931001		
	US 1994-267273	B2	19940628		
	US 1994-297004	A	19940831		
	WO 1994-US10096	W	19940909		
	US 1995-424740	B1	19950418		
OS	MARPAT 125:301598				

GI



- AB A solid pharmaceutical compn. is claimed, comprising a pharmaceutically acceptable adsorbent or a mixt. of pharmaceutically acceptable adsorbents to which is adsorbed a mixt. of (1) a pharmaceutically acceptable org. solvent or a mixt. of pharmaceutically acceptable org. solvents, (2) HIV protease inhibitor I, and (3) a pharmaceutically acceptable acid or a combination of pharmaceutically acceptable acids. Thus, e.g., a capsule compn. contg. (% by wt.): (all-S)-I = (2S,3S,5S)-5-[N-[N-[[N-methyl-N-[(2-isopropyl-4-thiazolyl)methyl]amino]carbonyl]valinyl]amino]-2-[N-[(5-thiazolyl)methoxycarbonyl]amino]-1,6-diphenyl-3-hydroxyhexane (prepn. given) (21.84); propylene glycol, USP (10.96); Ethanol, dehydrated USP, 200 proof (22.99); polysorbate 80, NF (5.31); Cremophor EL (4.4); HCl, reagent grade (1.18); Cab-o-sil (26.88) exhibited 89.6 mean % oral bioavailability in dogs vs. <2.0 for unformulated (all-S)-I in capsules.
- ST oral pharmaceutical dosage HIV protease inhibitor
- IT Acquired immune deficiency syndrome  
(solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)
- IT Castor oil  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(ethoxylated, component of pharmaceutical formulation; solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)
- IT Pharmaceutical dosage forms  
(oral, solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)
- IT 50-81-7, Ascorbic acid, biological studies 57-55-6, Propylene glycol, biological studies 64-17-5, Ethanol, biological studies 104-15-4, p-Toluenesulfonic acid, biological studies 7631-86-9, Silicon dioxide, biological studies 7647-01-0, Hydrochloric acid, biological studies 9005-65-6, Polysorbate 80 14807-96-6, Talc, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(component of pharmaceutical formulation; solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)
- IT 9004-34-6, Cellulose, biological studies  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(microcryst., component of pharmaceutical formulation; solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)
- IT 162990-01-4P  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)
- IT 144114-21-6, Retropepsin  
RL: BSU (Biological study, unclassified); MSC (Miscellaneous); BIOL

(Biological study)

(solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)

IT 143838-10-2P 144164-10-3P

RL: BYP (Byproduct); PREP (Preparation)

(solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)

IT 75-12-7, Formamide, reactions 105-39-5, Ethyl chloroacetate 534-07-6, 1,3-Dichloroacetone 563-83-7, Isobutyramide 6306-52-1, L-Valine methyl ester hydrochloride 6372-14-1, N-(Benzyloxycarbonyl)-L-phenylalaninol 7693-46-1, 4-Nitrophenyl chloroformate 24424-99-5, Di-tert-butyl dicarbonate 153441-77-1 156732-13-7 156732-15-9

RL: RCT (Reactant)

(solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)

IT 115-08-2P, Thioformamide 13515-65-6P, Thioisobutyramide 32955-21-8P, 2-Amino-5-(ethoxycarbonyl)thiazole 32955-22-9P, Ethyl thiazole-5-carboxylate 33142-21-1P, Ethyl 2-chloro-2-formylacetate 38585-74-9P, 5-(Hydroxymethyl)thiazole 59830-60-3P, N-(Benzyloxycarbonyl)-L-phenylalaninal 65386-28-9P, 4-(Chloromethyl)-2-isopropylthiazole hydrochloride 137649-69-5P 144141-68-4P 144163-43-9P 144163-44-0P 144163-85-9P 144163-97-3P 144164-11-4P 154212-59-6P 154212-60-9P 154212-61-0P 154248-99-4P 162537-10-2P, N-(4-Nitrophenoxycarbonyl)-L-valine Methyl Ester 162849-92-5P 162849-93-6P 162849-94-7P 162849-95-8P 162849-96-9P 162990-03-6P 165315-39-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)

IT 155213-67-5P

RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)

IT 153441-77-1


RL: RCT (Reactant)

(solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)

RN 153441-77-1 HCAPLUS

CN L-Valine, N-(phenoxycarbonyl)-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 162537-10-2P, N-(4-Nitrophenoxycarbonyl)-L-valine Methyl Ester

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)


(solid pharmaceutical compn. providing improved oral bioavailability for HIV protease inhibitors)

RN 162537-10-2 HCAPLUS

CN L-Valine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX

NAME)


Absolute stereochemistry.



L20 ANSWER 8 OF 12 HCAPLUS COPYRIGHT 2001 ACS  
AN 1995:928117 HCAPLUS  
DN 123:340946  
TI Novel amino acid ester and reagent composition for detecting  
leukocytes or elastase in bodily fluids.  
IN Yagi, Yuji  
PA Kyoto Daiichi Kagaku Co., Ltd., Japan  
SO Eur. Pat. Appl., 19 pp.  
CODEN: EPXXDW  
DT Patent  
LA English  
IC ICM C07D295-12  
ICS C07D213-81; C07D213-82; C07D307-68; C07D209-36; C07D213-64;  
C07D311-46; C07C311-19; C07D309-32; C12Q001-44  
CC 34-2 (Amino Acids, Peptides, and  
Proteins)  
Section cross-reference(s): 7, 9, 80


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 661280	A1	19950705	EP 1994-120650	19941224
	R: CH, DE, FR, GB, IT, LI				
	JP 07233131	A2	19950905	JP 1994-319986	19941222
PRAI	JP 1993-349879		19931229		
OS	MARPAT 123:340946				
GI					



I





I

AB Amino acid esters of formula (XOA)<sub>n</sub>Y [X = arom. or heterocyclic moiety; A = L-amino acid; n .gtoreq. 2; Y = moiety derived from compd. with 2 or more carbonyl or sulfonyl groups, bound to N of amino acid groups A] are claimed. The esters are hydrolyzed by leukocytes or elastase with high specificity, and are thus useful for detecting these in bodily fluids. For example, reaction of ClCO(CH<sub>2</sub>)<sub>4</sub>COCl (adipoyl dichloride) with excess L-alanine and pyridine gave 87.5% (CH<sub>2</sub>)<sub>4</sub>[CO-Ala-OH]<sub>2</sub>. This was treated with SOCl<sub>2</sub> to give its acid dichloride (100%), which reacted with 3-(dimethylamino)-5-piperidinophenol [prepn. given] to give 65.8% title compd. I. A test paper based on I and 1-diazo-2-naphthol-4-sulfonic acid gave color change (red-purple) for leukocyte liq. or elastase, but no change for .alpha.-chymotrypsin, trypsin, cathepsin, or butylcholinesterase, and only slight change for subtilisin. Thirty syntheses and a variety of diagnostic examples are provided.

ST amino acid ester detn leukocyte elastase

IT Diagnosis

Leukocyte

Urine

(prepn. of amino acid esters as **reagents** for detecting leukocytes and elastase in bodily fluids)

IT Amino acids, preparation

RL: ARG (Analytical reagent use); SPN (Synthetic preparation); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); PREP (Preparation); USES (Uses)

(esters, prepn. of amino acid esters as **reagents** for detecting leukocytes and elastase in bodily fluids)

IT 480-93-3P, 3-Hydroxyindole 13170-66-6P, Cyclohexane-1,4-dicarbonyl chloride 40248-00-8P 136458-21-4P 141187-06-6P 170487-71-5P 170487-72-6P 170487-73-7P 170487-74-8P 170487-75-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation) (intermediate; prepn. of amino acid esters as **reagents** for detecting leukocytes and elastase in bodily fluids)

IT 9004-06-2, Elastase

RL: ANT (Analyte); ANST (Analytical study) (prepn. of amino acid esters as **reagents** for detecting leukocytes and elastase in bodily fluids)

IT 170487-41-9P 170487-42-0P 170487-43-1P 170487-44-2P 170487-45-3P 170487-46-4P 170487-47-5P **170487-48-6P** 170487-49-7P

170487-50-0P 170487-51-1P 170487-52-2P 170487-53-3P 170487-54-4P  
 170487-55-5P 170487-56-6P 170487-57-7P 170487-58-8P 170487-59-9P  
 170487-60-2P 170487-61-3P 170487-62-4P 170487-63-5P 170487-64-6P  
 170487-65-7P 170487-66-8P 170487-67-9P 170487-68-0P 170487-69-1P  
 170487-70-4P

RL: ARG (Analytical reagent use); SPN (Synthetic preparation); THU  
 (Therapeutic use); ANST (Analytical study); BIOL (Biological study); PREP  
 (Preparation); USES (Uses)

(prepn. of amino acid esters as **reagents** for detecting  
 leukocytes and elastase in bodily fluids)

IT 56-40-6, Glycine, reactions 56-41-7, L-Alanine, reactions 61-90-5,  
 L-Leucine, reactions 72-18-4, L-Valine, reactions 83-72-7,  
 2-Hydroxy-1,4-naphthoquinone 89-05-4, Benzene-1,2,4,5-tetracarboxylic  
 acid 89-32-7 99-14-9, Propane-1,2,3-tricarboxylic acid 100-20-9,  
 1,4-Benzenedicarbonyl dichloride 108-73-6, 1,3,5-Trihydroxybenzene  
 110-89-4, Piperidine, reactions 111-50-2, Adipoyl chloride 119-80-2,  
 2,2'-Dithiodibenzoic acid 124-40-3, reactions 142-08-5,  
 2-Hydroxypyridine 533-75-5, Tropolone 585-47-7, Benzene-1,3-disulfonyl  
 chloride 608-08-2, Indoxyl acetate 636-78-2, 4-Sulfobenzoic acid  
 1076-38-6, 4-Hydroxycoumarin 1076-97-7, 1,4-Cyclohexanedicarboxylic acid  
 1141-38-4, Naphthalene-2,6-dicarboxylic acid 3119-64-0,  
 Diphenylmethane-4,4'-disulfonyl chloride 3387-26-6, Furan-3,4-  
 dicarboxylic acid 3739-94-4, Pyridine-2,6-dicarbonyl chloride  
 4808-48-4 13827-62-8, Naphthalene-2,6-disulfonyl chloride 15658-35-2,  
 6,6'-Dithiodinicotinic acid 33177-29-6 67294-61-5,  
 1,3,6-Naphthalenetrisulfonyl chloride 170487-76-0 170487-77-1  
 170487-78-2 170487-79-3 170487-80-6

RL: RCT (Reactant)

(starting material; prepn. of amino acid esters as **reagents**  
 for detecting leukocytes and elastase in bodily fluids)

IT **170487-48-6P**



RL: ARG (Analytical reagent use); SPN (Synthetic preparation); THU  
 (Therapeutic use); ANST (Analytical study); BIOL (Biological study); PREP  
 (Preparation); USES (Uses)

(prepn. of amino acid esters as **reagents** for detecting  
 leukocytes and elastase in bodily fluids)

RN 170487-48-6 HCAPLUS

CN L-Alanine, N,N'-[1,4-phenylenebis(oxycarbonyl)]bis-, bis[3-(dimethylamino)-  
 5-(1-piperidinyl)phenyl] ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



L20 ANSWER 9 OF 12 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1995:887807 HCAPLUS  
 DN 123:314522  
 TI Pharmaceutical composition for HIV protease inhibitor [ritonavir] with improved oral bioavailability  
 IN Al-razzak, Laman A.; Marsh, Kennan C.; Pyter, Richard A.  
 PA Abbott Laboratories, USA  
 SO PCT Int. Appl., 55 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM A61K009-14  
 ICS A61K009-16; A61K009-48  
 CC 34-2 (Amino Acids, Peptides, and Proteins)  
 Section cross-reference(s): 28, 63  
 FAN.CNT 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----

PI WO 9509614 A1 19950413 WO 1994-US10096 19940909  
 W: AU, CA, JP, KR  
 RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE  
 US 5559158 A 19960924 US 1994-297004 19940831  
 AU 9477229 A1 19950501 AU 1994-77229 19940909  
 AU 685509 B2 19980122  
 EP 721330 A1 19960717 EP 1994-928043 19940909  
 EP 721330 B1 20010328  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE  
 JP 09503501 T2 19970408 JP 1994-510810 19940909  
 AT 200023 E 20010415 AT 1994-928043 19940909  
 PRAI US 1993-130409 A 19931001  
 US 1994-267273 A 19940628  
 US 1994-297004 A 19940831  
 WO 1994-US10096 W 19940909  
 OS MARPAT 123:314522  
 GI



AB A solid pharmaceutical compn. is disclosed which comprises a pharmaceutically acceptable adsorbent or mixt. of adsorbents, to which is adsorbed a mixt. of: (1) a pharmaceutically acceptable org. solvent or mixt. of solvents; (2) an HIV protease-inhibiting compd.; and (3) one or more pharmaceutically acceptable acids. The solid compn. can optionally be encapsulated in a hard gelatin capsule. The compn. is particularly applicable to compd. I, and esp. its (2S,3S,5S,L)-isomer [ritonavir; II]. For example, oral administration of unformulated II to dogs gave < 2.0% mean bioavailability. In contrast, 89.6% mean bioavailability was obtained with the following capsule formulation: II 21.84, propylene glycol 10.96, ethanol 22.99, Polysorbate 80 5.31, Cremophor EL 4.4, HCl 1.18, and Cab-o-sil 26.88% by wt. Also described are addnl. oral formulations (comparative and invention), and several syntheses of II. For example, N-(benzyloxycarbonyl)-L-phenylalaninol was converted in 5 steps to (2S,3S,5S)-PhCH<sub>2</sub>CH(NHZ)CH(OH)CH<sub>2</sub>CH(NHZ)CH<sub>2</sub>Ph [Z = benzyloxycarbonyl], which was deprotected and reacted with 5-thiazolylmethyl nitrophenyl carbonate to give intermediate III and its isomer from acylation of the other amino group. Coupling of III with N-[[N-methyl-N-[(2-isopropyl-4-thiazolyl)methyl]amino]carbonyl]-L-valine

[prepn. given] using the carbodiimide reagent EDC and 1-hydroxybenzotriazole gave II.

ST HIV protease inhibitor ritonavir pharmaceutical bioavailability; oral bioavailability HIV protease inhibitor

IT Drug bioavailability  
Virucides and Virustats  
(pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)

IT Castor oil  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(ethoxylated, Cremophor EL, adjuvant; pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)

IT Pharmaceutical dosage forms  
(oral, pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)

IT 144114-21-6, Retropepsin  
RL: BSU (Biological study, unclassified); MSC (Miscellaneous); BIOL (Biological study)  
(HIV-1 and HIV-2; pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)

IT 50-81-7, Ascorbic acid, biological studies 57-55-6, Propylene glycol, biological studies 64-17-5, Ethanol, biological studies 77-92-9, Citric acid, biological studies 104-15-4, p-Toluenesulfonic acid, biological studies 7631-86-9, Silica, biological studies 7647-01-0, Hydrochloric acid, biological studies 9002-96-4 9004-34-6, Cellulose, biological studies 9005-25-8, Corn starch, biological studies 9005-65-6, Polysorbate 80 9050-36-6, Maltodextrin 14807-96-6, Talc, biological studies 41080-67-5  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(adjuvant; pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)

IT 115-08-2P, Thioformamide 13515-65-6P, 2-Methylpropanethioamide 32955-21-8P, 2-Amino-5-(ethoxycarbonyl)thiazole 32955-22-9P, Ethyl thiazole-5-carboxylate 33142-21-1P, Ethyl 2-chloro-2-formylacetate 38585-74-9P, 5-(Hydroxymethyl)thiazole 59830-60-3P, N-[[ (Benzyl)oxy]carbonyl]-L-phenylalaninal 65386-28-9P, 4-(Chloromethyl)-2-isopropylthiazole hydrochloride 137649-69-5P 143838-10-2P 144141-68-4P 144163-43-9P 144163-44-0P 144163-85-9P 144163-97-3P 144164-10-3P 144164-11-4P 154212-59-6P 154212-60-9P, 2-Isopropyl-4-[(N-methylamino)methyl]thiazole 154212-61-0P 154248-99-4P 162537-10-2P 162849-92-5P 162849-93-6P 162849-94-7P 162849-95-8P 162849-96-9P 162990-03-6P 165315-39-9P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(intermediate; pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)

IT 155213-67-5P 162990-01-4P  
RL: BPR (Biological process); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); PROC (Process); USES (Uses)  
(pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)

IT 62-56-6, Thiourea, reactions 74-89-5, Methylamine, reactions 75-12-7, Formamide, reactions 98-80-6, Phenylboronic acid 105-39-5, Ethyl chloroacetate 109-94-4, Ethyl formate 1534-07-6, 1,3-Dichloroacetone 563-83-7, Isobutyramide 6306-52-1, L-Valine methyl ester hydrochloride 6372-14-1, N-[[ (Benzyl)oxy]carbonyl]-L-phenylalaninol 7693-46-1, 4-Nitrophenyl chloroformate 24424-99-5, Di-tert-butyl dicarbonate 40635-67-4, .alpha.-Acetoxyisobutyryl bromide 126147-70-4, N-(Phenoxycarbonyl)-L-valine 156732-13-7 156732-15-9 169597-13-1  
RL: RCT (Reactant)

(starting material; pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)


IT 162537-10-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(intermediate; pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)

RN 162537-10-2 HCAPLUS

CN L-Valine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 126147-70-4, N-(Phenoxycarbonyl)-L-valine


RL: RCT (Reactant)

(starting material; pharmaceutical compn. of HIV protease inhibitor with improved oral bioavailability)

RN 126147-70-4 HCAPLUS

CN L-Valine, N-(phenoxycarbonyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L20 ANSWER 10 OF 12 HCAPLUS COPYRIGHT 2001 ACS

AN 1990:179761 HCAPLUS

DN 112:179761

TI Optical resolution of amino acid derivatives by high-performance liquid chromatography on tris(phenylcarbamate)s of cellulose and amylose

AU Okamoto, Yoshio; Kaida, Yuriko; Aburatani, Ryo; Hatada, Koichi

CS Fac. Eng. Sci., Osaka Univ., Toyonaka, 560, Japan

SO J. Chromatogr. (1989), 477(2), 367-76

CODEN: JOCRAM; ISSN: 0021-9673

DT Journal

LA English

CC 34-2 (Amino Acids, Peptides, and Proteins)

AB The optical resolu. of 10 N-protected alanine esters was examd. by HPLC using 6 cellulose and 5 amylose tris(phenylcarbamate) derivs. as chiral stationary phases. Tris(3,5-dimethylphenylcarbamate)s of both cellulose and amylose showed high resolving power for these racemates. The resolu. of 23 N-benzyloxycarbonyl .alpha.-amino acid ester was also tested on tris(3,5-dimethylphenylcarbamate)s of cellulose and amylose. All but 2 amino acid derivs. were completely resolved at least by one of the columns. On cellulose tris(3,5-dimethylphenylcarbamate), all L-amino

acids (except threonine) eluted first.

ST resoln amino ester **HPLC** cellulose trisphenylcarbamate; amylose trisphenylcarbamate resoln amino ester

IT Resolution  
(of protected amino acid esters by **HPLC** on cellulose and amylose tris(phenylcabamate) stationary phases)

IT Chromatography, column and liquid  
(high-performance, resoln. by, of protected amino acid esters with cellulose and amylose tris(phenylcarbmate) stationary phases)

IT Amino acids, esters  
RL: PROC (Process)  
(N-protected, esters, resoln. of, by **HPLC** with cellulose and amylose tris(phenylcarbamate) stationary phases)


IT 4515-20-2 5143-72-6 5446-46-8, N-Benzoyl-DL-alanine ethyl ester  
5513-39-3, N-Benzyloxycarbonyl-DL-alanine benzyl ester 5557-84-6  
23161-27-5, N-Benzyloxycarbonyl-DL-serine ethyl ester 23161-28-6,  
N-Benzyloxycarbonyl-DL-threonine ethyl ester 25282-53-5,  
DL-Phenylalanine benzyl ester 39978-35-3, N-Benzyloxycarbonyl-DL-leucine ethyl ester 40489-45-0, N-Benzyloxycarbonyl-DL-serine benzyl ester 42998-42-5, N-Benzyloxycarbonyl-DL-valine ethyl ester 42998-44-7, N-Benzyloxycarbonyl-DL-phenylalanine ethyl ester 42998-45-8  
46229-47-4, DL-Alanine benzyl ester 72604-32-1, N-tert-Butoxycarbonyl-DL-alanine ethyl ester 72604-33-2, N-Benzyloxycarbonyl-DL-alanine ethyl ester 85369-24-0 86827-19-2 **88406-41-1** 103063-37-2,  
N-Benzyloxycarbonyl-DL-phenylalanine benzyl ester 114285-13-1,  
N-Acetyl-DL-alanine benzyl ester 126400-85-9 126400-86-0 126400-87-1  
126400-88-2 126400-89-3 126400-90-6 126400-91-7 126400-92-8  
126400-93-9 126400-94-0, N-Benzoyl-DL-alanine benzyl ester 126400-95-1  
**126400-96-2** 126400-97-3, N-Benzyloxycarbonyl-DL-valine benzyl ester 126400-98-4, N-Benzyloxycarbonyl-DL-norvaline ethyl ester 126400-99-5, N-Benzyloxycarbonyl-DL-norvaline benzyl ester 126401-00-1, N-Benzyloxycarbonyl-DL-leucine benzyl ester 126401-01-2, N-Benzyloxycarbonyl-DL-norleucine ethyl ester 126401-02-3, N-Benzyloxycarbonyl-DL-norleucine benzyl ester 126401-03-4, N-Benzyloxycarbonyl-DL-isoleucine benzyl ester 126401-04-5 126401-05-6  
126401-06-7, N-Benzyloxycarbonyl-DL-threonine benzyl ester 126401-07-8  
126401-08-9 126401-09-0 126401-10-3 126401-11-4 126401-12-5  
126401-13-6 126401-14-7 126401-15-8 126401-16-9 126401-17-0  
126401-18-1 126401-19-2 126401-20-5 126401-21-6,  
N-Benzyloxycarbonyl-DL-methionine benzyl ester 126401-22-7 126401-23-8  
126433-60-1, N-Benzyloxycarbonyl-DL-methionine ethyl ester 126456-11-9  
126456-12-0 126456-13-1, N-Benzyloxycarbonyl-DL-proline ethyl ester 126456-14-2, N-Benzyloxycarbonyl-DL-proline benzyl ester  
RL: PROC (Process)  
(resoln. of, by **HPLC** with cellulose and amylose tris(phenylcarbamate) stationary phases)

IT 9047-05-6, Amylose tris(phenylcarbamate) 9047-07-8, Cellulose tris(phenylcarbamate) 103938-40-5, Cellulose tris(4-bromophenylcarbamate) 103938-45-0, Cellulose tris(4-ethylphenylcarbamate) 103938-46-1, Cellulose tris(4-fluorophenylcarbamate) 103938-49-4 107028-63-7, Amylose tris(3,5-dichlorophenylcarbamate) 115901-94-5, Amylose tris(4-chlorophenylcarbamate) 115901-96-7, Amylose tris(4-methylphenylcarbamate)


RL: RCT (Reactant)  
(stationary phase, for **HPLC** resoln. of protected alanine esters)

IT 103938-44-9, Cellulose tris(3,5-dimethylphenylcarbamate) 112049-40-8  
RL: RCT (Reactant)  
(stationary phase, for **HPLC** resoln. of protected amino acid

esters)  
 IT 88406-41-1 126400-96-2  
 RL: PROC (Process)  
 (resoln. of, by HPLC with cellulose and amylose  
 tris(phenylcarbamate) stationary phases)  
 RN 88406-41-1 HCAPLUS  
 CN Alanine, N-(phenoxy-carbonyl)-, ethyl ester (9CI) (CA INDEX NAME)



RN 126400-96-2 HCAPLUS  
 CN Alanine, N-(phenoxy-carbonyl)-, phenylmethyl ester (9CI) (CA INDEX NAME)



L20 ANSWER 11 OF 12 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1988:611486 HCAPLUS  
 DN 109:211486  
 TI Preparation of heterocyclipeptides as renin inhibitors  
 IN Ryono, Denis Evan; Weller, Harold Norris, III  
 PA Squibb, E. R., and Sons, Inc., USA  
 SO Eur. Pat. Appl., 143 pp.  
 CODEN: EPXXDW  
 DT Patent  
 LA English  
 IC ICM C07K005-06  
 ICS C07K005-08; C07D233-64; C07D417-12; C07D401-12; C07D403-12;  
 A61K037-02; A61K037-64; A61K031-415; A61K031-425; A61K031-44  
 ICI C07D417-12, C07D277-00, C07D233-00; C07D401-12, C07D213-00, C07D233-00;  
 C07D417-12, C07D235-00, C07D233-00  
 CC 34-3 (Amino Acids, Peptides, and  
 Proteins)  
 Section cross-reference(s): 1

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 231919	A2	19870812	EP 1987-101373	19870202
EP 231919	A3	19900718		
EP 231919	B1	19930120		
R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, LU, NL, SE				
ZA 8700563	A	19870930	ZA 1987-563	19870126
AU 8768153	A1	19870806	AU 1987-68153	19870130
AU 595578	B2	19900405		
DK 8700523	A	19870804	DK 1987-523	19870202
HU 44576	A2	19880328	HU 1987-352	19870202
HU 203368	B	19910729		
AT 84793	E	19930215	AT 1987-101373	19870202



ES 2043611	T3	19940101	ES 1987-101373	19870202
JP 62258365	A2	19871110	JP 1987-23434	19870203
CA 1310792	A1	19921124	CA 1987-528852	19870203
US 4885292	A	19891205	US 1989-373633	19890629
PRAI US 1986-825724		19860203		
US 1987-3446		19870115		
EP 1987-101373		19870202		
AB	<p>X-(NHCHR5CO)p-NHCHR4CONHCHR3CH(OH)R1 [I; X = R6(CH2)n, R6(CH2)nO2C, R6O(CH2)nCO, R6(CH2)nSO2, etc.; R1 = (substituted) N-contg. heterocyclyl; R3, R4, R5 = (halo)alkyl, arylalkyl, hydroxyalkyl, carboxyalkyl, guanidinoalkyl, imidazolylalkyl, etc.; R6 = alkyl, cycloalkyl, aryl, heterocyclyl; n = 0-5; p = 0, 1] useful as renin inhibitors, (no data) were prepd. N-[(1,1-Dimethylethoxy)carbonyl]-L-leucinal (prepn. from leucine given) in THF was added to a mixt. of BuLi and 1-[(phenylmethoxy)methyl]-1H-imidazole in THF at -70.degree.. The mixt. was kept at -70.degree. for 1 h, and at 0.degree. for 15 min followed by quenching with aq. NH4Cl and hydrolysis with HCl/EtOAc to give .alpha.-[(S)-1-amino-3-methylbutyl]-1-[(phenylmethoxy)methyl]-1H-imidazole-2-methanol-2HCl. The latter was coupled with N-[N-[(1,1-dimethylethoxy)carbonyl]-L-phenylalanyl]-1-[(phenylmethoxy)methyl]-L-histidine (prepn. given) using 1-hydroxybenzotriazole and DCC in DMF at ice temps. The coupling product was hydrogenolyzed in aq. MeOH contg. HCl over Pd/C to give N2-[N-[(1,1-dimethylethoxy)carbonyl]-L-phenylalanyl]-N-[(S)-1-[hydroxy(1H-imidazol-2-yl)methyl]-3-methylbutyl]-L-histidinamide-HOAc.</p>			
ST	<p>imidazolylpeptide prepn antihypertensive; Renin inhibitor peptide heterocyclyl; antihypertensive heterocyclylpeptide</p>			
IT	<p>Antihypertensives (heterocyclyl-contg. peptide alcs.)</p>			
IT	<p>Peptides, preparation RL: SPN (Synthetic preparation); PREP (Preparation) (heterocyclyl-contg., prepn. of, as renin inhibitors)</p>			
IT	<p>7693-46-1, 4-Nitrophenyl chloroformate RL: RCT (Reactant) (acylation by, of phenylalanine Me ester, in prepn. of antihypertensive)</p>			
IT	<p>123-75-1, Pyrrolidine, reactions RL: RCT (Reactant) (acylation of, by phosgene, in prepn. of antihypertensive)</p>			
IT	<p>100-39-0, Benzyl bromide RL: RCT (Reactant) (alkylation by, of di-Et benzylmalonate, in prepn. of antihypertensive)</p>			
IT	<p>607-81-8, Diethyl benzylmalonate RL: RCT (Reactant) (alkylation of, by benzyl bromide)</p>			
IT	<p>105-53-3, Diethyl malonate RL: RCT (Reactant) (alkylation of, by chloromethylnaphthalene, in prepn. of antihypertensive)</p>			
IT	<p>6638-79-5, O,N-Dimethylhydroxylamine hydrochloride RL: RCT (Reactant) (amidation by, of (dimethylethoxycarbonylamino)cyclohexanepropanoic acid, in prepn. of antihypertensive)</p>			
IT	<p>7524-50-7, L-Phenylalanine methyl ester hydrochloride RL: RCT (Reactant) (amidation by, of pyrrolidinecarbonyl chloride, in prepn. of antihypertensive)</p>			
IT	<p>13139-15-6 RL: RCT (Reactant) (amidation of, by dimethylhydroxylamine, in prepn. of antihypertensive)</p>			

IT 86-52-2, 1-Chloromethylnaphthalene  
 RL: RCT (Reactant)  
 (condensation of, with di-Et malonate, in prepn. of antihypertensive)

IT 61-90-5, L-Leucine, reactions  
 RL: RCT (Reactant)  
 (conversion of, to Et ester)

IT 71-00-1, L-Histidine, reactions  
 RL: PROC (Process)  
 (conversion of, to Me ester, in prepn. of antihypertensive)

IT 115888-02-3P  
 RL: SPN (Synthetic preparation); FORM (Formation, nonpreparative); PREP (Preparation)  
 (formation of, in prepn. of antihypertensive)

IT 288-47-1, Thiazole 49822-58-4  
 RL: RCT (Reactant)  
 (lithiation and condensation of, with leucinal deriv., in prepn. of antihypertensive)

IT 623-33-6, Glycine ethyl ester monohydrochloride 2666-93-5, L-Leucine methyl ester 13734-34-4 25024-53-7 25616-33-5 33014-68-5 83468-83-1 96867-10-6  
 RL: RCT (Reactant)  
 (peptide coupling of, in prepn. of antihypertensive)

IT 7389-87-9P, L-Histidine methyl ester dihydrochloride  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and N-protection of, in prepn. of antihypertensive)

IT 111629-38-0P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and coupling of, with phenylalanine deriv., in prepn. of antihypertensive)

IT 4372-32-1P 115766-40-0P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and decarboxylation of, in prepn. of antihypertensive)

IT 115766-26-2P 115766-27-3P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and deprotection of, in prepn. of antihypertensive)

IT 82010-31-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and oxidn. of, in prepn. of antihypertensive)

IT 115766-28-4P 115766-29-5P 115766-31-9P 115766-34-2P 115766-35-3P 115766-37-5P 115766-38-6P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and peptide coupling of, in prepn. of antihypertensive)

IT 7533-40-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and protection of, in prepn. of antihypertensive)

IT 20898-43-5P 111629-37-9P 115766-33-1P 115766-41-1P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and reaction of, in prepn. of antihypertensive)

IT 2743-40-0P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and redn. of)

IT 51987-73-6P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and redn. of, in prepn. of antihypertensive)

IT 111629-39-1P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and sapon. of, in prepn. of antihypertensive)

IT 115765-65-6P 115765-66-7P 115765-67-8P 115765-68-9P 115765-69-0P 115765-70-3P 115765-71-4P 115765-72-5P 115765-73-6P 115765-74-7P 115765-76-9P 115765-77-0P 115765-78-1P 115765-79-2P 115765-80-5P

115765-81-6P 115765-82-7P 115765-83-8P 115765-84-9P 115765-86-1P  
 115765-88-3P 115766-42-2P 115766-43-3P 115802-91-0P 115887-98-4P  
 115888-03-4P 115888-04-5P 115888-05-6P 115888-06-7P 115888-07-8P  
 115888-08-9P 115888-09-0P 115888-86-3P 115935-94-9P 115935-95-0P  
 115935-96-1P

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (prepn. of, as antihypertensive)

IT 597-55-7P 618-68-8P 2481-59-6P 30189-51-6P 37736-82-6P  
**54601-21-7P** 58521-45-2P 64152-76-7P 66605-57-0P  
 72155-45-4P 87694-50-6P 98105-42-1P 102639-04-3P 102639-05-4P  
 103322-56-1P 104539-18-6P 110695-91-5P 114457-62-4P 114473-20-0P  
 115765-89-4P 115765-90-7P 115765-91-8P 115765-92-9P 115765-93-0P  
 115765-94-1P 115765-95-2P 115765-96-3P 115765-97-4P 115765-98-5P  
 115765-99-6P 115766-00-2P 115766-01-3P 115766-02-4P 115766-03-5P  
 115766-04-6P 115766-05-7P 115766-06-8P 115766-07-9P 115766-08-0P  
 115766-09-1P 115766-10-4P 115766-11-5P 115766-12-6P 115766-13-7P  
 115766-14-8P 115766-15-9P 115766-16-0P 115766-17-1P 115766-18-2P  
 115766-19-3P 115766-20-6P 115766-22-8P 115766-23-9P 115766-24-0P  
 115766-25-1P 115766-32-0P 115766-36-4P 115766-39-7P 115802-92-1P  
 115802-93-2P 115802-94-3P 115802-95-4P 115887-99-5P 115888-00-1P  
 115888-01-2P

RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of, as intermediate for antihypertensive)

IT 115766-30-8P

RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of, as intermediates for antihypertensive)

IT 1070-83-3, 3,3-Dimethylbutanoic acid

RL: RCT (Reactant)  
 (reaction of, in prepn. of antihypertensive peptide deriv.)

IT 109-01-3, 1-Methylpiperazine 110-91-8, Morpholine, reactions

RL: RCT (Reactant)  
 (reaction of, with (nitrophenoxycarbonyl)phenylalanine)

IT 70-34-8, 2,4-Dinitrofluorobenzene

RL: RCT (Reactant)  
 (reaction of, with histidine contg. dipeptide, in prepn. of antihypertensive)

IT 68-11-1, Mercaptoacetic acid, uses and miscellaneous

RL: USES (Uses)  
 (reagent, for deprotection of dinitrophenylhistidine contg. peptide)

IT 3674-06-4

RL: RCT (Reactant)  
 (redn. of, in prepn. of antihypertensive)


IT **54601-21-7P**


RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of, as intermediate for antihypertensive)

RN 54601-21-7 HCAPLUS

CN L-Phenylalanine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.





L20 ANSWER 12 OF 12 HCAPLUS COPYRIGHT 2001 ACS

AN 1974:437818 HCAPLUS

DN 81:37818

TI Histidine derivatives

PA Tanabe Seiyaku Co., Ltd.

SO Brit., 9 pp.

CODEN: BRXXAA

DT Patent

LA English

IC C07D; A61K


CC 34-3 (Synthesis of Amino Acids, Peptides, and Proteins)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	GB 1352414	A	19740508	GB 1971-12019	19710428
GI	For diagram(s), see printed CA Issue.				
AB	L-Pyroglutamyl-L-proline amide (I), a thyrotropine-releasing or thyroid gland-stimulating factor, was prepd. from L-histidine Me ester (II). III-IV were prepd. from II by successive treatment with the appropriate chloride or azidoformate, hydrolysis, and tosylation. Condensation of these with L-proline amide at -10 to 10.degree. in the presence of a carbodiimide reagent, followed by treatment with N-benzyloxycarbonyl-L-pyroglutamic acid, and removal of the protecting groups, gave I.				
ST	histidine deriv				
IT	Protective groups (alkoxycarbonyl and sulfenyl groups, for amino group of histidine methyl ester, in peptide prepn.)				
IT	Peptides, preparation RL: PREP (Preparation) (histidine- and proline-contg.)				
IT	(o-Nitrophenyl)sulfenyl group (tert-Pentyloxy)carbonyl group [(p-Methoxybenzyl)oxy]carbonyl group tert-Butoxycarbonyl group (protective group, for amino group in peptide prepn.)				
IT	Amino group (protective groups for, alkoxycarbonyl and sulfenyl groups as, in peptide prepn.)				
IT	Imido group (protective groups for, benzyl and tert-alkyloxycarbonyl groups as, in peptide prepn.)				
IT	(Benzyloxy)carbonyl group (protective groups, for imido group in peptide prepn.)				
IT	7531-52-4 RL: RCT (Reactant) (condensation of, with histidines)				
IT	15354-08-2P	23241-50-1P	23241-51-2P	24305-27-9P	35899-43-5P
	35899-44-6P	35899-45-7P	35899-46-8P	35899-47-9P	35899-49-1P


53090-17-8P 53090-19-0P 53090-21-4P 53090-24-7P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 IT 98-79-3  
 RL: RCT (Reactant)  
 (protective group for imido group of, in peptide prepn.)  
 IT 7389-87-9  
 RL: RCT (Reactant)  
 (protective groups for .alpha.-amino group of, in peptide prepn.)  
 IT 5591-72-0 7669-54-7 24608-52-4 40438-33-3  
 RL: RCT (Reactant)  
 (reaction of, with histidine methyl ester)  
 IT 32159-21-0 53090-23-6  
 RL: RCT (Reactant)  
 (reaction of, with histidyl prolinamide)  
 IT 53090-17-8P 53090-19-0P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 53090-17-8 HCAPLUS  
 CN L-Histidine, N-[(4-methoxyphenoxy)carbonyl]-, methyl ester (9CI) (CA  
 INDEX NAME)

Absolute stereochemistry.




RN 53090-19-0 HCAPLUS  
 CN L-Histidine, N-[(4-methoxyphenoxy)carbonyl]-1-[(4-methylphenyl)sulfonyl]-,  
 compd. with N-cyclohexylcyclohexanamine (1:1) (9CI) (CA INDEX NAME)  
 CM 1  
 CRN 53090-18-9  
 CMF C21 H21 N3 O7 S  
 CDES 5:L

Absolute stereochemistry.




CM 2

CRN 101-83-7  
CMF C12 H23 N



L1

STR



## NODE ATTRIBUTES:

HCOUNT	IS	M1	AT	10
NSPEC	IS	R	AT	1
NSPEC	IS	R	AT	2
NSPEC	IS	R	AT	3
NSPEC	IS	R	AT	4
NSPEC	IS	R	AT	5
NSPEC	IS	R	AT	6
NSPEC	IS	C	AT	7
NSPEC	IS	C	AT	8
NSPEC	IS	C	AT	9
NSPEC	IS	C	AT	10
NSPEC	IS	C	AT	11
NSPEC	IS	C	AT	12
NSPEC	IS	C	AT	13
NSPEC	IS	C	AT	14

DEFAULT MLEVEL IS ATOM

MLEVEL IS CLASS AT 7 8 9 10 11 12 13 14

DEFAULT ECLEVEL IS LIMITED


## GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 14

## STEREO ATTRIBUTES: NONE

L3	501	SEA FILE=REGISTRY SSS FUL L1
L4	213	SEA FILE=HCAPLUS ABB=ON PLU=ON L3
L6	1	SEA FILE=HCAPLUS ABB=ON PLU=ON 2000:755247/AN
L14	75483	SEA FILE=HCAPLUS ABB=ON PLU=ON (AMINO ACIDS OR PEPTIDES OR PROTEINS)/CC
L15	83	SEA FILE=HCAPLUS ABB=ON PLU=ON L4 AND L14
L16	82	SEA FILE=HCAPLUS ABB=ON PLU=ON L15 NOT L6
L18	4	SEA FILE=HCAPLUS ABB=ON PLU=ON (HPLC OR CHROMATOGARPH?) AND L16
L19	9	SEA FILE=HCAPLUS ABB=ON PLU=ON (ENANTIO? OR SEPARAT? OR CHROMOPHO? OR REAGENT) AND L16
L20	12	SEA FILE=HCAPLUS ABB=ON PLU=ON L18 OR L19
L21		STR



NODE ATTRIBUTES:

HCOUNT	IS M1	AT	10
NSPEC	IS R	AT	1
NSPEC	IS R	AT	2
NSPEC	IS R	AT	3
NSPEC	IS R	AT	4
NSPEC	IS R	AT	5
NSPEC	IS R	AT	6
NSPEC	IS C	AT	7
NSPEC	IS C	AT	8
NSPEC	IS C	AT	9
NSPEC	IS C	AT	10
NSPEC	IS C	AT	11
NSPEC	IS C	AT	12
NSPEC	IS C	AT	13
NSPEC	IS C	AT	14
NSPEC	IS C	AT	15
NSPEC	IS C	AT	16
NSPEC	IS C	AT	17

DEFAULT MLEVEL IS ATOM

MLEVEL IS CLASS AT 7 8 9 10 11 12 13 14 15 16 17

DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED

NUMBER OF NODES IS 17

STEREO ATTRIBUTES: NONE

L23	87	SEA	FILE=REGISTRY	SUB=L3	SSS	FUL	L21
L24	56	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L23	
L25	50	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L24	NOT L20
L26	49	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L25	NOT L6
L27	24	SEA	FILE=HCAPLUS	ABB=ON	PLU=ON	L14	AND L26





L27 ANSWER 1 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 2001:416903 HCAPLUS  
 DN 135:33643  
 TI Preparation of 3-(2-aminoethylthio)methyl-4-oxo-4-(3-pyridyl)butanoic acid derivatives as neuroprotective agents  
 IN Bhagwat, Shripad; Palanki, Moorthy; Erdman, Paul; Doubleday, Mary; Sato, Hiroshi  
 PA Nippon Kayaku Co., Ltd., Japan  
 SO PCT Int. Appl., 119 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM C07D213-55  
 ICS C07D213-56; C07D417-12; A61K031-44; A61P025-00  
 CC 34-2 (Amino Acids, Peptides, and Proteins)  
 Section cross-reference(s): 1, 27

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001040187	A2	20010607	WO 2000-JP8090	20001116
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRAI US 1999-450245	A	19991129		

GI



AB The title compds. [I; R1 = (un)substituted alkyl, aryl, arylakyl, etc.; R2 = OR2a, NR2bR2c; R3 = H, :O, CO2R3a, etc.; R4 = H, (un)substituted alkyl, aryl, etc.; R3 and R4 taken together = (un)substituted heterocyclyl; R5 = H, (un)substituted alkyl; R2a = H, (un)substituted alkyl, aryl, etc.; R2b, R2c = H, (un)substituted alkyl, aryl, etc.; NR2bR2c = (un)substituted heterocyclyl; R3a = H, (un)substituted alkyl, aryl, etc.] which have

utility in the treatment of conditions which benefit from administration of neuroprotective agents generally, including treatment of central and peripheral nervous condition as well as for promoting nerve cell differentiation, were prepd. Thus, reacting 4-oxo-3-(piperidylmethyl)-4-(3-pyridyl)butanoic acid with Me N-acetyl-L-cysteine ester in EtOH afforded 85% (R)-II. Biol. data for compds. I were given.

ST aminoethylthiomethyloxopyridylbutanoic acid prepn neuroprotectant;  
pyridylbutanoic acid aminoethylthiomethyloxo prepn neuroprotectant

IT Cytoprotective agents

(neuroprotectants; prepn. of 3-(2-aminoethylthio)methyl-4-oxo-4-(3-pyridyl)butanoic acid derivs. as neuroprotective agents)

IT 343574-29-8P 343574-71-0P 343575-87-1P 343575-92-8P 343577-16-2P  
343577-17-3P 343577-23-1P 343577-91-3P

RL: BAC (Biological activity or effector, except adverse); RCT (Reactant);  
SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of 3-(2-aminoethylthio)methyl-4-oxo-4-(3-pyridyl)butanoic acid derivs. as neuroprotective agents)

IT 220301-04-2P 343573-75-1P 343573-76-2P 343573-77-3P 343573-78-4P  
343573-79-5P 343573-80-8P 343573-81-9P 343573-82-0P 343573-83-1P  
343573-84-2P 343573-85-3P 343573-86-4P 343573-87-5P 343573-88-6P  
343573-89-7P 343573-90-0P 343573-91-1P 343573-92-2P 343573-93-3P  
343573-94-4P 343573-95-5P 343573-96-6P 343573-97-7P 343573-98-8P  
343573-99-9P 343574-00-5P 343574-01-6P 343574-02-7P 343574-03-8P  
343574-04-9P 343574-05-0P 343574-06-1P 343574-07-2P 343574-08-3P  
343574-09-4P 343574-10-7P 343574-11-8P 343574-12-9P 343574-13-0P  
343574-14-1P 343574-15-2P 343574-16-3P 343574-17-4P 343574-18-5P  
343574-19-6P 343574-20-9P 343574-21-0P 343574-22-1P 343574-23-2P  
343574-24-3P 343574-25-4P 343574-26-5P 343574-27-6P 343574-28-7P  
343574-30-1P 343574-31-2P 343574-32-3P 343574-33-4P 343574-34-5P  
343574-35-6P 343574-36-7P 343574-37-8P 343574-38-9P 343574-39-0P  
343574-40-3P 343574-41-4P 343574-42-5P 343574-43-6P 343574-44-7P  
343574-45-8P 343574-46-9P 343574-47-0P 343574-48-1P 343574-49-2P  
343574-50-5P 343574-51-6P 343574-52-7P 343574-53-8P 343574-54-9P  
343574-55-0P 343574-56-1P 343574-57-2P 343574-58-3P 343574-59-4P  
343574-60-7P 343574-61-8P 343574-62-9P 343574-63-0P 343574-64-1P  
343574-65-2P 343574-66-3P 343574-67-4P 343574-68-5P 343574-69-6P  
343574-70-9P 343574-72-1P 343574-73-2P 343574-74-3P 343574-75-4P  
343574-76-5P 343574-77-6P 343574-78-7P 343574-79-8P 343574-80-1P  
343574-81-2P 343574-82-3P 343574-83-4P 343574-84-5P 343574-85-6P  
343574-86-7P 343574-87-8P 343574-88-9P 343574-90-3P 343574-91-4P  
343574-92-5P 343574-93-6P 343574-94-7P 343574-95-8P 343574-96-9P  
343574-97-0P 343574-98-1P 343574-99-2P 343575-00-8P 343575-01-9P  
343575-02-0P 343575-03-1P 343575-04-2P 343575-05-3P 343575-06-4P  
343575-07-5P 343575-08-6P 343575-09-7P 343575-10-0P 343575-11-1P  
343575-13-3P 343575-14-4P 343575-15-5P 343575-16-6P 343575-17-7P  
343575-18-8P 343575-19-9P 343575-20-2P 343575-21-3P 343575-22-4P  
343575-23-5P 343575-24-6P 343575-25-7P 343575-26-8P 343575-27-9P  
343575-28-0P 343575-29-1P 343575-30-4P 343575-31-5P 343575-32-6P  
343575-33-7P 343575-34-8P 343575-35-9P 343575-36-0P 343575-37-1P  
343575-38-2P 343575-39-3P 343575-40-6P 343575-41-7P 343575-42-8P  
343575-43-9P 343575-44-0P 343575-45-1P 343575-46-2P  
**343575-47-3P** 343575-48-4P 343575-49-5P 343575-50-8P  
343575-51-9P 343575-52-0P 343575-53-1P 343575-54-2P 343575-55-3P  
343575-56-4P 343575-57-5P 343575-58-6P 343575-59-7P 343575-60-0P  
343575-61-1P 343575-62-2P 343575-63-3P 343575-64-4P 343575-65-5P  
343575-66-6P 343575-67-7P 343575-68-8P 343575-69-9P 343575-70-2P  
343575-71-3P 343575-72-4P 343575-73-5P 343575-74-6P 343575-75-7P  
343575-76-8P 343575-77-9P 343575-78-0P 343575-79-1P 343575-80-4P  
343575-81-5P 343575-82-6P 343575-83-7P 343575-84-8P 343575-85-9P

343575-86-0P	343575-88-2P	343575-89-3P	343575-90-6P	343575-91-7P
343575-93-9P	343575-94-0P	343575-95-1P	343575-96-2P	343575-97-3P
343575-99-5P	343576-00-1P	343576-01-2P	343576-02-3P	343576-03-4P
343576-04-5P	343576-05-6P	343576-06-7P	343576-07-8P	343576-08-9P
343576-09-0P	343576-10-3P	343576-11-4P	343576-12-5P	343576-13-6P
343576-14-7P	343576-15-8P			

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of 3-(2-aminoethylthio)methyl-4-oxo-4-(3-pyridyl)butanoic acid derivs. as neuroprotective agents)

IT	343576-16-9P	343576-17-0P	343576-18-1P	343576-19-2P	343576-20-5P
	343576-21-6P	343576-22-7P	343576-23-8P	343576-24-9P	343576-25-0P
	343576-26-1P	343576-27-2P	343576-28-3P	343576-29-4P	343576-30-7P
	343576-31-8P	343576-32-9P	343576-33-0P	343576-34-1P	343576-35-2P
	343576-36-3P	343576-37-4P	343576-38-5P	343576-39-6P	343576-40-9P
	343576-41-0P	343576-42-1P	343576-43-2P	343576-44-3P	343576-45-4P
	343576-46-5P	343576-47-6P	343576-48-7P	343576-49-8P	343576-50-1P
	343576-51-2P	343576-52-3P	343576-53-4P	343576-54-5P	343576-55-6P
	343576-56-7P	343576-57-8P	343576-58-9P	343576-59-0P	343576-60-3P
	343576-61-4P	343576-62-5P	343576-63-6P	343576-64-7P	343576-65-8P
	343576-66-9P	343576-67-0P	343576-68-1P	343576-69-2P	343576-70-5P
	343576-71-6P	343576-72-7P	343576-73-8P	343576-74-9P	343576-75-0P
	343576-76-1P	343576-77-2P	343576-78-3P	343576-79-4P	343576-80-7P
	343576-81-8P	343576-82-9P	343576-83-0P	343576-84-1P	343576-85-2P
	343576-86-3P	343576-87-4P	343576-88-5P	343576-89-6P	343576-90-9P
	343576-91-0P	343576-92-1P	343576-93-2P	343576-94-3P	343576-95-4P
	343576-96-5P	343576-97-6P	343576-98-7P	343576-99-8P	343577-00-4P
	343577-01-5P	343577-02-6P	343577-03-7P	343577-04-8P	343577-05-9P
	343577-06-0P	343577-07-1P	343577-08-2P	343577-09-3P	343577-10-6P
	343577-11-7P	343577-12-8P	343577-13-9P	343577-14-0P	343577-15-1P
	343577-18-4P	343577-19-5P	343577-20-8P	343577-21-9P	343577-22-0P
	343577-24-2P	343577-25-3P	343577-26-4P	343577-27-5P	343577-28-6P
	343577-29-7P	343577-30-0P	343577-31-1P	343577-32-2P	343577-33-3P
	343577-34-4P	343577-35-5P	343577-36-6P	343577-37-7P	343577-38-8P
	343577-39-9P	343577-40-2P	343577-41-3P	343577-42-4P	343577-43-5P
	343577-44-6P	343577-45-7P	343577-46-8P	343577-47-9P	343577-48-0P
	343577-49-1P	343577-50-4P	343577-51-5P	343577-52-6P	343577-53-7P
	343577-54-8P	343577-55-9P	343577-56-0P	343577-57-1P	343577-58-2P
	343577-59-3P	343577-60-6P	343577-61-7P	343577-62-8P	343577-63-9P
	343577-64-0P	343577-65-1P	343577-66-2P	343577-67-3P	343577-68-4P
	343577-69-5P	343577-70-8P	343577-71-9P	343577-72-0P	343577-73-1P
	343577-74-2P	343577-75-3P	343577-76-4P	343577-77-5P	343577-78-6P
	343577-79-7P	343577-80-0P	343577-81-1P	343577-82-2P	343577-83-3P
	343577-84-4P	343577-85-5P	343577-86-6P	343577-87-7P	343577-88-8P
	343577-89-9P	343577-90-2P	343577-92-4P	343577-93-5P	343577-94-6P
	343577-95-7P	343577-96-8P	343577-97-9P	343577-99-1P	343578-00-7P
	343578-01-8P	343578-02-9P	343578-03-0P	343578-04-1P	343578-05-2P
	343578-06-3P	343578-07-4P	343578-08-5P	343578-09-6P	343578-10-9P
	343578-11-0P	343578-12-1P	343578-13-2P	343578-14-3P	343578-15-4P
	343578-16-5P	343578-17-6P	343578-18-7P	343578-19-8P	343578-20-1P
	343578-21-2P	343578-22-3P	343578-23-4P	343578-24-5P	343578-25-6P
	343578-31-4P	343964-43-2P			

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of 3-(2-aminoethylthio)methyl-4-oxo-4-(3-pyridyl)butanoic acid derivs. as neuroprotective agents)

IT 71-23-8, n-Propanol, reactions 86-84-0, 1-Naphthyl isocyanate 96-33-3, Methyl acrylate 104-12-1, 4-Chlorophenyl isocyanate 104-97-2,

Cyclopentanepropanoyl chloride 110-89-4, Piperidine, reactions  
 140-88-5, Ethyl acrylate 141-32-2 500-22-1, 3-Pyridinecarboxaldehyde  
 501-53-1, Benzyl chloroformate 622-58-2, 4-Methylphenyl isocyanate  
 700-87-8, 2-Methoxyphenyl isocyanate 1016-19-9, 3,4,5-Trimethoxyphenyl  
 isocyanate 1190-73-4, n-Acetylcysteamine 1195-45-5, 4-Fluorophenyl  
 isocyanate 1476-23-9, Allyl isocyanate 2859-67-8, 3-Pyridinepropanol  
 3025-95-4 3173-53-3, Cyclohexyl isocyanate 4192-31-8,  
 4-Oxo-4-(3-pyridyl)butanoic acid 4530-20-5 6068-72-0, 4-Cyanobenzoyl  
 chloride 7652-46-2 15159-40-7, Morpholine-4-carbonyl chloride  
 16744-98-2, 2-Fluorophenyl isocyanate 20938-74-3 36643-74-0, Indanol  
 54132-75-1, 3,5-Dimethylphenyl isocyanate 59587-09-6 67385-09-5  
 69812-29-9 129714-97-2, 3,5-Difluorobenzoyl chloride 220301-12-2  
 343578-28-9 343578-29-0 343578-30-3

RL: RCT (Reactant)

(prepn. of 3-(2-aminoethylthio)methyl-4-oxo-4-(3-pyridyl)butanoic acid  
 derivs. as neuroprotective agents)

IT 59086-27-0P 61192-47-0P 343573-70-6P 343573-71-7P 343573-72-8P  
 343578-26-7P 343578-27-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(prepn. of 3-(2-aminoethylthio)methyl-4-oxo-4-(3-pyridyl)butanoic acid  
 derivs. as neuroprotective agents)

IT 343575-47-3P


RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic  
 preparation); THU (Therapeutic use); BIOL (Biological study); PREP  
 (Preparation); USES (Uses)

(prepn. of 3-(2-aminoethylthio)methyl-4-oxo-4-(3-pyridyl)butanoic acid  
 derivs. as neuroprotective agents)

RN 343575-47-3 HCAPLUS

CN INDEX NAME NOT YET ASSIGNED

Absolute stereochemistry.



L27 ANSWER 2 OF 24 HCAPLUS COPYRIGHT 2001 ACS

AN 2000:592548 HCAPLUS

DN 133:177486

TI Preparation of substituted stilbene compounds with vascular damaging  
 activity

IN Davis, Peter David

PA Angiogene Pharmaceuticals Ltd., UK

SO PCT Int. Appl., 31 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K031-195

ICS A61P035-00; A61P017-00; A61P027-02

CC 34-2 (Amino Acids, Peptides, and  
 Proteins)

Section cross-reference(s): 1, 25

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000048590	A1	20000824	WO 2000-GB503	20000215
	<p>W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM</p> <p>RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG</p>				
PRAI	GB 1999-3403	A	19990216		
OS	MARPAT 133:177486				
AB	<p>A vascular damaging agent AXB (A = substituted cis-stilbene; X = linker bond, atom, or group; B = moiety derived from an inhibitor of the formation or action of NO in mammalian systems), is claimed. Thus, (Z)-1-[3-(N-.alpha.-tert-butoxycarbonyl-N-.omega.-nitroarginyloxy)-4-methoxyphenyl]-2-(3,4,5-trimethoxyphenyl)ethene was stirred with CF3CO2H in CH2Cl2 to give (Z)-1-(4-methoxy-3-NG-nitroarginyloxyphenyl)-2-(3,4,5-trimethoxyphenyl)ethene. The latter at 50 mg/kg i.p. in mice bearing CaNT or SaS tumors gave 95% redn. in vascular vol. and 91-100% tumor necrosis.</p>				
ST	<p>vascular damaging agent substituted stilbene prepn; neovascularization disease treatment substituted stilbene; nitroarginyloxystilbene prepn tumor vascular damaging activity; nitric oxide synthase inhibitor nitroarginyloxystilbene prepn</p>				
IT	<p>Angiogenesis (neovascularization, treatment of diseases involving neovascularization; prepn. of substituted stilbene compds. with vascular damaging activity)</p>				
IT	<p>Amino acids, preparation RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (nitroarginyloxystilbene derivs.; prepn. of substituted stilbene compds. with vascular damaging activity)</p>				
IT	<p>Antitumor agents (prepn. of substituted stilbene compds. with vascular damaging activity)</p>				
IT	<p>125978-95-2, Nitric oxide synthase RL: BSU (Biological study, unclassified); MSC (Miscellaneous); BIOL (Biological study) (inhibitors; prepn. of substituted stilbene compds. with vascular damaging activity)</p>				
IT	<p>288585-54-6P <b>288585-55-7P 288585-56-8P 288585-57-9P</b> RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (prepn. of substituted stilbene compds. with vascular damaging activity)</p>				
IT	<p>51298-62-5 61240-20-8, 3,4,5-Trimethoxybenzyltriphenylphosphonium bromide 97315-18-9 117048-59-6 288585-58-0 RL: RCT (Reactant) (prepn. of substituted stilbene compds. with vascular damaging activity)</p>				
IT	<p>288585-59-1P 288585-60-4P RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)</p>				

(prepn. of substituted stilbene compds. with vascular damaging activity)

RE.CNT 5

RE

- (1) Ajinomoto Kk; EP 0641767 A 1995 HCAPLUS
- (2) Aston Molecules Ltd; WO 9216486 A 1992 HCAPLUS
- (3) George, R; ANTI-CANCER DRUG DESIGN 1995; V10(4), P299
- (4) Koji, O; JOURNAL OF MEDICINAL CHEMISTRY 1998, V41(16), P3022
- (5) Ohsumi, K; BIOORGANIC & MEDICINAL CHEMISTRY LETTERS 1998, V8(22), P3153 HCAPLUS

IT 288585-55-7P 288585-56-8P 288585-57-9P


RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of substituted stilbene compds. with vascular damaging activity)

RN 288585-55-7 HCAPLUS

CN L-Ornithine, N5-[imino(nitroamino)methyl]-N2-[[2-methoxy-5-[(1Z)-2-(3,4,5-trimethoxyphenyl)ethenyl]phenoxy]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.  
Double bond geometry as shown.



RN 288585-56-8 HCAPLUS

CN L-Ornithine, N5-[imino(nitroamino)methyl]-N2-[[2-methoxy-5-[(1Z)-2-(3,4,5-trimethoxyphenyl)ethenyl]phenoxy]carbonyl]-, (9CI) (CA INDEX NAME)


Absolute stereochemistry.  
Double bond geometry as shown.



RN 288585-57-9 HCAPLUS

CN L-Ornithine, N5-[imino(nitroamino)methyl]-N2-[[2-methyl-5-[(1Z)-2-(3,4,5-trimethoxyphenyl)ethenyl]phenoxy]carbonyl]-, methyl ester (9CI) (CA INDEX NAME)




Absolute stereochemistry.  
Double bond geometry as shown.



L27 ANSWER 3 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
AN 1999:753019 HCAPLUS  
DN 132:12506  
TI Preparation of peptides for treating diseases of excessive bone loss or  
cartilage or matrix degradation as cysteine protease inhibitors  
IN Bondinell, William Edward; Desjarlais, Renee Louise; Veber, Daniel Frank;  
Yamashita, Dennis Shinji  
PA Smithkline Beecham Corporation, USA  
SO PCT Int. Appl., 128 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
IC ICM A61K  
CC 34-3 (Amino Acids, Peptides, and  
Proteins)  
Section cross-reference(s): 1, 7, 63

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9959526	A2	19991125	WO 1999-US11266	19990520
	WO 9959526	A3	20000120		
	W:	AE, AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZA, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
	EP 1067894	A2	20010117	EP 1999-924421	19990520
	R:	BE, CH, DE, ES, FR, GB, IT, LI, NL			
PRAI	US 1998-86557	P	19980521		
	WO 1999-US11266	W	19990520		
OS	MARPAT 132:12506				
GI					



AB The present invention provides peptides bis-aminomethyl carbonyl protease inhibitors I (R<sup>1</sup>, R<sup>2</sup> = alkyl; P = X, Y; R<sup>3</sup> selected from the group consisting of: CH<sub>2</sub>CH(CH<sub>3</sub>)<sub>2</sub>, CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>, CH<sub>2</sub>CH=CH<sub>2</sub>, or CH<sub>2</sub>Ph; R<sup>4</sup> is selected from the group consisting of alkyl; N-piperazine; N-tetrahydroisoquinoline; substituted alkyl, Ph, benzofuran, benzothiazole; quinoline; naphthyl; and benzoxazole; R<sup>5</sup> = Ph and Ph substituted with alkyl, N-piperidine, benzofuran; pyridine; Q = arylacyl) and pharmaceutically acceptable salts, hydrates and solvates thereof which inhibit proteases, including cathepsin K, pharmaceutical compns. of such compds., and methods for treating diseases of excessive bone loss or cartilage or matrix degrdn., including osteoporosis; gingival disease including gingivitis and periodontitis; arthritis, more specifically, osteoarthritis and rheumatoid arthritis; Paget's disease; hypercalcemia of malignancy; and metabolic bone disease, comprising inhibiting said bone loss or excessive cartilage or matrix degrdn. by administering to a patient in need thereof a compd. of the present invention. Thus, (S)-3N-(N-(thianaphthenyl-2-carbonyl)-leuciny)-amino-1N-(3-(2-(1-oxo)-pyridyl)phenylacetyl)-amino-butan-2-one was prepd. for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitor. Detn. of cathepsin K proteolytic catalytic activity of these compds. are reported.

ST osteoarthritis osteoporosis rheumatoid arthritis peptide prepn; bone loss cartilage degrdn peptide prepn protease inhibitor

IT Cartilage  
(degeneration; prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

IT Bone  
(demineralization; prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

IT Bone, disease  
Osteoarthritis  
Osteoporosis  
Rheumatoid arthritis

(prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

IT Peptides, preparation  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

IT 251458-51-2P

RL: BAC (Biological activity or effector, except adverse); RCT (Reactant);



SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

IT	247119-79-5P	247119-80-8P	247119-82-0P	247119-83-1P	251457-08-6P
	251457-09-7P	251457-10-0P	251457-14-4P	251457-15-5P	251457-16-6P
	251457-17-7P	251457-18-8P	251457-19-9P	251457-21-3P	251457-23-5P
	251457-24-6P	251457-27-9P	251457-28-0P	251457-30-4P	251457-32-6P
	251457-34-8P	251457-35-9P	251457-36-0P	251457-37-1P	251457-38-2P
	251457-39-3P	251457-40-6P	251457-41-7P	251457-42-8P	251457-43-9P
	251457-44-0P	251457-45-1P	251457-46-2P	251457-47-3P	251457-48-4P
	251457-49-5P	251457-50-8P	251457-51-9P	251457-52-0P	251457-53-1P
	251457-54-2P	251457-55-3P	251457-56-4P	251457-60-0P	251457-62-2P
	251457-64-4P	251457-65-5P	251457-66-6P	251457-68-8P	251457-69-9P
	251457-71-3P	251457-74-6P	251457-75-7P	251457-76-8P	251457-77-9P
	251457-78-0P	251457-79-1P	251457-80-4P	251457-82-6P	251457-84-8P
	251457-85-9P	251457-87-1P	251457-89-3P	251457-90-6P	251457-91-7P
	251457-92-8P	251457-93-9P	251457-94-0P	251457-95-1P	251457-96-2P
	251457-97-3P	251457-98-4P	251457-99-5P	251458-01-2P	251458-02-3P
	251458-04-5P	251458-05-6P	251458-06-7P	251458-08-9P	251458-09-0P
	251458-10-3P	251458-11-4P	251458-12-5P	251458-13-6P	251458-17-0P
	251458-18-1P	251458-19-2P	251458-20-5P	251458-24-9P	251458-29-4P
	251458-31-8P	251458-32-9P	251458-33-0P	251458-35-2P	251458-37-4P
	251458-40-9P	251458-41-0P	251458-42-1P	251458-43-2P	251458-44-3P
	251458-45-4P	251458-50-1P	251458-52-3P	251458-53-4P	251458-55-6P
	251458-57-8P	251458-59-0P	251458-61-4P	251458-62-5P	251458-63-6P
	251458-64-7P	251458-65-8P	251458-66-9P	251458-67-0P	251458-68-1P
	251458-69-2P	251458-70-5P	251458-71-6P	251458-78-3P	251458-79-4P
	251458-80-7P				

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

IT 37353-41-6, Cysteine protease

RL: BPR (Biological process); BIOL (Biological study); PROC (Process)

(prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

IT 50-43-1, 2,4,6-Trichlorobenzoic acid 51-44-5, 3,4-Dichlorobenzoic acid  
 65-85-0, Benzoic acid, reactions 75-08-1, Ethanethiol 79-09-4,  
 Propionic acid, reactions 86-59-9, 8-Quinoline-carboxylic acid  
 88-13-1, Thiophene-3-carboxylic acid 91-21-4, 1,2,3,4-  
 Tetrahydroisoquinoline 92-92-2, Biphenyl-4-carboxylic acid" 93-07-2,  
 3,4-Dimethoxybenzoic acid" 93-09-4, 2-Naphthoic acid 93-10-7,  
 2-Quinolinylcarboxylic acid 98-09-9, Benzenesulfonyl chloride 98-97-5,  
 Pyrazine-2-carboxylic acid" 99-96-7, 4-Hydroxybenzoic acid, reactions  
 100-00-5, 1-Chloro-4-nitro-benzene 100-09-4, 4-Methoxybenzoic acid"  
 109-01-3 109-52-4, Valeric acid, reactions 142-25-6 349-88-2,  
 4-Fluorophenylsulfonyl chloride 371-40-4, 4-Fluoro-aniline 402-54-0  
 455-24-3, 4-(Trifluoromethyl)benzoic acid 456-22-4, 4-Fluoro-benzoic  
 acid 496-41-3, Benzofuran-2-carboxylic acid 501-52-0,  
 3-Phenylpropionic acid 504-29-0, 2-Amino-pyridine 536-69-6,  
 5-Butylpyridine-2-carboxylic acid 585-76-2, 3-Bromo-benzoic acid  
 611-95-0, 4-(Benzoyl)benzoic acid 616-29-5, 1,3-Diamino-propan-2-ol  
 619-45-4, 4-Methoxycarbonyl-aniline 620-86-0 622-40-2,  
 4-(2-Hydroxyethyl)morpholine 636-82-8, 1-Cyclohexene-1-carboxylic acid  
 646-07-1 661-69-8, Hexamethylditin 828-51-3, Adamantane-1-carboxylic  
 acid 883-21-6, 1-Methoxy-2-naphthoic acid 883-62-5,  
 3-Methoxy-2-naphthoic acid 1457-59-6, 5-Methylimidazolyl-4-carboxylic  
 acid 1477-50-5, Indole-2-carboxylic acid 1486-51-7 1623-92-3,

4-Phenoxybenzenesulfonyl chloride 1670-82-2, Indole-6-carboxylic acid 1723-27-9, Thieno[3,2-b]thiophene-2-carboxylic acid 1878-67-7, 3-Bromo phenyl acetic acid 2008-75-5, 2-(Piperidin-1-yl)ethyl chloride hydrochloride" 2666-93-5 2991-42-6, 4-Trifluoromethylbenzenesulfonyl chloride 3328-70-9, 5-Formylsalicylaldehyde 3510-66-5, 2-Bromo-5-methylpyridine 3622-35-3, 6-Benzothiazolecarboxylic acid 3647-69-6, 2-(4-Morpholinyl)ethyl chloride hydrochloride 4457-32-3, 4-Nitrobenzyl chloroformate 4467-07-6, 3-(2-Pyridyl)benzoic acid 4926-28-7, 2-Bromo-4-methylpyridine 5315-25-3, 2-Bromo-6-methylpyridine 5731-13-5 6314-28-9, Benzo[b]thiophene-2-carboxylic acid 6315-89-5, 3,4-Dimethoxy-aniline 6480-68-8, 3-Quinolinecarboxylic acid 6702-50-7, Methyl 3-hydroxy-4-methoxy-benzoate 6973-60-0 7250-53-5, 5-Quinoline-carboxylic acid 10147-37-2, Isopropyl sulfonyl chloride 10349-57-2, 6-Quinolinecarboxylic acid 15112-41-1, 5-Benzoxazolecarboxylic acid 16309-45-8 16419-60-6 19438-10-9, Methyl 3-hydroxybenzoate 20029-52-1, 4-Cyclohexylbenzoic acid 27298-97-1 31462-59-6, Pyrimidine-4-carboxylic acid 32315-10-9, Triphosgene 40274-67-7 50551-56-9 50793-29-8, 4-(4-Cyanophenoxy)benzoic acid 57260-71-6, 1-(tert-Butoxycarbonyl)piperazine 65007-00-3 66715-65-9, 2-Pyridinesulfonyl chloride 68947-43-3, N-Methylpiperidine-4-carboxylic acid 73579-08-5 78161-82-7, 4-(4-(Trifluoromethyl)phenoxy)benzoic acid 81432-12-4 90433-20-8 92198-45-3 98327-87-8, 2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl 98437-24-2, 2-Benzofuran boronic acid 144059-86-9 150529-73-0, Methyl 3-bromophenylacetate 150798-78-0 154235-77-5, Benzoxazole-6-carboxylic acid 154739-53-4 175137-58-3 175201-51-1 190661-70-2 210304-58-8 215947-98-1 228419-12-3 248923-01-5 251457-20-2 251457-33-7 **251457-72-4** 251457-88-2 251458-03-4 251458-15-8 251458-26-1 251458-30-7 251458-34-1 251458-38-5 251458-39-6 251458-73-8

RL: RCT (Reactant)

(prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

IT 23948-77-8P, [1,1'-Biphenyl]-3-acetic acid 51061-68-8P 69038-74-0P  
75852-28-7P 99370-68-0P 203503-06-4P 215309-01-6P 216316-23-3P  
216316-24-4P 216316-27-7P 224643-30-5P 227178-31-6P 227178-32-7P  
229630-60-8P 247119-96-6P 247119-99-9P 247122-09-4P 250726-45-5P  
250726-46-6P 250726-47-7P 251457-05-3P 251457-06-4P 251457-07-5P  
251457-11-1P 251457-12-2P 251457-13-3P 251457-22-4P 251457-25-7P  
251457-26-8P 251457-29-1P 251457-31-5P 251457-57-5P 251457-58-6P  
251457-59-7P 251457-61-1P 251457-63-3P 251457-67-7P 251457-70-2P  
251457-73-5P 251458-00-1P 251458-07-8P 251458-14-7P 251458-16-9P  
251458-21-6P 251458-22-7P 251458-23-8P 251458-25-0P 251458-27-2P  
251458-28-3P 251458-36-3P 251458-46-5P 251458-47-6P 251458-48-7P  
251458-49-8P 251458-72-7P 251458-74-9P 251458-75-0P 251458-76-1P  
251458-77-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

IT **251457-72-4**


RL: RCT (Reactant)

(prepn. of peptides for treating diseases of excessive bone loss or cartilage or matrix degrdn. as cysteine protease inhibitors)

RN 251457-72-4 HCAPLUS

CN L-Leucine, N-[(4-nitrophenoxy)carbonyl]-, 1,1-dimethylethyl ester (9CI)  
(CA INDEX NAME)

Absolute stereochemistry.



L27 ANSWER 4 OF 24 HCAPLUS COPYRIGHT 2001 ACS

AN 1998:749736 HCAPLUS

DN 130:110572

TI Self-Immolative Nitrogen Mustard Prodrugs for Suicide Gene Therapy  
 AU Niculescu-Duvaz, Dan; Niculescu-Duvaz, Ion; Friedlos, Frank; Martin, Janet; Spooner, Robert; Davies, Lawrence; Marais, Richard; Springer, Caroline J.

CS UK

SO J. Med. Chem. (1998), 41(26), 5297-5309

CODEN: JMCMAR; ISSN: 0022-2623

PB American Chemical Society


DT Journal

LA English


CC 34-2 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 1, 25


GI



I



II



III

AB Four new potential self-immolative prodrugs derived from phenol and aniline nitrogen mustards, four model compds. derived from their corresponding fluoroethyl analogs and two new self-immolative linkers were designed and synthesized for use in the suicide gene therapy termed GDEPT (gene-directed enzyme prodrug therapy). The self-immolative prodrugs were designed to be activated by the enzyme carboxypeptidase G2 (CPG2) releasing an active drug by a 1,6-elimination mechanism via an unstable intermediate. Thus, glutamate derivs. I (Y = O, NH; Y = O, NH; X = Cl)

were synthesized. They are bifunctional alkylating agents in which the activating effects of the phenolic hydroxyl or amino functions are masked through an oxycarbonyl or a carbamoyl bond to a benzylic spacer which is itself linked to a glutamic acid by an oxycarbonyl or a carbamoyl bond. The corresponding fluoroethyl compds. I (X = F) were also synthesized. The rationale was to obtain model compds. with greatly reduced alkylating abilities that would be much less reactive with nucleophiles compared to the corresponding chloroethyl derivs. This enabled studies of these model compds. as substrates for CPG2, without incurring the rapid and complicated decompn. pathways of the chloroethyl derivs. The prodrugs were designed to be activated to their corresponding phenol and aniline nitrogen mustard drugs by CPG2 for use in GDEPT. The synthesis of the analogous novel parent drugs II is also described. A colorectal cell line was engineered to express CPG2 tethered to the outer cell surface. The phenylenediamine compds. were found to behave as prodrugs, yielding IC50 prodrug/IC50 drug ratios between 20- and 33-fold for I (X = Cl, Y = NH; Z = O, NH) and differentials of 12-14-fold between CPG2-expressing and control LacZ-expressing clones. The drugs released are up to 70-fold more potent than benzoic acid III that results from the glutamate prodrug (CMDA) which has been used previously for GDEPT. These data demonstrate the viability of this strategy and indicate that self-immolative prodrugs can be synthesized to release potent mustard drugs selectively by cells expressing CPG2 tethered to the cell surface in GDEPT.

- ST glutamate nitrogen mustard prodrug prepn suicide gene therapy; immolative nitrogen mustard prodrug suicide gene therapy; carboxypeptidase G2 cleavage glutamate nitrogen mustard prodrug prepn; anticancer agent glutamate nitrogen mustard prodrug prepn; cytotoxicity glutamate nitrogen mustard prodrug prepn
- IT Chemotherapy  
(gene-directed enzyme prodrug; prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)
- IT Antitumor agents  
Cytotoxicity  
Gene therapy  
Prodrugs  
(prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)
- IT 122665-70-7  
RL: BAC (Biological activity or effector, except adverse); BIOL (Biological study)  
(prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)
- IT 1204-69-9, 4-[Bis(2-chloroethyl)amino]phenol 2067-58-5  
RL: BAC (Biological activity or effector, except adverse); RCT (Reactant); BIOL (Biological study)  
(prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)
- IT 219591-87-4P 219592-00-4P  
RL: BAC (Biological activity or effector, except adverse); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)
- IT 122665-73-ODP, N-4-[(2-Chloroethyl)(2-mesyloxyethyl)amino]benzoyl-L-glutamic acid, dichloro and difluoro carbonate and carbamate analogs  
180838-98-6P 180839-01-4P 180839-03-6P 180839-05-8P 219591-76-1P  
219591-80-7P 219591-82-9P 219591-86-3P 219591-92-1P 219591-94-3P  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)

IT 9074-87-7, Carboxypeptidase G2  
 RL: BPR (Biological process); BIOL (Biological study); PROC (Process)  
 (prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)

IT 219591-77-2P 219591-79-4P  
 RL: BYP (Byproduct); PREP (Preparation)  
 (prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)

IT 219591-73-8P  
 RL: BYP (Byproduct); RCT (Reactant); PREP (Preparation)  
 (prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)

IT 123-08-0, 4-Hydroxybenzaldehyde 619-73-8, 4-Nitrobenzyl alcohol  
 18226-17-0 20845-16-3 32677-01-3, Di-tert-butyl L-glutamate  
 hydrochloride 101582-69-8  
 RL: RCT (Reactant)  
 (prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)

IT 18483-99-3P 18484-05-4P 57529-05-2P 113068-95-4P 161803-03-8P  
 161803-04-9P 161803-05-0P 161803-06-1P 180838-97-5P 180839-00-3P  
 180839-02-5P 180839-04-7P 180839-06-9P 180839-11-6P 180839-13-8P  
 180839-14-9P 180839-15-0P 180839-16-1P 180839-18-3P 180839-19-4P  
**180839-20-7P 180839-21-8P** 180841-62-7P 219591-81-8P  
 219591-83-0P 219591-85-2P 219591-91-0P 219591-93-2P 219591-96-5P  
 219591-97-6P 219591-98-7P 219592-01-5P 219592-02-6P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)

IT 219591-75-0P 219591-89-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)

RE.CNT 29

RE

- (1) Alexander, R; Tetrahedron Lett 1991, V32, P3269 HCAPLUS
- (2) Bridgewater, G; Eur J Cancer 1995, V31A, P2362
- (3) Carl, P; J Med Chem 1981, V24, P479 HCAPLUS
- (4) Corey, E; J Am Chem Soc 1972, V94, P6190 HCAPLUS
- (5) Dowell, R; J Med Chem 1996, V39, P1100 HCAPLUS
- (6) Dubowchik, G; Tetrahedron Lett 1997, V38, P5261 HCAPLUS
- (7) Friedlos, F; J Med Chem 1997, V40, P1270 HCAPLUS
- (8) Greene, T; Protective groups in organic chemistry 1991
- (9) Hanessian, S; Can J Chem 1975, V53, P2975 HCAPLUS
- (10) Hollitzer, O; Angew Chem Int Ed Engl 1976, V5, P444
- (11) Huber, B; Proc Natl Acad Sci U S A 1991, V88, P8039 HCAPLUS
- (12) Jarman, M; Chem Br 1989, V25, P51 HCAPLUS
- (13) Kruse, C; J Org Chem 1979, V44, P1847 HCAPLUS
- (14) Marais, R; Cancer Res 1996, V56, P4735 HCAPLUS
- (15) Marais, R; Nature Biotechnol 1997, V15, P1373 HCAPLUS
- (16) Minton, N; J Bacteriol 1983, V156, P1222 HCAPLUS
- (17) Niculescu-Duvaz, I; Bioconjugate Chem 1998, V9, P4 HCAPLUS
- (18) Olofson, R; Tetrahedron Lett 1980, V21, P819 HCAPLUS
- (19) Palmer, B; J Med Chem 1990, V33, P112 HCAPLUS
- (20) Perrone, E; EP 0236880 A2 1987 HCAPLUS
- (21) Ross, W; Biological alkylating agents 1962
- (22) Schmidt, F; Bioorg Med Chem Lett 1997, V7, P1071 HCAPLUS
- (23) Schmidt, H; Chem Ber 1979, V112, P727 HCAPLUS
- (24) Sherwood, R; Eur J Biochem 1985, V24, P447
- (25) Springer, C; J Med Chem 1990, V33, P677 HCAPLUS
- (26) Springer, C; J Med Chem 1994, V37, P2361 HCAPLUS

- (27) Springer, C; J Med Chem 1995, V38, P5051 HCAPLUS  
 (28) Wakselman, M; N J Chim 1983, V7, P439 HCAPLUS  
 (29) Zhang, W; Adv Pharmacol (San Diego) 1995, V6, P289


IT 180839-20-7P 180839-21-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of self-immolative nitrogen mustard prodrugs for suicide gene therapy)

RN 180839-20-7 HCAPLUS

CN L-Glutamic acid, N-[[4-[[[(4-nitrophenoxy)carbonyl]oxy]methyl]phenoxy]carbonyl]-, bis(1,1-dimethylethyl) ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 180839-21-8 HCAPLUS

CN L-Glutamic acid, N-[[4-[[[(4-nitrophenoxy)carbonyl]oxy]methyl]phenoxy]carbonyl]-, di-2-propenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A



PAGE 1-B



L27 ANSWER 5 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1998:147326 HCAPLUS  
 DN 128:205147

TI Preparation of non-peptide bombesin receptor antagonists  
 IN Horwell, David Christopher; Pritchard, Martyn Clive  
 PA Warner-Lambert Company, USA; Horwell, David Christopher; Pritchard, Martyn Clive  
 SO PCT Int. Appl., 112 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM C07D401-12  
 ICS A61K031-395; C07D209-20; C07D213-89; C07D213-40; C07D213-56; C07D405-12; C07D417-12  
 CC 34-3 (Amino Acids, Peptides, and Proteins)  
 Section cross-reference(s): 1

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9807718	A1	19980226	WO 1997-US13871	19970806
	W: AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, GH, HU, IL, IS, JP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	BR 9711342	A	19990817	BR 1997-11342	19970222
	AU 9741466	A1	19980306	AU 1997-41466	19970806
	AU 733226	B2	20010510		
	EP 920424	A1	19990609	EP 1997-939359	19970806
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	JP 2001500850	T2	20010123	JP 1998-510779	19970806
	US 6194437	B1	20010227	US 1999-230933	19990203
	NO 9900788	A	19990219	NO 1999-788	19990219
PRAI	US 1996-24323	P	19960822		
	WO 1997-US13871	W	19970806		
OS	MARPAT 128:205147				
AB	Compds. Ar(CR1R8)0-1(CH2)0-1NR4CONR5CR7(CH2Ar1)CONR6(CH2)0-3(CR2R9)0-1(CH2)0-2R3 [Ar = Ph, (un)substituted pyridyl; R1, R2 = H, alkyl, cycloalkyl; R8, R9 = H or forms a ring with R1 or R2, resp; Ar1 = Ar or pyridyl-N-oxide, indolyl, pyridyl, imidazole; R4, R5, R6, R7 = H, Me; R3 = Ar or H, OH, Me2N, N-methylpyrrole, etc.] or their pharmaceutically acceptable salts were prepd. as bombesin receptor antagonists. Thus, 2-[3-(2,6-diisopropylphenyl)ureido]-3-(1H-indol-3-yl)-2-methyl-N-(1-pyridin-2-ylcyclohexylmethyl)propionamide was prepd. by condensation of .alpha.-methyl-L-tryptophan with 2,6-diisopropylphenyl isocyanate, followed by amidation with 1-pyridin-2-ylcyclohexylmethylamine. Affinity binding data (IC50 values) for the product were detd. to be 5 and <10 nM for the NMB and GRP receptors, resp.				
ST	pseudopeptide prepn bombesin receptor antagonist				
IT	Bombesin receptors				
	RL: BPR (Biological process); BIOL (Biological study); PROC (Process) (prepn. of non-peptide bombesin receptor antagonists)				
IT	Peptides, preparation				
	RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses) (pseudopeptides; prepn. of non-peptide bombesin receptor antagonists)				
IT	204066-87-5P 204067-04-9P				
	RL: BAC (Biological activity or effector, except adverse); RCT (Reactant);				

SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of non-peptide bombesin receptor antagonists)

IT 142627-75-6P 185215-75-2P 204066-70-6P 204066-71-7P 204066-72-8P  
204066-73-9P 204066-74-0P 204066-75-1P 204066-76-2P 204066-77-3P  
204066-78-4P 204066-79-5P 204066-80-8P 204066-81-9P 204066-82-0P  
204066-83-1P 204066-84-2P 204066-85-3P 204066-86-4P 204066-89-7P  
204066-91-1P 204066-93-3P 204066-95-5P 204066-97-7P 204066-99-9P  
204067-01-6P 204067-02-7P 204067-03-8P 204067-05-0P 204067-06-1P  
204067-38-9P 204067-40-3P

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of non-peptide bombesin receptor antagonists)

IT 204067-30-1P

RL: BYP (Byproduct); PREP (Preparation)

(prepn. of non-peptide bombesin receptor antagonists)

IT 54-12-6, Tryptophan 62-53-3, Benzenamine, reactions 73-22-3,  
L-Tryptophan, reactions 100-28-7, p-Nitrophenyl isocyanate 111-24-0,  
1,5-Dibromopentane 153-91-3, Tryptophan, .alpha.-methyl- 487-59-2  
1121-78-4 1206-13-9 1548-13-6 2739-97-1, 2-Pyridylacetonitrile  
3218-02-8, Cyclohexanemethanamine 4000-72-0 4442-85-7,  
2-Cyclohexylethylamine 5766-79-0 7693-46-1, p-Nitrophenyl  
chloroformate 16709-25-4, .alpha.-Methyl-L-tryptophan 17540-18-0  
18502-05-1, 4-Imidazoleacetonitrile 23357-52-0 24544-04-5,  
2,6-Diisopropylaniline 25756-29-0, n-Methylcyclohexylmethylamine  
28178-42-9, 2,6-Diisopropylphenyl isocyanate 30806-83-8 35019-66-0  
40465-45-0 50528-53-5 56452-52-9 61341-86-4, s 1-Aminoindan  
70258-01-4 81428-13-9 103110-98-1 114524-80-0 114779-79-2  
135207-25-9 142854-50-0 164323-86-8 204067-18-5 204067-20-9  
204067-24-3 204067-28-7

RL: RCT (Reactant)

(prepn. of non-peptide bombesin receptor antagonists)

IT 5815-73-6P 13139-14-5P 13458-33-8P 35392-66-6P 37982-29-9P  
55270-47-8P 64464-46-6P 75342-32-4P 75342-33-5P 127978-70-5P  
135627-41-7P 135627-42-8P 142946-11-0P 187610-49-7P 187610-50-0P  
190333-61-0P 204067-07-2P 204067-08-3P 204067-09-4P 204067-10-7P  
204067-11-8P 204067-12-9P 204067-13-0P 204067-14-1P 204067-15-2P  
204067-16-3P 204067-17-4P 204067-19-6P 204067-21-0P 204067-22-1P  
204067-23-2P 204067-25-4P 204067-26-5P **204067-27-6P**  
204067-29-8P 204067-31-2P 204067-32-3P 204067-33-4P 204067-34-5P  
204067-35-6P 204067-36-7P 204067-37-8P 204067-39-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(prepn. of non-peptide bombesin receptor antagonists)


IT **204067-27-6P**

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(prepn. of non-peptide bombesin receptor antagonists)

RN 204067-27-6 HCAPLUS

CN Tryptophan, .alpha.-methyl-N-[(4-nitrophenoxy)carbonyl]-, methyl ester  
(9CI) (CA INDEX NAME)





L27 ANSWER 6 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1998:17976 HCAPLUS  
 DN 128:61798  
 TI Preparation of epoxide peptidomimetics as irreversible HIV protease inhibitors  
 IN Yoon, Heungsik; Choy, Nakyeon; Kim, Sung Chun; Choi, Ho Il; Son, Young Chan; Park, Chi Hyo; Moon, Kwang-yul; Jung, Wonhee; Kim, Chung Ryeol; Lee, Chang Sun; Koh, Jong Sung; Kim, Sang Soo  
 PA LG Chemical Ltd., S. Korea  
 SO U.S., 50 pp. Cont.-in-part of U.S. Ser. No. 341,352, abandoned.  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 IC ICM A61K031-44  
 ICS A61K031-47  
 NCL 514314000  
 CC 34-3 (Amino Acids, Peptides, and Proteins)  
 Section cross-reference(s): 1, 63

FAN.CNT 7

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 5696134	A	19971209	US 1995-473877	19950607
	US 5587388	A	19961224	US 1993-159382	19931130
	KR 125117	B1	19971205	KR 1994-13423	19940615
	US 5773468	A	19980630	US 1995-572402	19951214
	US 5744621	A	19980428	US 1996-667888	19960620
	US 5763631	A	19980609	US 1996-667133	19960620
PRAI	US 1993-159382	A2	19931130		
	KR 1994-13423	A	19940615		
	US 1994-341352	B2	19941117		
	KR 1992-23088	A	19921202		
	KR 1992-23089	A	19921202		
	KR 1993-10811	A	19930614		
	KR 1993-21298	A	19931014		
	KR 1993-21299	A	19931014		
	KR 1993-21300	A	19931014		
	US 1995-473877	A2	19950607		
	KR 1995-37292	A	19951026		
OS	MARPAT 128:61798				
GI					

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Novel cis-epoxide compds. I [R1, R2 = independently H, alkyl; R3 = aryl or alkyl (un)substituted with arom.; C3-8 cycloalkyl; R4 = H, C1-4 alkyl; n = 0-2; X = CO, COCO, S(O), SO2, CS; Y = O, CH2, NH, NMe; m = 0, 1; R5 = heterocycle; straight, branched, or cyclic C1-8 alkyl; alkyl substituted with heterocycle or cycloalkyl; straight, branched, or cyclic C1-8 alkoxy; aryl-substituted alkoxy; NR6R7; R6 = straight or branched C1-8 alkyl, cycloalkyl, alkyl substituted with cycloalkyl; R7 = H, alkyl; Z = O, NH, NMe; R8, R9 = independently alkyl (un)substituted with arom. hydrocarbon or cycloalkyl; C3-8 cycloalkyl; arom.] are useful for treating or preventing diseases caused by HIV infection. The novel HIV protease inhibitors I have specific structures to form stable bonding with the enzyme active site, which entails a highly enhanced irreversible

inhibition against HIV protease. Thus deprotection and peptide coupling of olefin II (prepd. in 4 steps from protected L-phenylalaninal and (S)-2-amino-3-methyl-1-phenylbutane) with penicillamine-derived sulfone III (prepd. in 3 steps from L-penicillamine), followed by epoxidn. with MCPBA gave title epoxide deriv. IV. IV showed irreversible inactivation of HIV-1 protease, with a stoichiometric ratio of inhibitor to enzyme of 1:1. IV also showed antiviral activity against HIV-1 with IC50 = 1 nM.

ST epoxide peptidomimetic prepn HIV protease inhibitor; virucide HIV epoxide peptidomimetic prepn; AIDS treatment epoxide peptidomimetic prepn; immunomodulator epoxide peptidomimetic prepn

IT Peptidomimetics

(epoxide; prepn. of epoxide peptidomimetics as irreversible HIV protease inhibitors)

IT Epoxides

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(peptidomimetics; prepn. of epoxide peptidomimetics as irreversible HIV protease inhibitors)

IT Anti-AIDS drugs

Antiviral agents

Human immunodeficiency virus 1

Immunomodulators

(prepn. of epoxide peptidomimetics as irreversible HIV protease inhibitors)

IT	174562-29-9P	174562-30-2P	174562-31-3P	174562-32-4P	174562-33-5P
	174562-34-6P	174562-35-7P	174562-36-8P	174562-37-9P	174562-38-0P
	174562-39-1P	174562-40-4P	174562-41-5P	174562-42-6P	174562-43-7P
	174562-44-8P	174562-45-9P	174562-46-0P	174562-47-1P	174562-48-2P
	174562-49-3P	174562-50-6P	174562-51-7P	174562-52-8P	174562-53-9P
	174562-54-0P	174562-55-1P	174562-56-2P	174562-57-3P	174562-58-4P
	174562-59-5P	174562-60-8P	174562-61-9P	174562-62-0P	174562-63-1P
	174562-65-3P	174562-66-4P	174562-67-5P	174562-68-6P	174562-69-7P
	174562-70-0P	174562-71-1P	174562-72-2P	174562-73-3P	174562-74-4P
	174562-75-5P	174562-76-6P	174562-77-7P	174562-78-8P	174562-79-9P
	174562-80-2P	174562-81-3P	200262-27-7P		

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of epoxide peptidomimetics as irreversible HIV protease inhibitors)

IT 144114-21-6, Retropepsin

RL: BPR (Biological process); BIOL (Biological study); PROC (Process)

(prepn. of epoxide peptidomimetics as irreversible HIV protease inhibitors)

IT 59-67-6, 3-Pyridinecarboxylic acid, reactions 78-77-3, Isobutyl bromide 78-82-0, Isobutyronitrile 88-14-2, 2-Furancarboxylic acid 93-10-7, 2-Quinolinecarboxylic acid 96-41-3, Cyclopentanol 98-00-0, 2-Furanylmethanol 98-59-9, p-Toluenesulfonyl chloride 98-98-6, 2-Pyridinecarboxylic acid 100-46-9, Benzylamine, reactions 100-55-0, 3-Pyridylcarbinol 110-68-9, N-Methyl-N-butylamine 503-74-2, Isovaleric acid 527-72-0, 2-Thiophenecarboxylic acid 574-98-1, N-(2-Bromoethyl)phthalimide 586-95-8, 4-Pyridylcarbinol 586-98-1, 2-Pyridylcarbinol 603-35-0, Triphenylphosphine, reactions 617-89-0, 2-Furanylmethanol 625-45-6, Methoxyacetic acid 1113-41-3, L-Penicillamine 2516-33-8, Cyclopropylmethanol 4083-57-2, 3-Amino-2,4-dimethylpentane 5163-20-2, N-Methyl-N-cyclopropylamine 6921-34-2, Benzylmagnesium chloride 6964-21-2, 3-Thiopheneacetic acid 7693-46-1, p-Nitrophenyl chloroformate 13734-34-4 23844-66-8 24939-24-0, p-Aminobenzenesulfonyl chloride 33445-07-7, Isopropoxyacetic

acid 59830-60-3, N-Benzyloxycarbonyl-L-phenylalaninal 80866-93-9  
96521-86-7 96928-87-9 111491-96-4 123617-80-1, 3-Furanacetic acid  
136465-98-0

RL: RCT (Reactant)

(prepn. of epoxide peptidomimetics as irreversible HIV protease inhibitors)

IT 65273-64-5P 82894-53-9P 97589-56-5P 112898-22-3P 156641-79-1P  
156641-81-5P 156641-83-7P 160742-44-9P 160742-45-0P 160742-70-1P  
160742-71-2P 174562-82-4P 174562-83-5P 174562-84-6P 174562-85-7P  
174562-86-8P 174562-88-0P 174562-89-1P 174562-90-4P  
**174562-91-5P** 174562-92-6P 174562-94-8P 196515-98-7P  
200262-28-8P 200262-29-9P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of epoxide peptidomimetics as irreversible HIV protease inhibitors)

IT 156715-06-9P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of epoxide peptidomimetics as irreversible HIV protease inhibitors)


IT **174562-91-5P**

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of epoxide peptidomimetics as irreversible HIV protease inhibitors)

RN 174562-91-5 HCAPLUS

CN L-Valine, 3-(methylthio)-N-[(4-nitrophenoxy)carbonyl]-, phenylmethyl ester  
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L27 ANSWER 7 OF 24 HCAPLUS COPYRIGHT 2001 ACS

AN 1997:440050 HCAPLUS

DN 127:66223

TI Preparation of urea moiety-containing peptide derivatives as neutral endopeptidase and angiotensin converting enzyme inhibitors

IN Nagano, Masanobu; Takenaka, Yasuhei; Kato, Takeshi

PA Fujisawa Pharmaceutical Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 51 pp.

CODEN: JKXXAF

DT Patent

LA Japanese

IC ICM C07C275-16

ICS A61K031-195; A61K031-215; A61K031-445; A61K038-00; C07C275-18;  
C07D207-06; C07D207-16; C07D211-62; A61K031-40; C12N009-99;  
C07M007-00

CC 34-3 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 1

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

PI JP 09118662 A2 19970506 JP 1996-237543 19960909  
 PRAI GB 1995-18553 19950911  
 OS MARPAT 127:66223  
 AB The title compds. R1CONHACHR2YNR3CONR4CHR6R5 [R1 = heterocyclic ring, etc.; R2 = carboxy, etc.; R3 = alkyl, etc.; R5 = carboxy, etc.; A = alkylene; Y = alkylene; R4 = H, etc.; R6 = (aryl-substituted) alkyl, etc.] are prepd. (2R or 2S)-N-[N-butyl-N-[2-carboxy-3-[(S)-prolylamino]propyl]carbamoyl]-4-phenyl-(S)-phenylalanine trifluoroacetic acid salt (prepn. given) in vitro showed IC50 of  $7.2 \times 10^{-8}$  M against neutral endopeptidase.  
 ST ureidopeptide prepn neutral endopeptidase inhibitor; angiotensin converting enzyme inhibitor ureidopeptide prepn; antihypertensive ureidopeptide prepn prepn  
 IT Anti-inflammatory drugs  
 Antihypertensives  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (urea moiety-contg. peptide derivs. with neutral endopeptidase and angiotensin converting enzyme inhibiting activity)  
 IT 191422-90-9P 191422-91-0P 191422-92-1P 191422-93-2P 191422-94-3P  
 191422-96-5P 191422-98-7P 191422-99-8P 191423-01-5P 191423-02-6P  
 191423-03-7P 191423-04-8P 191423-05-9P 191423-06-0P 191423-07-1P  
 191423-08-2P 191423-09-3P 191423-10-6P 191423-11-7P 191423-12-8P  
 191423-13-9P 191423-14-0P 191423-15-1P 191423-16-2P 191423-17-3P  
 191423-18-4P 191423-19-5P 191423-20-8P 191423-21-9P 191423-22-0P  
 191423-24-2P 191423-26-4P 191423-28-6P 191423-30-0P 191423-32-2P  
 191423-34-4P 191423-36-6P 191423-38-8P 191423-40-2P 191423-42-4P  
 191423-44-6P 191423-46-8P 191423-48-0P 191423-50-4P 191423-52-6P  
 191423-54-8P 191423-56-0P 191423-58-2P 191423-60-6P 191423-62-8P  
 191423-64-0P 191423-66-2P 191423-68-4P 191423-70-8P 191423-72-0P  
 191423-74-2P 191423-76-4P 191423-78-6P 191423-79-7P 191423-80-0P  
 191423-81-1P 191423-83-3P 191423-85-5P 191423-87-7P 191423-89-9P  
 191423-91-3P 191423-93-5P 191423-95-7P 191423-96-8P 191423-97-9P  
 191423-98-0P 191423-99-1P 191424-00-7P 191424-01-8P 191424-02-9P  
 191424-03-0P 191424-04-1P 191424-05-2P 191424-06-3P 191424-07-4P  
 191424-08-5P 191424-09-6P 191424-10-9P 191424-12-1P 191424-13-2P  
 191424-14-3P 191424-15-4P 191424-17-6P 191424-18-7P 191424-19-8P  
 191424-21-2P 191424-23-4P 191424-24-5P 191424-27-8P 191424-28-9P  
 191424-29-0P 191424-31-4P 191424-32-5P 191424-33-6P 191424-34-7P  
 191424-37-0P 191424-40-5P 191424-42-7P 191424-45-0P 191424-48-3P  
 191424-50-7P 191424-53-0P 191424-56-3P 191424-59-6P 191424-61-0P  
 191424-63-2P 191424-65-4P 191424-67-6P 191424-69-8P 191424-70-1P  
 191424-73-4P 191424-75-6P 191424-77-8P 191424-80-3P 191424-82-5P  
 191424-84-7P 191424-86-9P 191424-88-1P 191424-90-5P 191424-92-7P  
 191424-93-8P 191424-94-9P 191424-96-1P 191424-98-3P 191425-00-0P  
 191425-02-2P 191425-04-4P 191425-06-6P 191425-08-8P 191425-10-2P  
 191425-12-4P 191425-13-5P 191425-14-6P 191425-16-8P 191425-18-0P  
 191425-19-1P 191425-21-5P 191425-23-7P 191425-24-8P  
 RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (prepn. of urea moiety-contg. peptide derivs. as neutral endopeptidase and angiotensin converting enzyme inhibitors)  
 IT 9015-82-1, Angiotensin converting enzyme 82707-54-8, Neutral endopeptidase  
 RL: BSU (Biological study, unclassified); MSC (Miscellaneous); BIOL (Biological study)  
 (prepn. of urea moiety-contg. peptide derivs. as neutral endopeptidase and angiotensin converting enzyme inhibitors)  
 IT 75-36-5, Acetyl chloride 95-56-7, 2-Bromophenol 100-39-0, Benzyl bromide 109-73-9, Butylamine, reactions 123-72-8, Butylaldehyde

123-75-1, Pyrrolidine, reactions 617-52-7, Dimethyl itaconate  
 2812-46-6 5419-55-6, Triisopropyl borate 6165-69-1, 3-Thienylboronic  
 acid 7693-46-1 9000-92-4, Amylase 24424-99-5, Di-tert-butyl  
 dicarbonate 28310-65-8 30418-59-8, 3-Aminophenylboronic acid  
 32315-10-9, Triphosgene 63024-30-6 71989-31-6 118786-32-6

**191426-93-4** 191426-94-5

RL: RCT (Reactant)

(prepn. of urea moiety-contg. peptide derivs. as neutral endopeptidase  
 and angiotensin converting enzyme inhibitors)

IT 6793-92-6P 31575-75-4P 53087-13-1P 78887-39-5P 118602-51-0P  
 118602-52-1P 137255-87-9P 137255-91-5P 146631-00-7P 150351-64-7P  
 156682-54-1P 190661-29-1P 191425-25-9P **191425-26-0P**  
 191425-27-1P 191425-28-2P 191425-29-3P 191425-30-6P 191425-31-7P  
 191425-32-8P 191425-33-9P 191425-34-0P 191425-35-1P 191425-36-2P  
**191425-37-3P** 191425-38-4P 191425-39-5P 191425-40-8P  
 191425-41-9P 191425-42-0P 191425-43-1P 191425-44-2P 191425-45-3P  
 191425-46-4P 191425-47-5P 191425-48-6P 191425-49-7P 191425-50-0P  
 191425-51-1P 191425-52-2P 191425-53-3P 191425-54-4P  
**191425-55-5P 191425-56-6P 191425-57-7P**  
**191425-58-8P 191425-59-9P 191425-60-2P**  
 191425-61-3P 191425-62-4P 191425-63-5P 191425-64-6P 191425-65-7P  
 191425-66-8P 191425-67-9P 191425-68-0P 191425-69-1P 191425-70-4P  
 191425-71-5P 191425-72-6P 191425-73-7P 191425-74-8P 191425-75-9P  
 191425-77-1P 191425-79-3P 191425-81-7P 191425-83-9P 191425-85-1P  
 191425-87-3P 191425-88-4P 191425-90-8P 191425-92-0P 191425-94-2P  
 191425-95-3P 191425-97-5P 191426-01-4P 191426-03-6P 191426-05-8P  
 191426-07-0P 191426-09-2P 191426-11-6P 191426-13-8P 191426-15-0P  
 191426-16-1P 191426-17-2P 191426-18-3P 191426-21-8P 191426-23-0P  
**191426-26-3P 191426-27-4P 191426-29-6P**  
**191426-31-0P** 191426-33-2P 191426-35-4P 191426-37-6P  
 191426-39-8P 191426-40-1P 191426-41-2P 191426-42-3P 191426-43-4P  
 191426-44-5P 191426-45-6P 191426-46-7P 191426-47-8P 191426-48-9P  
 191426-49-0P 191426-50-3P 191426-51-4P 191426-52-5P 191426-53-6P  
 191426-54-7P 191426-55-8P 191426-56-9P 191426-57-0P 191426-58-1P  
 191426-59-2P 191426-60-5P 191426-61-6P 191426-62-7P 191426-63-8P  
 191426-64-9P 191426-67-2P 191426-68-3P 191426-69-4P 191426-70-7P  
 191426-71-8P 191426-72-9P 191426-73-0P 191426-74-1P  
**191426-75-2P 191426-76-3P 191426-77-4P**  
**191426-78-5P** 191426-80-9P 191426-81-0P 191426-82-1P  
 191426-83-2P 191426-84-3P 191426-85-4P 191426-86-5P  
**191426-87-6P** 191426-88-7P 191426-89-8P 191426-90-1P  
 191426-91-2P 191426-92-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(prepn. of urea moiety-contg. peptide derivs. as neutral endopeptidase  
 and angiotensin converting enzyme inhibitors)

IT **191426-93-4**


RL: RCT (Reactant)

(prepn. of urea moiety-contg. peptide derivs. as neutral endopeptidase  
 and angiotensin converting enzyme inhibitors)

RN 191426-93-4 HCAPLUS

CN L-Tyrosine, O-(1,1-dimethylethyl)-N-[(4-nitrophenoxy)carbonyl]-,  
 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 191425-26-0P 191425-37-3P 191425-55-5P  
 191425-56-6P 191425-57-7P 191425-58-8P  
 191425-59-9P 191425-60-2P 191426-26-3P  
 191426-27-4P 191426-29-6P 191426-31-0P  
 191426-75-2P 191426-76-3P 191426-77-4P  
 191426-78-5P 191426-87-6P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of urea moiety-contg. peptide derivs. as neutral endopeptidase  
 and angiotensin converting enzyme inhibitors)

RN 191425-26-0 HCAPLUS

CN [1,1'-Biphenyl]-4-propanoic acid, .alpha.-[[[4-nitrophenoxy)carbonyl]amino]-, methyl ester, (S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 191425-37-3 HCAPLUS

CN L-Phenylalanine, N-[(4-nitrophenoxy)carbonyl]-4-(3-thienyl)-, methyl ester  
 (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 191425-55-5 HCAPLUS


CN L-Phenylalanine, N-[(4-nitrophenoxy)carbonyl]-, 1,1-dimethylethyl ester  
 (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 191425-56-6 HCAPLUS  
 CN 2-Benzothiazolepropanoic acid, .alpha.-[[ (4-nitrophenoxy) carbonyl] amino]-, methyl ester, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 191425-57-7 HCAPLUS  
 CN L-Tyrosine, O-(1-methylethyl)-N-[[ (4-nitrophenoxy) carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 191425-58-8 HCAPLUS  
 CN 2-Naphthalenepropanoic acid, .alpha.-[[ (4-nitrophenoxy) carbonyl] amino]-, methyl ester, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 191425-59-9 HCAPLUS  
 CN 1,3-Benzodioxole-5-propanoic acid, .alpha.-[[ (4-nitrophenoxy) carbonyl] amino]-, methyl ester, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 191425-60-2 HCAPLUS  
 CN L-Tyrosine, N-[(4-nitrophenoxy) carbonyl]-O-phenyl-, methyl ester (9CI)  
 (CA INDEX NAME)

Absolute stereochemistry.



RN 191426-26-3 HCAPLUS  
 CN [1,1'-Biphenyl]-4-propanoic acid, 4'-fluoro-.alpha.-[[ (4-nitrophenoxy) carbonyl] amino]-, methyl ester, (S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 191426-27-4 HCAPLUS  
 CN [1,1'-Biphenyl]-4-propanoic acid, .alpha.-[[ (4-nitrophenoxy) carbonyl] amino]-4'-(trifluoromethyl)-, methyl ester, (S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.






RN 191426-29-6 HCAPLUS  
 CN [1,1'-Biphenyl]-4-propanoic acid, 4'-methyl-.alpha.-[[[4-nitrophenoxy)carbonyl]amino]-, methyl ester, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 191426-31-0 HCAPLUS  
 CN L-Serine, O-[(4-fluorophenyl)methyl]-N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 191426-75-2 HCAPLUS  
 CN [1,1'-Biphenyl]-4-propanoic acid, .alpha.-[[[4-nitrophenoxy)carbonyl]amino]-2'-(phenylmethoxy)-, methyl ester, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 191426-76-3 HCAPLUS  
CN [1,1'-Biphenyl]-4-propanoic acid, .alpha.-[[[4-nitrophenoxy)carbonyl]amino]-3'-(phenylmethoxy)-, methyl ester, (S)- (9CI)  
(CA INDEX NAME)

Absolute stereochemistry.




RN 191426-77-4 HCAPLUS  
CN [1,1'-Biphenyl]-4-propanoic acid, .alpha.-[[[4-nitrophenoxy)carbonyl]amino]-4'-(phenylmethoxy)-, methyl ester, (S)- (9CI)  
(CA INDEX NAME)

Absolute stereochemistry.




RN 191426-78-5 HCAPLUS  
CN [1,1'-Biphenyl]-4-propanoic acid, 3'-(acetylamino)-.alpha.-[[[4-nitrophenoxy)carbonyl]amino]-, methyl ester, (S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 191426-87-6 HCAPLUS  
CN L-Lysine, N2-[(4-nitrophenoxy)carbonyl]-N6-[(phenylmethoxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.




L27 ANSWER 8 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1996:637443 HCAPLUS  
 DN 125:329473  
 TI Preparation of aminediol-containing peptide analogs as retroviral protease inhibitors  
 IN Gordon, Eric M.; Barrish, Joel C.; Bisacchi, Gregory S.; Sun, Chong-qing; Tino, Joseph A.; Vite, Gregory D.; Zahler, Robert  
 PA E. R. Squibb & Sons, Inc., USA  
 SO U.S., 219 pp. Cont.-in-part of U.S. Ser. No. 927,027, abandoned.  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 IC ICM C07D401-12  
 NCL 552303000  
 CC 34-3 (Amino Acids, Peptides, and Proteins)  
 Section cross-reference(s): 1

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 5559256	A	19960924	US 1993-79978	19930625
	AU 9341659	A1	19940127	AU 1993-41659	19930630
	AU 677194	B2	19970417		
	HU 67090	A2	19950130	HU 1993-2080	19930719
	CA 2100894	AA	19940121	CA 1993-2100894	19930720
	NO 9302620	A	19940121	NO 1993-2620	19930720
	EP 580402	A2	19940126	EP 1993-305691	19930720
	EP 580402	A3	19970305		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
	ZA 9305243	A	19940217	ZA 1993-5243	19930720
	CN 1085546	A	19940420	CN 1993-108954	19930720
	JP 06206857	A2	19940726	JP 1993-201016	19930720
	US 5760036	A	19980602	US 1995-455295	19950531
	US 5776933	A	19980707	US 1995-456125	19950531
PRAI	US 1992-916916		19920720		
	US 1992-927027		19920806		
	US 1993-79978		19930625		
OS	MARPAT 125:329473				
GI					



I



AB Aa-E-NR8CHR9H(OH)CH2NHCH2CH(OH)CHR9NR8-E-Ab [Aa, Ab = H, alkyl, R3C(:Z), R3SO2, R3R4NSO2, R3R4NC(:Z), R3SC(:O), R5R6R7COC(:Z); E = a single bond or a peptide chain contg. 1 to 4 amino acids, the N-terminus of which is bonded to Aa or Ab; R3, R4 = H, alkyl, aryl, carbocyclyl; R5, R6, R7 = H, alkyl, aryl, carbocyclyl, fluorenyl, alkynyl, alkenyl; R5, R6, and R7 may, independently, be joined together with the carbon atom to which they are bonded, to form a mono-, bi- or tricyclic carbocyclic ring system; R8 = H, alkyl; R9 = arylalkyl; Z = O, S; wherein: wherever they appear alone or as part of another group, unless otherwise indicated, the terms "alk." or "alkyl" denote a straight or branched chain satd. radical contg. 1 to 12 carbons in the normal chain, optionally substituted by one or more groups selected from (un)protected OH, oxo (with the proviso that the carbon bearing the oxo group is not adjacent to a heteroatom), CO2H, halo, alkoxy, aryloxy, alkoxy carbonyl, etc.] or salts thereof, which inhibit retroviral protease and are particularly useful in the treatment and/or prevention of HIV infection (AIDS), are prepd. Thus, bis(3-amino-2-hydroxy-4-phenylbutyl)amine deriv. (I; R = H) was condensed with L-tert-leucine deriv. (HO-Q) using 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride and HOBT in DMF/CH2CH2 at 0.degree. to room temp. to give the title compd. I (R = Q). The latter compd. at 10 .mu.M in vitro inhibited 99% HIV protease and showed IC50 of 0.012 .mu.M which was the concn. of drug that increased the formazan prodn. in CEM-SS cells infected with the RF strain of HIV to 50% of that produced by uninfected cells in the absence of drug.

ST aminediol contg peptide analog prepn; retroviral protease inhibitor; HIV infection AIDS treatment

IT Acquired immune deficiency syndrome  
Virucides and Virustats

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of aminediol-contg. peptide analogs as retroviral protease inhibitors for treatment of HIV infection (AIDS))

IT Peptides, preparation

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(analog, prepn. of aminediol-contg. peptide analogs as retroviral protease inhibitors for treatment of HIV infection (AIDS))

IT Virus, animal

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(human immunodeficiency, prepn. of aminediol-contg. peptide analogs as retroviral protease inhibitors for treatment of HIV infection (AIDS))

IT 272-01-5P, Furo[2,3-b]pyridine 609-71-2P 2508-01-2P 3356-88-5P  
3694-86-8P 7423-92-9P 13031-76-0P 15833-82-6P 15833-84-8P  
22455-69-2P 27038-48-8P, Furo[2,3-b]pyridin-3(2H)-one 41036-01-5P

52532-02-2P 62030-47-1P 83096-36-0P 97024-23-2P 100841-12-1P  
 109274-92-2P, Furo[2,3-b]pyridine-2-carboxaldehyde 116005-23-3P  
 144731-95-3P 161302-38-1P 161302-39-2P 161302-40-5P 162537-26-0P  
 162537-37-3P 162537-75-9P 162537-76-0P 162537-77-1P 162537-78-2P  
 162537-79-3P 162537-80-6P 162537-81-7P 162537-82-8P,  
 Furo[2,3-b]pyridine-2-methanol 162537-83-9P 162537-84-0P  
 162537-86-2P 162537-87-3P 162537-88-4P 162537-89-5P 162537-91-9P  
 162537-95-3P 162537-96-4P 162537-97-5P 162537-98-6P 162537-99-7P  
 162538-00-3P 162538-01-4P 162538-02-5P 162538-08-1P 162538-10-5P  
 162538-11-6P 162538-12-7P 162538-13-8P 162538-14-9P 162538-15-0P  
 162538-18-3P 162538-19-4P 162538-20-7P 162538-21-8P 162538-22-9P  
 162538-23-0P 162538-24-1P 162538-25-2P 162538-28-5P 162538-29-6P  
 162538-31-0P 162538-33-2P 162538-37-6P 162538-38-7P 162538-39-8P  
 162538-40-1P 162538-42-3P 162538-45-6P 162538-47-8P 162538-48-9P  
 162538-50-3P 162538-51-4P 162538-52-5P 162538-53-6P 162538-54-7P  
 162538-55-8P 162538-57-0P 162538-58-1P 162538-59-2P 162538-60-5P  
 162538-61-6P 162538-63-8P 162538-64-9P 162538-65-0P 162538-66-1P  
 162538-67-2P 162538-69-4P 162538-70-7P 162538-71-8P 162538-72-9P  
 162538-73-0P 162538-74-1P 162538-76-3P 162538-80-9P 162538-81-0P  
 162538-83-2P 162538-84-3P 162538-85-4P 162538-90-1P 162538-91-2P  
 162538-92-3P 162538-93-4P 162538-94-5P 162538-95-6P 162538-96-7P  
 162538-97-8P 162539-02-8P 162539-03-9P 162539-05-1P 162539-07-3P  
 162539-09-5P 162539-13-1P 162539-15-3P 162539-17-5P 162539-18-6P  
 162539-19-7P 162539-20-0P 162539-21-1P 162539-22-2P 162539-23-3P  
 162539-25-5P 162539-27-7P 162539-29-9P 162539-32-4P 162539-33-5P  
 162539-34-6P 162539-35-7P 162539-36-8P 162539-37-9P 162539-38-0P  
 162539-39-1P 162539-41-5P 162539-43-7P 162539-44-8P 162539-45-9P  
 162539-46-0P 162539-47-1P 162539-48-2P 162539-50-6P 162539-54-0P  
 162539-57-3P 162539-58-4P 162539-60-8P 162539-61-9P 162539-62-0P  
 162539-63-1P 162539-64-2P 162539-65-3P 162539-66-4P 162539-67-5P  
 162539-68-6P 162539-69-7P 162539-70-0P 162539-71-1P 162539-72-2P  
 162539-73-3P 162539-74-4P 162539-75-5P 162539-76-6P 162539-80-2P  
 162539-81-3P 162539-85-7P 162539-86-8P 162539-87-9P 162539-88-0P  
 162539-89-1P 162539-90-4P 162539-91-5P 162539-92-6P 162539-93-7P  
 162539-94-8P 162539-95-9P 162539-96-0P 162539-97-1P 162539-98-2P  
 162539-99-3P 162540-00-3P 162540-01-4P 162540-03-6P 162540-04-7P  
 162540-05-8P 162540-06-9P 162540-07-0P 162540-08-1P 162540-09-2P  
 162540-10-5P 162540-11-6P 162540-12-7P 162540-13-8P 162540-14-9P  
 162540-15-0P 162540-17-2P 162540-18-3P 162540-19-4P 162540-20-7P  
 162540-21-8P 162540-22-9P 162540-23-0P 162540-24-1P 162540-25-2P  
 162540-26-3P 162540-27-4P 162540-28-5P 162540-29-6P 162540-30-9P  
 162540-31-0P 162540-32-1P 162540-33-2P 162540-34-3P 162540-35-4P  
 162540-36-5P 162540-37-6P 162540-38-7P 162540-39-8P 162540-40-1P  
 162540-41-2P 162540-42-3P 162540-43-4P 162540-44-5P 162540-45-6P  
 162540-46-7P 162540-47-8P 162540-48-9P 162540-49-0P 162540-50-3P  
 162540-51-4P 162540-52-5P 162540-53-6P 162540-54-7P 162540-55-8P

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of aminediol-contg. peptide analogs as retroviral protease inhibitors for treatment of HIV infection (AIDS).)

IT	162540-56-9P	162540-57-0P	162540-58-1P	162540-59-2P	162540-60-5P
	162540-61-6P	162540-63-8P	162540-65-0P	162540-66-1P	162540-67-2P
	162540-68-3P	162540-69-4P	162540-70-7P	162540-71-8P	162540-72-9P
	162540-73-0P	162540-74-1P	162540-75-2P	162540-76-3P	162540-77-4P
	162540-78-5P	162540-79-6P	162540-80-9P	162540-81-0P	162540-82-1P
	162540-83-2P	162540-84-3P	162540-85-4P	162540-86-5P	162540-87-6P
	162540-88-7P	162540-89-8P	162540-90-1P	162540-91-2P	162540-92-3P
	162540-93-4P	162540-94-5P	162540-95-6P	162540-96-7P	162540-97-8P
	162540-98-9P	162540-99-0P	162541-00-6P	162541-01-7P	162541-02-8P

162541-03-9P	162541-04-0P	162541-05-1P	162541-06-2P	162541-07-3P
162541-08-4P	162541-09-5P	162541-11-9P	162541-12-0P	162541-13-1P
162541-14-2P	162541-15-3P	162541-16-4P	162541-17-5P	162541-18-6P
162541-19-7P	162541-20-0P	162541-21-1P	162541-22-2P	162541-23-3P
162541-24-4P	162541-25-5P	162541-97-1P	162541-98-2P	162541-99-3P
162542-00-9P	162542-01-0P	162542-02-1P	162542-03-2P	162542-05-4P
162542-08-7P	162542-09-8P	162542-10-1P	162677-24-9P	162677-29-4P
162677-30-7P	162677-32-9P	162677-33-0P	162677-34-1P	162677-35-2P
162677-36-3P	162677-37-4P	162677-38-5P	162677-39-6P	162677-40-9P
162677-42-1P	162677-45-4P	162677-48-7P	162677-50-1P	162677-52-3P
162677-54-5P	162677-56-7P	162677-58-9P	162677-59-0P	162677-60-3P
162677-61-4P	162677-62-5P	162677-63-6P	162677-64-7P	162677-65-8P
162677-66-9P	162677-69-2P	162677-70-5P	162677-71-6P	162677-72-7P
162677-73-8P	162677-74-9P	162677-75-0P	162677-76-1P	162677-77-2P
162677-78-3P	162677-79-4P	162677-80-7P	162677-81-8P	162677-82-9P
162677-83-0P	162677-84-1P	162677-85-2P	162677-86-3P	162677-87-4P
162677-88-5P	162677-89-6P	162677-90-9P	162677-92-1P	162677-93-2P
162677-94-3P	162677-95-4P	162677-96-5P	162677-97-6P	162677-98-7P
162677-99-8P	162678-00-4P	162678-01-5P	162678-02-6P	162678-03-7P
162678-04-8P	162678-05-9P	162678-06-0P	162678-07-1P	162678-08-2P
162678-09-3P	162678-10-6P	162678-12-8P	162678-13-9P	162678-14-0P
162678-15-1P	162678-16-2P	162678-17-3P	162678-18-4P	162678-23-1P
162678-25-3P	162678-31-1P	162678-33-3P	162678-34-4P	162678-38-8P
170996-47-1P	170996-48-2P	171228-69-6P	175233-62-2P	175417-50-2P
175417-51-3P	183161-02-6P	183161-31-1P	183161-35-5P	183162-38-1P
183162-39-2P	183162-40-5P	183162-41-6P	183162-42-7P	183162-44-9P
183162-45-0P	183162-49-4P	183162-51-8P	183162-58-5P	183162-62-1P
183162-63-2P	183162-64-3P	183162-67-6P	183162-69-8P	183162-75-6P
183162-77-8P	183162-79-0P	183255-85-8P	183255-86-9P	183255-88-1P
183255-89-2P	183255-90-5P	183255-92-7P	183255-93-8P	183255-94-9P
183256-02-2P				

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of aminediol-contg. peptide analogs as retroviral protease inhibitors for treatment of HIV infection (AIDS))

IT 144114-21-6, Retropepsin

RL: BPR (Biological process); BSU (Biological study, unclassified); MSC (Miscellaneous); BIOL (Biological study); PROC (Process)

(prepn. of aminediol-contg. peptide analogs as retroviral protease inhibitors for treatment of HIV infection (AIDS))

IT 50-00-0, Formaldehyde, reactions 52-52-8, 1-Aminocyclopentanecarboxylic acid 56-12-2, 4-Aminobutyric acid, reactions 64-18-6, Formic acid, reactions 68-12-2, Dimethylformamide, reactions 70-25-7 72-18-4, L-Valine, reactions 74-88-4, Methyl iodide, reactions 74-89-5, Methylamine, reactions 75-16-1, Methylmagnesium bromide 75-36-5, Acetyl chloride 75-44-5, Phosgene 75-66-1, tert-Butyl mercaptan 75-98-9, Trimethylacetic acid 76-83-5, Triphenylmethyl chloride 77-76-9, 2,2-Dimethoxypropane 79-14-1, reactions 79-22-1, Methyl chloroformate 79-44-7, Dimethylcarbonyl chloride 79-50-5, DL-Pantolactone 83-33-0, 1-Indanone 91-62-3, 6-Methylquinoline 93-10-7, Quinaldic acid 95-54-5, o-Phenylenediamine, reactions 95-55-6, o-Aminophenol 96-49-1, Ethylene carbonate 98-59-9, Tosyl chloride 98-80-6, Phenylboronic acid 98-88-4, Benzoyl chloride 100-39-0, Benzyl bromide 100-44-7, Benzyl chloride, reactions 100-46-9, Benzylamine, reactions 100-52-7, Benzaldehyde, reactions 100-86-7, .alpha.,.alpha.-Dimethylphenethyl alcohol 103-74-2, 2-(2-Hydroxyethyl)pyridine 105-36-2, Ethyl bromoacetate 108-23-6, Isopropyl chloroformate 108-24-7, Acetic anhydride 109-00-2, 3-Hydroxypyridine 109-86-4, 2-Methoxyethanol 110-06-5, tert-Butyl

disulfide 110-15-6, Butanedioic acid, reactions 110-91-8, Morpholine, reactions 111-42-2, reactions 119-67-5, 2-Carboxybenzaldehyde 122-59-8, Phenoxyacetic acid 122-98-5, 2-Anilinoethanol 122-99-6, 2-Phenoxyethanol 137-07-5, 2-Aminothiophenol 288-32-4, Imidazole, reactions 335-08-0, 1,1,1-Trifluoroacetone cyanohydrin 353-80-0 358-23-6, Triflic anhydride 453-20-3, 3-Hydroxytetrahydrofuran 473-85-8, 1,4-Anhydro-D-threitol 500-22-1, 3-Pyridinecarboxaldehyde 501-53-1, Benzyl chloroformate 503-38-8, Trichloromethyl chloroformate 534-03-2, 2-Amino-1,3-propanediol 539-74-2, Ethyl 3-bromopropionate 540-51-2, 2-Bromoethanol 541-47-9, 3,3-Dimethylacrylic acid 558-30-5, Isobutylene oxide 586-98-1, 2-Pyridylcarbinol 591-80-0, 4-Pentenoic acid 593-56-6, Methoxyamine hydrochloride 594-56-9, 2,3,3-Trimethylbutene 598-21-0, Bromoacetyl bromide 611-71-2 617-35-6, Ethyl pyruvate 617-94-7, Dimethylphenyl carbinol 622-08-2, 2-Benzyloxyethanol 622-40-2, 4-(2-Hydroxyethyl)morpholine 623-08-5, N-Methyl-p-toluidine 624-83-9, Methyl isocyanate 625-38-7, 3-Butenoic acid 627-18-9 628-41-1, 1,4-Cyclohexadiene 630-19-3, Pivalaldehyde 644-36-0, o-Tolylacetic acid 670-95-1, 4-Phenylimidazole 672-15-1, L-Homo-serine 677-22-5, tert-Butylmagnesium chloride 687-47-8, (S)-Ethyl lactate 693-89-0, 1-Methylcyclopentene 759-24-0, Diethyl tert-butylmalonate 775-06-4, DL-Meta-tyrosine 821-09-0, 4-Penten-1-ol 917-54-4, Methyllithium 937-14-4, m-Chloroperbenzoic acid 1003-04-9 1070-83-3, tert-Butylacetic acid 1120-87-2, 4-Bromopyridine 1122-62-9, 2-Acetylpyridine 1142-20-7 1145-80-8 1148-11-4 1149-26-4 1161-13-3 1193-47-1, 2,2-Dimethylcyclohexanone 1462-03-9, 1-Methyl-1-cyclopentanol 1609-86-5, tert-Butyl isocyanate 1664-54-6, 3-Amino-3-phenylpropionic acid 1779-49-3, Methyltriphenylphosphonium bromide 1826-67-1, Vinylmagnesium bromide 2018-66-8 2130-96-3 2212-75-1 2370-61-8 2976-75-2, (1-Naphthoxy)acetic acid 2987-16-8, 3,3-Dimethylbutyraldehyde 3160-59-6 3173-56-6, Benzyl isocyanate 3240-94-6, 4-(2-Chloroethyl)morpholine 3262-72-4 3587-60-8, Benzyl chloromethyl ether 3731-51-9, 2-(Aminomethyl)pyridine 4436-24-2, Benzyloxirane 4530-20-5 4541-32-6, 2,2-Dimethylcyclopentanone 4857-04-9, 2-(Chloromethyl)benzimidazole 5034-06-0, Trimethylsulfoxonium chloride 5333-74-4, Ethyl tert-butylglyoxylate 5470-11-1, Hydroxylamine hydrochloride 6278-91-7, 4-Benzyloxy-2-butanone 6290-49-9, Methyl methoxyacetate 6306-52-1, L-Valine methyl ester hydrochloride 6351-10-6, 1-Indanol 6829-40-9, Diethyl aminomalonate 7326-19-4, D-Phenyllactic acid 7364-25-2, Indazolinone 7432-21-5 7486-35-3, Vinyltributyltin 7536-55-2 7677-24-9, Trimethylsilyl cyanide 7693-46-1, p-Nitrophenyl chloroformate 10147-11-2, 3-Phenyl-1-propyne 13031-04-4 13139-15-6 13139-16-7 13139-17-8, N-Benzyloxycarbonyloxy succinimide 13329-18-5, 5-Benzyloxy-2-pentanone 13570-08-6, 2-Benzimidazoleacetic acid 13575-16-1, Ethyl 5-Phenyloxazole-2-carboxylate 13734-34-4 13734-41-3 14347-78-5, (R)-2,2-Dimethyl-1,3-dioxolane-4-methanol 14397-64-9, 1-Ethoxycarbonyl-2-indanone 15761-39-4 16520-62-0, 4-Phenyl-1-butyne 16677-29-5 17199-29-0 17392-83-5, (R)-Methyl lactate 17463-43-3, DL-3,3,3-Trifluoroalanine 18162-48-6, tert-Butyldimethylsilyl chloride 18942-49-9 19575-07-6, Methyl quinaldate 19728-63-3, Z-Thr-OH 19752-84-2, 3-Hydroxytetrahydropyran 19810-31-2, Benzyloxyacetyl chloride 20117-47-9, 1-Methyl-1-cyclobutanol 20160-60-5, 2-Trimethylsilylethyl chloroformate 20412-38-8, Neopentyl chloroformate 20662-89-9, 4-Phenyloxazole 20859-02-3, L-tert-Leucine 21641-92-9 22146-57-2 22323-82-6, (S)-2,2-Dimethyl-1,3-dioxolane-4-methanol 24424-99-5, Di-tert-butyl dicarbonate 26628-22-8, Sodium azide 26782-71-8, D-tert-Leucine 28920-43-6, 9-Fluorenylmethyl chloroformate 28954-12-3, L-Allothreonine 29943-42-8, Tetrahydro-4H-pyran-4-one 30525-89-4, Paraformaldehyde 32366-02-2, N-Benzyl-N-methyl carbamoyl chloride 36024-28-9 37595-74-7, N-Phenyltriflimide 37736-82-6,

N-tert-Butoxycarbonyl-L-cyclohexylalanine 40299-87-4,  
 4-(Bromoacetyl)morpholine 41242-94-8, 2-Hydroxymethyl quinoxaline  
 52373-72-5 53333-76-9, 2,2-Dimethyl-1-propanesulfonyl chloride  
 58632-95-4, Boc-ON 59562-82-2 60456-21-5 67478-50-6 68835-89-2,  
 Di-tert-amyl dicarbonate 69739-34-0, tert-Butyldimethylsilyl triflate  
 76513-69-4, 2-(Trimethylsilyl)ethoxymethyl chloride 78879-20-6  
 80360-23-2 85613-64-5 86087-23-2, (S)-(+)-3-Hydroxytetrahydrofuran  
 106167-47-9 107202-43-7 112372-06-2, Furo[2,3-c]pyridine-2-  
 carboxaldehyde 127862-89-9 162537-72-6, Furo[2,3-c]pyridine-2-methanol  
 162537-73-7 162541-63-1 162678-30-0 162870-63-5

RL: RCT (Reactant)

(prepn. of aminediol-contg. peptide analogs as retroviral protease  
 inhibitors for treatment of HIV infection (AIDS))

IT 95-13-6P, 1H-Indene 102-14-7P 111-32-0P 272-62-8P,  
 Furo[3,2-b]pyridine 334-88-3P, Diazomethane 374-35-6P 558-43-0P  
 587-33-7P 815-17-8P 1184-93-6P 1191-31-7P 1615-14-1P,  
 1H-Imidazole-1-ethanol 1780-17-2P, 2-Quinolinemethanol 1796-25-4P  
 2215-63-6P 2258-42-6P, Acetic formic anhydride 2280-28-6P 2644-82-8P  
 2842-44-6P 2849-93-6P, 1H-Benzimidazole-2-carboxylic acid 3587-64-2P  
 3724-55-8P 4026-20-4P 4113-04-6P, 6-Quinolinedicarboxaldehyde  
 4441-30-9P, 4-Morpholinepropanol 4647-42-1P 4647-43-2P 4754-27-2P  
 4856-97-7P, 1H-Benzimidazole-2-methanol 5105-78-2P 5367-24-8P  
 6970-72-5P 7467-35-8P 7525-64-6P 7748-36-9P, 3-Oxetanol  
 13737-35-4P 14477-66-8P 14598-96-0P 15546-08-4P 17450-34-9P  
 18096-68-9P, 1H-Indene-2-methanol 19458-29-8P 19539-50-5P,  
 Furo[2,3-c]pyridine 20120-24-5P 20361-09-5P 22892-29-1P  
 22929-52-8P 23249-97-0P, 1H-Benzimidazole-2-propanoic acid 24580-44-7P  
 24621-70-3P, 1H-Indole-2-methanol 25854-85-7P 25854-87-9P  
 30293-86-8P 31562-43-3P 33905-47-4P 34637-40-6P 35677-88-4P  
 37535-57-2P 37859-42-0P, 2-Benzothiazolemethanol 39497-64-8P  
 40594-83-0P 42417-65-2P 50411-26-2P 50531-59-4P 51110-97-5P,  
 2-Benzoxazolepropanol 53346-03-5P 56365-70-9P 57443-39-7P  
 59524-02-6P 60398-41-6P 60651-97-0P 62565-28-0P 62965-10-0P  
 64360-69-6P 66866-64-6P 67706-63-2P 70448-03-2P 73282-11-8P  
 77186-95-9P, 2-Benzoxazolemethanol 80466-51-9P 85328-36-5P  
 85951-09-3P 85995-53-5P 86096-65-3P 86562-71-2P 88246-12-2P  
 89464-59-5P 90819-30-0P 91968-72-8P 94882-74-3P 98737-29-2P  
 98760-08-8P 98955-64-7P 98997-01-4P 100516-88-9P,  
 6-Quinolinemethanol 100868-72-2P 102123-74-0P 102123-85-3P  
 102152-03-4P 102229-10-7P 102831-44-7P 104948-22-3P 106513-42-2P  
 108957-20-6P 112372-05-1P, Furo[3,2-b]pyridine-2-carboxaldehyde  
 113247-51-1P 113459-50-0P 114645-18-0P 115916-75-1P 127041-02-5P  
 127382-65-4P 128018-44-0P 131424-20-9P 134807-06-0P 134807-20-8P  
 134807-28-6P 134807-29-7P 134807-30-0P 137515-66-3P 138432-95-8P  
 141978-97-4P 143372-45-6P 143372-46-7P 143372-47-8P 143576-95-8P  
 143688-65-7P 144186-00-5P 144186-52-7P 144825-44-5P 149357-61-9P  
 153291-20-4P 154117-17-6P 154612-75-6P 156474-21-4P 156474-22-5P  
 159259-43-5P 160232-54-2P 162125-34-0P 162536-40-5P 162536-41-6P  
 162536-42-7P 162536-43-8P 162536-44-9P 162536-45-0P 162536-46-1P  
 162536-47-2P 162536-48-3P 162536-50-7P 162536-54-1P 162536-55-2P  
 162536-56-3P 162536-57-4P 162536-58-5P 162536-59-6P 162536-60-9P  
 162536-63-2P 162536-64-3P 162536-65-4P 162536-67-6P 162536-68-7P  
 162536-69-8P 162536-70-1P 162536-71-2P 162536-72-3P 162536-73-4P  
 162536-74-5P 162536-77-8P 162536-78-9P 162536-79-0P 162536-80-3P  
 162536-81-4P 162536-82-5P 162536-83-6P 162536-84-7P 162536-85-8P  
 162536-86-9P 162536-87-0P 162536-88-1P 162536-89-2P 162536-90-5P  
 162536-91-6P 162536-92-7P 162536-93-8P 162536-94-9P 162536-95-0P  
 162536-96-1P 162536-97-2P 162536-98-3P 162536-99-4P 162537-00-0P  
 162537-01-1P 162537-03-3P 162537-10-2P 162537-11-3P  
 162537-12-4P 162537-13-5P 162537-14-6P 162537-15-7P 162537-16-8P



162537-17-9P	162537-20-4P	162537-21-5P	162537-22-6P	162537-27-1P
162537-31-7P	162537-32-8P	162537-33-9P	162537-34-0P	162537-35-1P
162537-36-2P	162537-39-5P	162537-40-8P	162537-41-9P	162537-42-0P
162537-43-1P	162537-44-2P	162537-45-3P	162537-46-4P	162537-47-5P
162537-48-6P	162537-49-7P	162537-50-0P	162537-51-1P	162537-53-3P
162537-54-4P	162537-55-5P	162537-56-6P	162537-61-3P,	
Furo[3,2-b]pyridine-2-methanol		162537-62-4P	162537-63-5P	
162537-64-6P	162537-65-7P	162537-66-8P	162537-67-9P	162537-68-0P
162537-69-1P	162537-70-4P	162537-74-8P		

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of aminediol-contg. peptide analogs as retroviral protease  
 inhibitors for treatment of HIV infection (AIDS))

IT	162538-06-9P	162538-07-0P	162538-09-2P	162538-16-1P	162538-17-2P
	162538-30-9P	162538-32-1P	162538-34-3P	162538-36-5P	162538-41-2P
	162538-43-4P	162538-44-5P	162538-46-7P	162538-49-0P	162538-56-9P
	162538-68-3P	162538-77-4P	162538-82-1P	162538-86-5P	162538-98-9P
	162538-99-0P	162539-00-6P	162539-01-7P	162539-04-0P	162539-06-2P
	162539-08-4P	162539-11-9P	162539-12-0P	162539-14-2P	162539-16-4P
	162539-24-4P	162539-26-6P	162539-28-8P	162539-30-2P	162539-31-3P
	162539-40-4P	162539-42-6P	162539-51-7P	162539-52-8P	162539-53-9P
	162539-55-1P	162539-56-2P	162539-77-7P	162539-78-8P	162539-79-9P
	162540-62-7P	162541-27-7P	162541-28-8P	162541-29-9P	162541-30-2P
	162541-31-3P	162541-32-4P	162541-33-5P	162541-34-6P	162541-35-7P
	162541-38-0P	162541-39-1P	162541-40-4P	162541-41-5P	162541-42-6P
	162541-43-7P	162541-44-8P	162541-45-9P	162541-46-0P	162541-48-2P
	162541-49-3P	162541-50-6P	162541-52-8P	162541-53-9P	162541-54-0P
	162541-55-1P	162541-57-3P	162541-58-4P	162541-60-8P	162541-61-9P
	162541-62-0P	162541-64-2P	162541-65-3P	162541-66-4P	162541-67-5P
	162541-68-6P	162541-69-7P	162541-70-0P	162541-71-1P	162541-72-2P
	162541-73-3P	162541-74-4P	162541-75-5P	162541-76-6P	162541-77-7P
	162541-78-8P	162541-79-9P	162541-80-2P	162541-81-3P	162541-82-4P
	162541-84-6P	162541-85-7P	162541-86-8P	162541-87-9P	162541-88-0P
	162541-89-1P	162541-92-6P	162541-94-8P	162541-95-9P	162541-96-0P
	162677-20-5P	162677-23-8P	162677-31-8P	162677-41-0P	162677-43-2P
	162677-46-5P	162677-47-6P	162677-49-8P	162677-51-2P	162677-53-4P
	162677-55-6P	162677-57-8P	162677-91-0P	162678-21-9P	162678-22-0P
	162678-26-4P	162678-29-7P	162678-36-6P	162678-37-7P	162776-41-2P
	165524-61-8P	165727-45-7P	170996-45-9P	171230-81-2P	175390-83-7P
	183161-49-1P	183161-50-4P	183161-56-0P	183161-57-1P	183161-58-2P
	183161-59-3P	183161-60-6P	183161-61-7P	183161-62-8P	183161-63-9P
	183161-64-0P	183161-65-1P	183161-67-3P	183161-68-4P	183161-69-5P
	183161-71-9P	183161-72-0P	183161-73-1P	183161-74-2P	183161-75-3P
	183161-76-4P	183161-77-5P	183161-78-6P	183161-81-1P	183161-82-2P
	183161-83-3P	183161-84-4P	183161-85-5P	183161-86-6P	183161-87-7P
	183161-89-9P	183161-90-2P	183161-91-3P	183161-92-4P	183161-93-5P
	183161-94-6P	183161-95-7P	183161-96-8P	183161-97-9P	183161-98-0P
	183161-99-1P	183162-00-7P	183162-01-8P	183162-02-9P	183162-03-0P
	183162-04-1P	183162-05-2P	183162-06-3P	183162-07-4P	183162-08-5P
	183162-09-6P	183162-10-9P	183162-11-0P	183162-12-1P	183162-13-2P
	183162-14-3P	183162-15-4P	183162-16-5P	183162-17-6P	183162-18-7P
	183162-19-8P	183162-20-1P	183162-21-2P	183162-22-3P	183162-23-4P
	183162-24-5P	183162-25-6P	183162-26-7P	183162-27-8P	183162-28-9P
	183162-29-0P	183162-30-3P	183162-32-5P	183162-33-6P	183162-34-7P
	183162-35-8P	183162-36-9P	183162-37-0P	183255-95-0P	183255-96-1P
	183255-97-2P	183255-98-3P	183255-99-4P	183256-00-0P	183256-01-1P
	183256-04-4P	183256-05-5P	183256-06-6P	183256-07-7P	183256-08-8P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of aminediol-contg. peptide analogs as retroviral protease  
 inhibitors for treatment of HIV infection (AIDS))


IT 162537-10-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of aminediol-contg. peptide analogs as retroviral protease  
inhibitors for treatment of HIV infection (AIDS))

RN 162537-10-2 HCAPLUS

CN L-Valine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX  
NAME)

Absolute stereochemistry.



L27 ANSWER 9 OF 24 HCAPLUS COPYRIGHT 2001 ACS

AN 1996:551382 HCAPLUS

DN 125:196378

TI Nitrogen mustard prodrugs with novel lipophilic protecting groups, and  
processes for their production

IN Springer, Caroline Joy; Niculescu-Duvaz, Ion

PA Cancer Research Campaign Technology Limited, UK

SO PCT Int. Appl., 44 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM C07C275-32

ICS C07C271-54; C07C219-34; C07C271-28; A61K031-17; A61K031-27

CC 34-2 (Amino Acids, Peptides, and  
Proteins)

Section cross-reference(s): 1, 25, 63

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9622277	A1	19960725	WO 1996-GB112	19960119
	W:	AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI			
	RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE			
	CA 2210347	AA	19960725	CA 1996-2210347	19960119
	AU 9644535	A1	19960807	AU 1996-44535	19960119
	AU 709251	B2	19990826		
	EP 804413	A1	19971105	EP 1996-900627	19960119
	EP 804413	B1	20000517		
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE			
	JP 10512565	T2	19981202	JP 1996-522127	19960119
	AT 193012	E	20000615	AT 1996-900627	19960119
	ES 2149445	T3	20001101	ES 1996-900627	19960119
	US 6005002	A	19991221	US 1997-875099	19970716
PRAI	GB 1995-1052	A	19950119		
	WO 1996-GB112	W	19960119		
OS	MARPAT 125:196378				
GI					

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

- AB The invention provides compds. I and II [X, Y = Cl, Br, iodo, CH<sub>3</sub>SO<sub>3</sub>, or OSO<sub>2</sub>Ph (wherein Ph is optionally substituted by 1-5 alkyl, halo, cyano, and/or nitro); R<sub>1</sub>, R<sub>2</sub> each = 1-4 optional substituents; Z<sub>1</sub>, Z<sub>2</sub> = O, NH; R<sub>3</sub> = H, t-Bu, allyl; Z<sub>3</sub> = hydrocarbyl group such as carboxyethyl, optionally contg. heteroatoms] and their physiol. acceptable derivs. The compds. can be converted in situ into nitrogen mustard agents by the actions of enzymes such as carboxypeptidase or nitroreductase and are useful for the treatment of cancer (no data). For example, the glutamate ester L-Me<sub>3</sub>COCOCH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>CMe<sub>3</sub> [III; R = NH<sub>2</sub>] was converted to the isocyanate III [R = NCO], which reacted with 4-hydroxybenzaldehyde to give 72% III [R = NHCO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>CHO-4]. This was reduced with NaBH<sub>3</sub>CN to give 86% of III [R = NHCO<sub>2</sub>C<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>OH-4], which was coupled with the nitrogen mustard 4-(OCN)C<sub>6</sub>H<sub>4</sub>N(CH<sub>2</sub>CH<sub>2</sub>Cl)<sub>2</sub> in 90% yield, followed by deesterification with formic acid (87%), to give title compd. IV.
- ST glutamate nitrogen mustard prodrug antitumor prepn
- IT Neoplasm inhibitors  
(prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)
- IT Amino acids, preparation  
Nitrogen mustards  
RL: BAC (Biological activity or effector, except adverse); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)
- IT Therapeutics  
(pharmaco-, GDEPT (gene-directed enzyme prodrug therapy); prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)
- IT Pharmaceutical dosage forms  
(prodrugs, prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)
- IT 9031-98-5, Carboxypeptidase 9037-41-6, Nitroreductase  
RL: BSU (Biological study, unclassified); MSC (Miscellaneous); BIOL (Biological study)  
(activation by; prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)
- IT 18483-99-3P, 4-Nitrobenzyl 2-tetrahydropyranyl ether 18484-05-4P, 4-Aminobenzyl 2-tetrahydropyranyl ether 113068-95-4P, 4-Isocyanatobenzyl tert-butyldiphenylsilyl ether 161803-03-8P, 4-Nitrobenzyl tert-butyldiphenylsilyl ether 161803-04-9P, 4-Aminobenzyl tert-butyldiphenylsilyl ether 161803-05-0P 161803-06-1P 180839-06-9P, 4-[N,N-Bis(2-chloroethyl)amino]phenyl trimethylsilyl ether 180839-08-1P 180839-11-6P 180839-12-7P 180839-13-8P 180839-14-9P 180839-15-0P 180839-16-1P 180839-17-2P 180839-18-3P 180839-19-4P 180839-20-7P 180839-21-8P 180841-62-7P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(intermediate; prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)
- IT 180838-97-5P 180839-00-3P 180839-02-5P 180839-04-7P  
RL: BAC (Biological activity or effector, except adverse); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)
- IT 180838-98-6P 180839-01-4P 180839-03-6P 180839-05-8P

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)

IT 110-87-2 123-08-0, 4-Hydroxybenzaldehyde 619-73-8, 4-Nitrobenzyl alcohol 623-04-1, 4-Aminobenzyl alcohol 623-05-2, 4-Hydroxybenzyl alcohol 1204-69-9, 4-[N,N-Bis(2-chloroethyl)amino]phenol 7693-46-1, 4-Nitrophenyl chloroformate 20845-16-3 32677-01-3, Di-tert-butyl L-glutamate hydrochloride 57529-05-2, 4-(1,3-Dithian-2-yl)phenol 58479-61-1, tert-Butyldiphenylchlorosilane 82484-59-1, 4-[N,N-Bis(2-chloroethyl)amino]phenyl isocyanate

RL: RCT (Reactant)

(starting material; prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)

IT 180839-20-7P 180839-21-8P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(intermediate; prepn. of lipophilic glutamate-based nitrogen mustard prodrugs as anticancer agents)

RN 180839-20-7 HCAPLUS

CN L-Glutamic acid, N-[[4-[[[(4-nitrophenoxy)carbonyl]oxy]methyl]phenoxy]carbonyl]-, bis(1,1-dimethylethyl) ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 180839-21-8 HCAPLUS

CN L-Glutamic acid, N-[[4-[[[(4-nitrophenoxy)carbonyl]oxy]methyl]phenoxy]carbonyl]-, di-2-propenyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.

PAGE 1-A




CH<sub>2</sub>

L27 ANSWER 10 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1996:509810 HCAPLUS  
 DN 125:168650  
 TI High-purity N-[4-[N,N-bis(2-iodoethyl)amino]phenoxy-carbonyl]-L-glutamic acid and hydrogen iodide salt as prodrugs for ADEPT therapy  
 IN Heaton, David William; Dines, Susan; Dowell, Robert Ian  
 PA Zeneca Limited, UK  
 SO PCT Int. Appl., 18 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM C07C271-54  
 ICS A61K031-325  
 CC 34-2 (Amino Acids, Peptides, and Proteins)  
 Section cross-reference(s): 1, 63

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9620169	A1	19960704	WO 1995-GB2997	19951221
W: AL, AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
ZA 9510936	A	19960624	ZA 1995-10936	19951221
CA 2204741	AA	19960704	CA 1995-2204741	19951221
AU 9642700	A1	19960719	AU 1996-42700	19951221
AU 702337	B2	19990218		
GB 2309031	A1	19970716	GB 1997-8977	19951221
GB 2309031	B2	19980812		
EP 799193	A1	19971008	EP 1995-941219	19951221
EP 799193	B1	20000816		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE				
CN 1171100	A	19980121	CN 1995-196993	19951221
HU 77288	A2	19980330	HU 1997-2207	19951221
BR 9510463	A	19980609	BR 1995-10463	19951221
AT 195511	E	20000915	AT 1995-941219	19951221
FI 9702630	A	19970618	FI 1997-2630	19970618
NO 9702883	A	19970620	NO 1997-2883	19970620
US 5981791	A	19991109	US 1997-860880	19970623
PRAI GB 1994-26133	A	19941223		
WO 1995-GB2997	W	19951221		
OS MARPAT 125:168650				
GI				



- AB A new salt of the prodrug N-[4-[N,N-bis(2-iodoethyl)amino]phenoxy]carbonyl-L-glutamic acid (I), which is useful in antibody directed enzyme prodrug therapy (ADEPT), is disclosed. This salt, the hydrogen iodide salt I.HI, is obtained in cryst. form with m.p. 142-145.degree. and has a specified X-ray powder diffraction spectrum. Prepn. of cryst. I.HI enables I to be prepd. in a highly pure form. I.HI also has advantages of good thermal stability, easy synthesis, and reduced hygroscopic/deliquescence properties. Improved stability may be due to reversal of degrading of the mustard alkylating arms of the mol. by HI. A multi-step prepn. of I from L-glutamic acid and p-O<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>OCOC<sub>2</sub>H<sub>5</sub> is given. Mouse tumor xenograft expts. using therapy with F(ab')<sub>2</sub>ASB7-CPG2 conjugate (2.5 mg/kg i.v.) and I.HI (3 doses at 70 mg/kg i.p. each) gave tumor regression and >30 day growth delays with only minor effects on body wt. and peripheral white blood cell counts.
- ST iodoethylaminophenoxy carbonyl glutamic acid hydriodide antitumor prodrug ADEPT; bisiodoethylaminophenoxy carbonyl glutamic acid prepn antitumor prodrug ADEPT
- IT Neoplasm inhibitors  
(prepn. of high-purity [[bis(iodoethyl)amino]phenoxy carbonyl] glutamic acid and hydriodide salt as antitumor prodrugs for ADEPT)
- IT Pharmaceutical dosage forms  
(prodrugs, antibody-directed enzyme prodrug therapy (ADEPT); prepn. of high-purity [[bis(iodoethyl)amino]phenoxy carbonyl] glutamic acid and hydriodide salt as antitumor prodrugs for ADEPT)
- IT Pharmaceutical dosage forms  
(prodrugs, prepn. of high-purity [[bis(iodoethyl)amino]phenoxy carbonyl] glutamic acid and hydriodide salt as antitumor prodrugs for ADEPT)
- IT 5269-43-2P, L-Glutamic acid bis(trimethylsilyl) ester **156079-00-4P**  
156079-01-5P 156079-89-9P 172974-17-3P 180031-70-3P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(intermediate; prepn. of high-purity [[bis(iodoethyl)amino]phenoxy carbonyl] glutamic acid and hydriodide salt as antitumor prodrugs for ADEPT)
- IT 180031-69-0P  
RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or effector, except adverse); IMF (Industrial manufacture); PRP (Properties); PUR (Purification or recovery); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(prepn. of high-purity [[bis(iodoethyl)amino]phenoxy carbonyl] glutamic acid and hydriodide salt as antitumor prodrugs for ADEPT)
- IT 156079-88-8P  
RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or effector, except adverse); IMF (Industrial manufacture); PUR (Purification or recovery); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(prepn. of high-purity [[bis(iodoethyl)amino]phenoxy carbonyl] glutamic acid and hydriodide salt as antitumor prodrugs for ADEPT)


IT 56-86-0, L-Glutamic acid, reactions 75-21-8, Ethylene oxide, reactions 75-77-4, Chlorotrimethylsilane, reactions 115-11-7, Isobutylene, reactions 7693-46-1, p-Nitrophenyl chloroformate  
 RL: RCT (Reactant)  
 (starting material; prepn. of high-purity [[bis(iodoethyl)amino]phenoxy carbonyl]glutamic acid and hydriodide salt as antitumor prodrugs for ADEPT)

IT **156079-00-4P**  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (intermediate; prepn. of high-purity [[bis(iodoethyl)amino]phenoxy carbonyl]glutamic acid and hydriodide salt as antitumor prodrugs for ADEPT)

RN 156079-00-4 HCAPLUS

CN L-Glutamic acid, N-[(4-nitrophenoxy)carbonyl]-, bis(1,1-dimethylethyl) ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L27 ANSWER 11 OF 24 HCAPLUS COPYRIGHT 2001 ACS

AN 1996:171803 HCAPLUS

DN 124:233139

TI Preparation of sulfonylamino acid amides containing cis-epoxide as irreversible HIV protease inhibitors

IN Yoon, Heungsik; Choy, Nakyeon; Kim, Sung Chun; Choi, Ho II; Son, Young Chan; Park, Chi Hyo; Moon, Kwang-Yul; Jung, Wonhee; Kim, Chung Ryeol; et al.

PA IG Chemical Ltd., S. Korea

SO Eur. Pat. Appl., 58 pp.  
 CODEN: EPXXDW

DT Patent

LA English


IC ICM C07D303-36  
 ICS C07D303-40; C07D405-12; C07D407-12; A61K031-335; A61K031-435

CC 34-2 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 1

FAN.CNT 7

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 687675	A2	19951220	EP 1995-108908	19950609
	EP 687675	A3	19960306		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
	KR 125117	B1	19971205	KR 1994-13423	19940615
	JP 08193077	A2	19960730	JP 1995-172733	19950615
	JP 2987313	B2	19991206		
PRAI	KR 1994-13423	A	19940615		
OS	MARPAT 124:233139				
GI					



AB Novel cis-epoxide compds. [I; R1, R2 = H, alkyl; R3 = (un)substituted aryl or alkyl; R4 = H, C1-4 alkyl; n = 0,1,2; A = (X)(Y)mR5, NR6R7, ZCHR8R9; wherein X = CO, COCO, CO, SO2, CS; Y = O, CH2, NH, NMe; m = 0,1; R5 = heterocyclyl, straight or branched or cyclic C1-8 alkyl or alkoxy, heterocyclylalkyl, cycloalkylalkyl, arylalkoxy; R6 = straight or branched C1-8 alkyl, cycloalkyl, cycloalkylalkyl; R7 = H, alkyl; Z = O, NH, NMe; R8, R9 = alkyl optionally substituted by arom. hydrocarbyl or cycloalkyl, C3-8 cycloalkyl, aryl], useful for treating or preventing diseases caused by HIV infection, are prepd. The novel HIV protease inhibitor I has a specific structure to form a stable bonding with the enzyme active site, which entails a highly enhanced irreversible inhibition against HIV protease. An anti-AIDS or immunomodulator contains a therapeutically effective amt. of said cis-epoxide I. Thus, (S)-5-[(N-benzyloxycarbonyl)amino]-6-phenyl-(cis)-3-hexene-1-carboxylic acid was condensed with (S)-2-amino-3-methyl-1-phenylbutane using N-ethyl-N'-(3-(dimethylamino)propyl)carbodiimide hydrochloride (EDC) and HOBT in DMF followed by epoxidn. with m-chloroperbenzoic acid in CH2Cl2 to give the cis-epoxide (II; R = PhCH2O2C), which was hydrogenolyzed in the presence of 10% Pd-C in MeOH under an atm. of H2 coupled with N-benzyloxycarbonyl-.beta.-(S-methyl)-L-valine using EDC and HOBT in DMF, and oxidized with m-chloroperbenzoic acid in CH2Cl2 to give the title compd. II (R = Q). The latter compd. in vitro inhibited HIV protease with the inhibition const. Kina/Ki min-1M-1 109-1010 (Kina = a rate const. indicating rate of chem. reaction forming covalent bond between an enzyme and an inhibitor in Michaelis-Menten complex; Ki = an inhibition const. indicating the disocn. rate of Michaelis-Menten complex into an enzyme and an inhibitor) and in vitro showed IC50 of 1 nM for inhibiting the HIV-1 infection of H9 or Sup T1 cell lines.

ST cis epoxide prepn HIV protease inhibitor; irreversible HIV protease inhibitor; sulfonylamino acid amide contg cis epoxide

IT Acquired immune deficiency syndrome  
Virucides and Virustats

(prepn. of sulfonylamino acid amides contg. cis-epoxide as irreversible HIV protease inhibitors for treating AIDS)

IT Amides, preparation

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(amino, prepn. of sulfonylamino acid amides contg. cis-epoxide as irreversible HIV protease inhibitors for treating AIDS)

IT Epoxides

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic



preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(cis-, prepn. of sulfonylamino acid amides contg. cis-epoxide as irreversible HIV protease inhibitors for treating AIDS)

IT Virus, animal  
(human immunodeficiency 1, prepn. of sulfonylamino acid amides contg. cis-epoxide as irreversible HIV protease inhibitors for treating AIDS)

IT 174562-29-9P 174562-30-2P 174562-31-3P 174562-32-4P 174562-33-5P  
174562-34-6P 174562-35-7P 174562-36-8P 174562-37-9P 174562-38-0P  
174562-39-1P 174562-40-4P 174562-41-5P 174562-42-6P 174562-43-7P  
174562-44-8P 174562-45-9P 174562-46-0P 174562-47-1P 174562-48-2P  
174562-49-3P 174562-50-6P 174562-51-7P 174562-52-8P 174562-53-9P  
174562-54-0P 174562-55-1P 174562-56-2P 174562-57-3P 174562-58-4P  
174562-59-5P 174562-60-8P 174562-61-9P 174562-62-0P 174562-63-1P  
174562-64-2P 174562-65-3P 174562-66-4P 174562-67-5P 174562-68-6P  
174562-69-7P 174562-70-0P 174562-71-1P 174562-72-2P 174562-73-3P  
174562-74-4P 174562-75-5P 174562-76-6P 174562-77-7P 174562-78-8P  
174562-79-9P 174562-80-2P 174562-81-3P

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of sulfonylamino acid amides contg. cis-epoxide as irreversible HIV protease inhibitors for treating AIDS)

IT 144114-21-6, Retropepsin  
RL: BPR (Biological process); BIOL (Biological study); PROC (Process)  
(prepn. of sulfonylamino acid amides contg. cis-epoxide as irreversible HIV protease inhibitors for treating AIDS)

IT 59-67-6, 3-Pyridinecarboxylic acid, reactions 74-88-4, Iodomethane, reactions 78-77-3, Isobutyl bromide 78-82-0, Isobutyronitrile 88-14-2, 2-Furancarboxylic acid 96-41-3, Cyclopentyl alcohol 98-00-0, 2-Furanmethanol 98-59-9, Toluenesulfonyl chloride 98-98-6, 2-Pyridinecarboxylic acid 100-39-0, Benzyl bromide 100-46-9, Benzylamine, reactions 100-55-0, 3-Pyridylcarbinol 110-68-9, Methyl-N-butylamine 501-53-1, Benzyl chloroformate 503-74-2, Isovaleric acid 527-72-0, 2-Thiophenic acid 574-98-1, N-(2-Bromoethyl)phthalimide 586-95-8, 4-Pyridylcarbinol 586-98-1, 2-Pyridylcarbinol 603-35-0, Triphenylphosphine, reactions 617-89-0, 2-Furanylmethylamine 625-45-6, Methoxyacetic acid 1113-41-3, L-Penicillamine 1779-49-3, Methyltriphenylphosphonium bromide 2516-33-8, Cyclopropylmethanol 2516-47-4, Cyclopropylmethylamine 4083-57-2, 3-Amino-2,4-dimethylpentane 6921-34-2, Benzylmagnesium chloride 7693-46-1, p-Nitrophenyl chloroformate 13734-34-4, 23844-66-8, (R)-1-Amino-2-methyl-1-phenylpropane 24424-99-5, Di-tert-Butyl dicarbonate 24939-24-0, p-Aminobenzenesulfonyl chloride 33445-07-7, Isopropoxyacetic acid 37222-66-5, Oxone 59830-60-3, N-Benzyloxycarbonyl-L-phenylalaninal 68906-26-3, (S)-1-Amino-2-methyl-1-phenylpropane 69492-74-6, Thiopheneacetic acid 74124-79-1, Disuccinimidyl carbonate 96928-87-9 111491-96-4 137867-58-4, Furanacetic acid 174562-82-4

RL: RCT (Reactant)

(prepn. of sulfonylamino acid amides contg. cis-epoxide as irreversible HIV protease inhibitors for treating AIDS)


IT 65273-64-5P 82894-53-9P 97589-56-5P 100217-05-8P 112898-22-3P  
156641-79-1P 156641-80-4P 156641-81-5P 156641-82-6P 156641-83-7P  
156715-06-9P 160742-44-9P 160742-45-0P 174562-83-5P 174562-84-6P  
174562-85-7P 174562-86-8P 174562-87-9P 174562-88-0P 174562-89-1P  
174562-90-4P 174562-91-5P 174562-92-6P 174562-93-7P  
174562-94-8P 174689-86-2P 174689-87-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(prepn. of sulfonylamino acid amides contg. cis-epoxide as irreversible

HIV protease inhibitors for treating AIDS)  
 IT 174562-91-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of sulfonylamino acid amides contg. cis-epoxide as irreversible  
 HIV protease inhibitors for treating AIDS)  
 RN 174562-91-5 HCAPLUS  
 CN L-Valine, 3-(methylthio)-N-[(4-nitrophenoxy)carbonyl]-, phenylmethyl ester  
 (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L27 ANSWER 12 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1995:994162 HCAPLUS  
 DN 124:87790  
 TI Pharmaceutical compositions containing HIV protease inhibitors and their  
 preparation.  
 IN Al-Razzak, Laman; Marsh, Kennan C.; Manning, Lourdes P.; Kaul, Dilip  
 PA Abbott Laboratories, USA  
 SO PCT Int. Appl., 58 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM A61K031-425  
 ICS A61K009-08; A61K047-10; A61K047-12; A61K047-14  
 CC 34-2 (Amino Acids, Peptides, and  
 Proteins)  
 Section cross-reference(s): 63

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9520384	A1	19950803	WO 1995-US232	19950103
W: AU, CA, JP, KR, MX				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
IL 111991	A1	20000726	IL 1994-111991	19941215
CA 2178632	AA	19950803	CA 1995-2178632	19950103
AU 9515248	A1	19950815	AU 1995-15248	19950103
AU 700942	B2	19990114		
EP 732923	A1	19960925	EP 1995-906790	19950103
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
JP 09508383	T2	19970826	JP 1995-520059	19950103
US 5484801	A	19960116	US 1995-440277	19950512
PRAI US 1994-189021	A	19940128		
US 1994-283239	A	19940729		
WO 1995-US232	W	19950103		


GI




- AB A pharmaceutical compn. which comprises a soln. of an HIV protease inhibiting compd. (e.g., I) in a pharmaceutically acceptable org. solvent comprising a mixt. of (1): (a) a solvent selected from propylene glycol and polyethylene glycol or (b) a solvent selected from polyoxyethyleneglycerol triricinoleate, polyethylene glycol 40 hydrogenated castor oil, fractioned coconut oil, polyoxyethylene 20 sorbitan monooleate and 2-(2-ethoxyethoxy)ethanol or (c) a mixt. thereof; and (2) ethanol or propylene glycol, is claimed. I was prepd. in many steps and its bioavailability in various formulations was studied.
- ST hiv protease inhibitor pharmaceutical compn; valinyldiphenylhexane prepn  
hiv protease inhibitor
- IT Pharmaceutical dosage forms  
(pharmaceutical compns. contg. HIV protease inhibitors and their prepn.)
- IT 155213-67-5P  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(pharmaceutical compns. contg. HIV protease inhibitors and their prepn.)
- IT 143838-10-2P 144164-10-3P  
RL: BYP (Byproduct); PREP (Preparation)  
(pharmaceutical compns. contg. HIV protease inhibitors and their prepn.)
- IT 62-56-6, Thiourea, reactions 75-12-7, Formamide, reactions 105-39-5, Ethyl chloroacetate 109-94-4, Ethyl formate 534-07-6, 1,3-Dichloroacetone 563-83-7, Isobutyramide 4070-48-8, Valine methyl ester 6372-14-1, N-Benzyloxycarbonylphenylalaninol 40635-67-4, .alpha.-Acetoxyisobutyryl bromide 156732-13-7 156732-15-9  
RL: RCT (Reactant)  
(pharmaceutical compns. contg. HIV protease inhibitors and their prepn.)
- IT 115-08-2P, Thioformamide 13515-65-6P 32955-21-8P, 2-Amino-5-(ethoxycarbonyl)thiazole 32955-22-9P, Ethyl thiazole-5-carboxylate 33142-21-1P, Ethyl 2-chloro-2-formylacetate 38585-74-9P, 5-Hydroxymethylthiazole 59830-60-3P 65386-28-9P 137649-69-5P 144141-68-4P 144163-44-0P 144163-85-9P 144163-97-3P 144164-11-4P 154212-59-6P 154212-60-9P 154212-61-0P 154248-99-4P 162537-10-2P 162849-92-5P 162849-93-6P 162849-94-7P 162849-95-8P 162849-96-9P, 2-Amino-5-(ethoxycarbonyl)thiazole hydrochloride 162990-03-6P 165315-39-9P  
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(pharmaceutical compns. contg. HIV protease inhibitors and their prepn.)
- IT 162990-01-4  
RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
(pharmaceutical compns. contg. HIV protease inhibitors and their prepn.)

prepn.)  
 IT 162537-10-2P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (pharmaceutical comps. contg HIV protease inhibitors and their  
 prepn.)  
 RN 162537-10-2 HCAPLUS  
 CN L-Valine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX  
 NAME)

Absolute stereochemistry.



L27 ANSWER 13 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1995:965043 HCAPLUS  
 DN 124:117909  
 TI Optimization of Alkylating Agent Prodrugs Derived from Phenol and Aniline  
 Mustards: A New Clinical Candidate Prodrug (ZD2767) for Antibody-Directed  
 Enzyme Prodrug Therapy  
 AU Springer, Caroline J.; Dowell, Robert; Burke, Philip J.; Hadley, Elma;  
 Davies, D. Huw; Blakey, David C.; Melton, Roger G.; Niculescu-Duvaz, Ion  
 CS Cancer Research Campaign Centre for Cancer Therapeutics, Institute of  
 Cancer Research, Sutton, SM2 5NG, UK  
 SO J. Med. Chem. (1995), 38(26), 5051-65  
 CODEN: JMCMAR; ISSN: 0022-2623  
 DT Journal  
 LA English  
 CC 34-2 (Amino Acids, Peptides, and  
 Proteins)  
 Section cross-reference(s): 1  
 GI



AB Sixteen novel potential prodrugs I [R = H, 2-Me, 2-Cl, 3-Me, 3-Me2CH, 3-F, 2,3-(CH:CHCH:CH), 3-CN; Z = O, NH; X, Y = Cl, Br, iodo, O3SMe] derived from phenol or aniline mustards and their 16 corresponding drugs II with ring substitution and/or different alkylating functionalities were designed. They are bifunctional alkylating agents in which the activating effect of the phenolic hydroxyl or amino function is masked through an oxycarbonyl or a carbamoyl bond to a glutamic acid. These prodrugs were

designed to be activated to their corresponding phenol and aniline nitrogen mustard drugs at a tumor site by prior administration of a monoclonal antibody conjugated to the bacterial enzyme carboxypeptidase G2 (CPG2) in antibody-directed enzyme prodrug therapy (ADEPT). The synthesis of the analogous novel parent drugs II is also described. The viability of a colorectal cell line (LoVo) was monitored with the potential prodrugs and the parent drugs. The differential in the cytotoxicity between the potential prodrugs and their corresponding active drugs ranged between 12 and >195 fold. Some compds. I exhibited substantial prodrug activity, since a cytotoxicity differential of >100 was achieved compared to the analogous II. The ability of the potential prodrugs to act as substrates for CPG2 was detd. (kinetic parameters KM and kcat), and the chem. stability was measured for all the compds. The unsubstituted phenols with different alkylating functionalities (I; R = H, Z = O) proved to have the highest ratio of substrates kcat:KM. From these studies, III (ZD2767) emerges as a new ADEPT clin. trial candidate due to its physicochem. and biol. characteristics.

ST ZD 2767 prepn biol alkylating agent; glutamate aniline mustard prepn ADEPT; phenol glutamate mustard prepn ADEPT; structure activity alkylating agent prodrug

IT Neoplasm inhibitors  
(optimization of alkylating agent prodrugs derived from phenol and aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)

IT Alkylating agents, biological  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(optimization of alkylating agent prodrugs derived from phenol and aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)

IT Antibodies  
Enzymes  
RL: BPR (Biological process); BIOL (Biological study); PROC (Process)  
(optimization of alkylating agent prodrugs derived from phenol and aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)

IT Molecular structure-biological activity relationship  
(cytotoxic, optimization of alkylating agent prodrugs derived from phenol and aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)

IT Pharmaceutical dosage forms  
(prodrugs, optimization of alkylating agent prodrugs derived from phenol and aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)

IT 21667-05-0P 156078-84-1P 156079-29-7P 156079-30-0P 156079-31-1P  
156079-32-2P 156079-34-4P 156079-35-5P 172974-00-4P 172974-01-5P  
172974-02-6P 172974-03-7P 172974-19-5P 172974-20-8P 172974-21-9P  
172974-22-0P

RL: BAC (Biological activity or effector, except adverse); RCT (Reactant);  
SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(optimization of alkylating agent prodrugs derived from phenol and aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)

IT 156078-82-9P 156078-91-0P 156078-94-3P 156078-98-7P 156079-02-6P  
156079-03-7P 156079-04-8P 156079-05-9P 156079-06-0P 156079-07-1P  
156079-56-0P 156079-57-1P 156079-88-8P, ZD 2767 156079-91-3P  
172974-23-1P 172974-24-2P

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(optimization of alkylating agent prodrugs derived from phenol and

aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)

IT 75-21-8, Oxirane, reactions 111-42-2, reactions 350-30-1,  
3-Chloro-4-fluoronitrobenzene 369-34-6, 3,4-Difluoronitrobenzene  
399-96-2, 4-Amino-2-fluorophenol 2791-84-6 2834-90-4,  
4-Amino-1-naphthol 2835-96-3, 4-Amino-2-methylphenol 2835-99-6,  
4-Amino-3-methylphenol 3964-52-1, 4-Amino-2-chlorophenol 5854-73-9  
7693-46-1, 4-Nitrophenyl chloroformate 17417-09-3, 2-Fluoro-5-  
nitrobenzonitrile 17609-80-2, 4-Amino-3-chlorophenol 32677-01-3,  
Di-tert-butyl glutamate hydrochloride 82774-61-6 156639-14-4  
172974-26-4

RL: RCT (Reactant)

(optimization of alkylating agent prodrugs derived from phenol and aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)

IT 65976-57-0P 65976-66-1P 156078-83-0P 156078-85-2P 156078-86-3P  
156078-92-1P 156078-93-2P 156078-95-4P 156078-99-8P  
**156079-00-4P** 156079-01-5P 156079-08-2P 156079-09-3P  
156079-10-6P 156079-11-7P 156079-12-8P 156079-13-9P 156079-15-1P  
156079-16-2P 156079-17-3P 156079-18-4P 156079-19-5P 156079-20-8P  
156079-22-0P 156079-23-1P 156079-24-2P 156079-25-3P 156079-27-5P  
156079-28-6P 156079-36-6P 156079-37-7P 156079-38-8P 156079-39-9P  
156079-40-2P 156079-41-3P 156079-58-2P 156079-59-3P 156079-60-6P  
156079-61-7P 156079-65-1P 156079-66-2P 156079-89-9P 156079-90-2P  
156079-92-4P 156639-24-6P 156639-26-8P 172973-90-9P 172973-91-0P  
172973-92-1P 172973-93-2P 172973-94-3P 172973-95-4P 172973-96-5P  
172973-97-6P 172973-98-7P 172973-99-8P 172974-04-8P 172974-05-9P  
172974-06-0P 172974-07-1P 172974-08-2P 172974-09-3P 172974-10-6P  
172974-11-7P 172974-12-8P 172974-13-9P 172974-14-0P 172974-15-1P  
**172974-16-2P** 172974-17-3P 172974-18-4P 172974-25-3P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(optimization of alkylating agent prodrugs derived from phenol and aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)


IT **156079-00-4P 172974-16-2P**

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(optimization of alkylating agent prodrugs derived from phenol and aniline mustards in prepn. of prodrug ZD2767 for antibody-directed enzyme prodrug therapy)

RN 156079-00-4 HCAPLUS

CN L-Glutamic acid, N-[(4-nitrophenoxy)carbonyl]-, bis(1,1-dimethylethyl) ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 172974-16-2 HCAPLUS

CN L-Glutamic acid, N-[(3-fluoro-4-nitrophenoxy)carbonyl]-, bis(1,1-dimethylethyl) ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L27 ANSWER 14 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1995:958521 HCAPLUS  
 DN 124:176946  
 TI Preparation of retroviral protease inhibiting peptide analogs.  
 IN Norbeck, Daniel W.; Sham, Hing L.; Kempf, Dale J.; Zhao, Chen  
 PA Abbott Laboratories, USA  
 SO U.S., 66 pp. Cont.-in-part of U.S. Ser. No. 23,226, abandoned.  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 IC ICM A61K031-44  
 ICS A61K031-425; A61K031-42; C07D413-14; C07D417-14; C07D263-30;  
 C07D277-20

NCL 514333000

CC 34-3 (Amino Acids, Peptides, and  
 Proteins)

Section cross-reference(s): 1

FAN.CNT 2

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 5461067	A	19951024	US 1994-185666	19940201
	WO 9419332	A1	19940901	WO 1994-US1457	19940208
	W: CA, JP				
	RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
	CA 2155338	AA	19940901	CA 1994-2155338	19940208
	EP 683772	A1	19951129	EP 1994-908018	19940208
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
	JP 08507061	T2	19960730	JP 1994-519025	19940208
	US 5621109	A	19970415	US 1995-455922	19950531
	US 5631376	A	19970520	US 1995-455458	19950531
	US 5990135	A	19991123	US 1995-455052	19950531
PRAI	US 1993-23226		19930225		
	US 1994-185666		19940201		
	WO 1994-US1457		19940208		
OS	MARPAT 124:176946				
AB	R6YmNHCHR1CH(OH)CH2NR2NH(Y1)nR5 [R1, R2 = H, alkyl, aryl, thioalkoxyalkyl, aralkyl, cycloalkyl, guanidinoalkyl, arylthioalkoxyalkyl, cycloalkyloxyalkyl, cycloalkylsulfonylalkyl, aminocarbonylalkyl, etc.; Y = NHCHR4CO, NHR4CO, etc.; Y1 = COCHR3NH, CONR3NH, etc.; R5, R6 = C(:T)GR7; T = O, S; G = CH2, O, S; NR8; R8 = H, alkyl, cycloalkyl; R7 = alkyl, cycloalkyl, aryl, arylalkyl, arylalkoxyalkyl, aminoalkyl, N-protected aminoalkyl, alkylaminoalkyl, N-protected alkylaminoalkyl, dialkylaminoalkyl, carboxyalkoxyalkyl, (alkoxycarbonyl)alkoxyalkyl, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, N-protected aminocarbonylalkyl, alkylaminocarbonylalkyl, etc.; m, n = 0, 1], were prepd. Thus, (5S)-[[[(5-thiazolyl)methoxy]carbonyl]amino-2-[(N-2-oxazolyl)methoxycarbonyl]amino-4S-hydroxy-1-(3-furanyl)-6-phenyl-2-azahexane (soln. phase prepn. given) inhibited HIV-13B in MT4 cells with IC50 = 0.029-0.032 .mu.M.				
ST	peptide analog prepn retroviral protease inhibitor; aminopropylhydrazine				

azolylmethoxycarbonyl amino acid prepn virucide

IT Virucides and Virustats  
(prepn. of retroviral protease inhibiting peptide analogs)

IT Peptides, preparation  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(analog, prep. of retroviral protease inhibiting peptide analogs)

IT Virus, animal  
(human immunodeficiency 1, treatment of HIV infections; prep. of retroviral protease inhibiting peptide analogs)

IT 144114-21-6, Retropepsin  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(HIV protease inhibitors; prep. of retroviral protease inhibiting peptide analogs)

IT 150767-06-9P 150767-07-0P 162739-20-0P 162739-21-1P 162739-22-2P  
162739-23-3P 162739-24-4P 162739-25-5P 162739-26-6P 162739-27-7P  
162739-28-8P 162739-31-3P 162739-32-4P 162739-33-5P 162739-38-0P  
162739-39-1P 162739-40-4P 162739-41-5P 162739-42-6P 162739-43-7P  
162739-46-0P 173767-16-3P 173767-17-4P 173767-18-5P 173767-19-6P  
173767-20-9P 173767-21-0P 173767-22-1P 173767-23-2P 173767-24-3P  
173767-25-4P 173767-26-5P 173767-27-6P 173767-28-7P 173767-29-8P  
173767-30-1P 173767-31-2P 173767-32-3P 173767-33-4P 173767-34-5P  
173767-35-6P 173767-36-7P 173767-37-8P 173767-38-9P 173767-39-0P  
173767-40-3P 173767-41-4P 173767-42-5P 173767-43-6P 173767-44-7P  
173767-45-8P 173767-46-9P 173767-47-0P 173767-48-1P 173767-49-2P  
173767-50-5P 173767-51-6P 173767-52-7P 173767-53-8P 173767-54-9P  
173767-55-0P 173767-56-1P 173767-57-2P 173767-58-3P 173767-59-4P  
173767-60-7P 173767-61-8P 173767-62-9P 173767-63-0P 173767-64-1P  
173767-65-2P 173767-66-3P 173767-67-4P 173767-68-5P 173767-69-6P  
173767-70-9P 173767-71-0P 173767-72-1P 173767-73-2P 173767-74-3P  
173767-75-4P 173767-76-5P 173767-77-6P 173767-78-7P 173767-79-8P  
173767-80-1P 173767-81-2P 173767-82-3P 173767-83-4P 173767-84-5P  
173767-85-6P 173767-86-7P 173767-87-8P 173767-88-9P 173767-89-0P  
173767-90-3P 173767-91-4P 173767-92-5P 173767-93-6P 173767-94-7P  
173767-95-8P 173767-96-9P 173767-97-0P 173767-98-1P 173767-99-2P  
173768-00-8P 173768-01-9P 173768-02-0P 173768-03-1P 173768-04-2P  
173768-05-3P 173768-06-4P 173768-07-5P 173768-08-6P 173768-09-7P  
173768-10-0P 173768-11-1P 173768-12-2P 173768-13-3P 173768-14-4P  
173768-15-5P 173768-16-6P 173768-17-7P 173768-18-8P 173768-19-9P  
173768-20-2P 173768-21-3P 173768-22-4P 173768-23-5P 173768-24-6P  
173768-25-7P 173768-26-8P 173768-27-9P 173768-28-0P 173768-29-1P  
173768-30-4P 173768-31-5P 173768-32-6P 173768-33-7P 173768-34-8P  
173768-35-9P 173768-36-0P 173768-37-1P 173768-38-2P 173768-39-3P  
173768-40-6P 173768-41-7P 173768-42-8P 173768-43-9P 173768-44-0P  
173768-45-1P 173768-46-2P 173768-47-3P 173768-48-4P 173768-49-5P  
173768-50-8P 173768-51-9P 173768-52-0P 173768-53-1P 173768-54-2P  
173768-55-3P 173768-56-4P 173768-57-5P 173768-58-6P 173768-59-7P  
173768-60-0P 173768-61-1P 173768-62-2P 173768-63-3P 173768-64-4P  
173768-65-5P 173768-66-6P 173768-67-7P 173768-68-8P 173768-69-9P  
173768-70-2P 173768-71-3P 173768-72-4P 173768-73-5P 173768-74-6P  
173768-75-7P 173768-76-8P 173768-77-9P 173768-78-0P 173768-79-1P  
173768-80-4P 173768-81-5P 173768-82-6P 173768-83-7P 173768-84-8P  
173768-85-9P 173768-86-0P 173768-87-1P 173768-88-2P 173768-89-3P  
173768-90-6P 173768-91-7P 173768-92-8P 173768-93-9P 173768-94-0P  
173768-95-1P 173768-96-2P 173768-97-3P 173768-98-4P 173768-99-5P  
173769-00-1P 173769-01-2P 173769-02-3P 173769-03-4P 173769-04-5P  
173769-05-6P 173769-06-7P 173769-07-8P 173769-08-9P 173769-09-0P  
173769-10-3P 173769-11-4P 173769-12-5P 173769-13-6P 173769-14-7P  
173769-15-8P 173769-16-9P 173769-17-0P 173769-18-1P 173769-19-2P



173769-20-5P 173769-21-6P 173769-22-7P 173769-23-8P 173769-24-9P  
 173769-25-0P 173769-26-1P 173769-27-2P 173769-28-3P 173769-29-4P  
 RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic  
 preparation); THU (Therapeutic use); BIOL (Biological study); PREP  
 (Preparation); USES (Uses)

(prepn. of retroviral protease inhibiting peptide analogs)

IT	173769-30-7P	173769-31-8P	173769-32-9P	173769-33-0P	173769-34-1P
	173769-35-2P	173769-36-3P	173769-37-4P	173769-38-5P	173769-39-6P
	173769-40-9P	173769-41-0P	173769-42-1P	173769-43-2P	173769-44-3P
	173769-45-4P	173769-46-5P	173769-47-6P	173769-48-7P	173769-49-8P
	173769-50-1P	173769-51-2P	173769-52-3P	173769-53-4P	173769-54-5P
	173769-55-6P	173769-56-7P	173769-57-8P	173769-58-9P	173769-59-0P
	173769-60-3P	173769-61-4P	173769-62-5P	173769-63-6P	173769-64-7P
	173769-65-8P	173769-66-9P	173769-67-0P	173769-68-1P	173769-69-2P
	173769-70-5P	173769-71-6P	173769-72-7P	173769-73-8P	173769-74-9P
	173769-75-0P	173769-76-1P	173769-77-2P	173769-78-3P	173769-79-4P
	173769-80-7P	173769-81-8P	173769-82-9P	173769-83-0P	173769-84-1P
	173769-85-2P	173769-86-3P	173769-87-4P	173769-88-5P	173769-89-6P
	173769-90-9P	173769-91-0P	173769-92-1P	173769-93-2P	173769-94-3P
	173769-95-4P	173769-96-5P	173769-97-6P	173769-98-7P	173769-99-8P
	173770-00-8P	173770-01-9P	173770-02-0P	173770-03-1P	173770-04-2P
	173770-05-3P	173770-06-4P	173770-07-5P	173770-08-6P	173770-09-7P
	173770-10-0P	173770-11-1P	173770-12-2P	173770-13-3P	173770-14-4P
	173770-15-5P	173770-16-6P	173770-17-7P	173770-18-8P	173770-19-9P
	173770-20-2P	173770-21-3P	173770-22-4P	173770-23-5P	173770-24-6P
	173770-25-7P	173770-26-8P	173770-27-9P	173770-28-0P	173770-29-1P
	173770-30-4P	173770-31-5P	173770-32-6P	173770-33-7P	173770-34-8P
	173770-35-9P	173770-36-0P	173770-37-1P	173770-38-2P	173770-39-3P
	173770-40-6P	173770-41-7P	173770-42-8P	173770-43-9P	173770-44-0P
	173770-45-1P	173770-46-2P	173770-47-3P	173770-48-4P	173770-49-5P
	173770-50-8P	173770-51-9P	173770-52-0P	173770-53-1P	173770-54-2P
	173770-55-3P	173770-56-4P	173770-57-5P	173770-58-6P	173770-59-7P
	173770-60-0P	173770-61-1P	173770-62-2P	173770-63-3P	173770-64-4P
	173770-65-5P	173770-66-6P	173770-67-7P	173770-68-8P	173770-69-9P
	173770-70-2P	173770-71-3P	173770-72-4P	173770-73-5P	173770-74-6P
	173770-75-7P	173770-76-8P	173770-77-9P	173770-78-0P	173770-79-1P
	173770-80-4P	173770-81-5P	173770-82-6P	173770-83-7P	173770-84-8P
	173770-85-9P	173770-86-0P	173770-87-1P	173770-88-2P	173770-89-3P
	173770-90-6P	173770-91-7P	173770-92-8P	173770-93-9P	173770-94-0P
	173770-95-1P	173770-96-2P	173770-97-3P	173770-98-4P	173770-99-5P
	173771-00-1P	173771-01-2P	173771-02-3P	173771-03-4P	173771-04-5P
	173771-05-6P	173771-06-7P	173771-07-8P	173771-08-9P	173771-09-0P
	173771-10-3P	173771-11-4P	173771-12-5P	173771-13-6P	173771-14-7P
	173771-15-8P	173771-16-9P	173771-17-0P	173771-18-1P	173771-19-2P
	173771-20-5P	173771-21-6P	173771-22-7P	173771-23-8P	173771-24-9P
	173771-25-0P	173771-26-1P	173771-27-2P	173771-28-3P	173771-29-4P
	173771-30-7P	173771-31-8P	173771-32-9P	173771-33-0P	173771-34-1P
	173771-35-2P	173771-36-3P	173771-37-4P	173771-38-5P	173771-39-6P
	173771-40-9P	173771-41-0P	173771-42-1P	173771-43-2P	173771-44-3P
	173771-45-4P	173771-46-5P	173771-47-6P	173771-48-7P	173771-49-8P
	173771-50-1P	173771-51-2P	173771-52-3P	173771-53-4P	173771-54-5P
	173771-55-6P	173771-56-7P	173771-57-8P	173771-58-9P	173771-59-0P
	173771-60-3P	173771-61-4P	173771-62-5P	173771-63-6P	173771-64-7P

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic  
 preparation); THU (Therapeutic use); BIOL (Biological study); PREP  
 (Preparation); USES (Uses)

(prepn. of retroviral protease inhibiting peptide analogs)

IT	173771-65-8P	173771-66-9P	173771-67-0P	173771-68-1P	173771-69-2P
	173771-70-5P	173771-71-6P	173771-72-7P	173771-73-8P	173771-74-9P
	173771-75-0P	173771-76-1P	173771-77-2P	173771-78-3P	173771-79-4P

173771-80-7P	173771-81-8P	173771-82-9P	173771-83-0P	173771-84-1P
173771-85-2P	173771-86-3P	173771-87-4P	173771-88-5P	173771-89-6P
173771-90-9P	173771-91-0P	173771-92-1P	173771-93-2P	173771-94-3P
173771-95-4P	173771-96-5P	173771-97-6P	173771-98-7P	173771-99-8P
173772-00-4P	173772-01-5P	173772-02-6P	173772-03-7P	173772-04-8P
173772-05-9P	173772-06-0P	173772-07-1P	173772-08-2P	173772-09-3P
173772-10-6P	173772-11-7P	173772-12-8P	173772-13-9P	173772-14-0P
173772-15-1P	173772-16-2P	173772-17-3P	173772-18-4P	173772-19-5P
173772-20-8P	173772-21-9P	173772-22-0P	173772-23-1P	173772-24-2P
173772-25-3P	173772-26-4P	173772-27-5P	173772-28-6P	173772-29-7P
173772-30-0P	173772-31-1P	173772-32-2P	173772-33-3P	173772-34-4P
173772-35-5P	173772-36-6P	173772-37-7P	173772-38-8P	173772-39-9P
173772-40-2P	173772-41-3P	173772-42-4P	173772-43-5P	173772-44-6P
173772-45-7P	173772-46-8P	173772-47-9P	173772-50-4P	

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of retroviral protease inhibiting peptide analogs)

IT 67-64-1, Acetone, reactions 70-23-5, Ethyl bromopyruvate 74-89-5, Methanamine, reactions 75-12-7, Formamide, reactions 75-44-5, Carbonic dichloride 89-98-5, 2-Chlorobenzaldehyde 93-55-0, Propiophenone 98-01-1, 2-Furaldehyde, reactions 98-83-9, reactions 98-86-2, Acetophenone, reactions 100-52-7, Benzaldehyde, reactions 100-55-0, Pyridine-3-methanol 100-83-4, 3-Hydroxybenzaldehyde 104-87-0, p-Tolualdehyde 104-88-1, 4-Chlorobenzaldehyde, reactions 105-39-5, Ethyl chloroacetate 109-94-4, Ethyl formate 110-91-8, Morpholine, reactions 122-78-1, Phenylacetaldehyde 123-08-0, 4-Hydroxybenzaldehyde 123-11-5, p-Anisaldehyde, reactions 123-75-1, Pyrrolidine, reactions 459-57-4, 4-Fluorobenzaldehyde 498-60-2, 3-Furaldehyde 529-20-4, o-Tolualdehyde 534-07-6, 1,3-Dichloroacetone 563-83-7, Isobutyramide 586-98-1, Pyridine-2-methanol 587-04-2, 3-Chlorobenzaldehyde 591-31-1, 688-99-3, 3-Hydroxy-5-hexene 870-46-2, tert-Butyl carbazate 872-85-5, Pyridine-4-carboxaldehyde 925-90-6, Ethylmagnesium bromide 930-45-0, (S,S)-2-Aminocyclopentanol 1121-60-4, Pyridine-2-carboxaldehyde 1779-49-3, Triphenylmethylphosphonium bromide 2043-61-0, Cyclohexanecarboxaldehyde 3731-51-9, 2-Aminomethylpyridine 6089-04-9, 6306-52-1, Valine methyl ester hydrochloride 6372-14-1 6972-05-0, N,N-Dimethylthiourea 10200-59-6, 2-Thiazolecarboxaldehyde 14337-43-0, Ethyl chlorooximidoacetate 16332-06-2, 2-Methoxyacetamide 74111-21-0, (S,S)-2-Aminocyclohexanol 82625-45-4 162740-04-7 162740-05-8 173772-49-1

RL: RCT (Reactant)

(prepn. of retroviral protease inhibiting peptide analogs)

IT 115-08-2P, Methanethioamide 934-53-2P 13242-92-7P 13515-65-6P 14294-10-1P, 4-Morpholinecarbothioamide 15536-75-1P 16689-34-2P 16689-35-3P 24469-50-9P 30293-86-8P 32939-32-5P 32955-22-9P 33142-21-1P 38585-74-9P, 5-Thiazolemethanol 39624-97-0P 40398-36-5P, 1-Pyrrolidinecarbothioamide 41337-78-4P 53370-84-6P 57699-48-6P 57699-55-5P 57699-57-7P 59830-60-3P 65386-28-9P 69353-16-8P 98019-60-4P, 5-Isioxazolemethanol 99805-29-5P 100868-72-2P 126533-95-7P 126533-96-8P 130782-46-6P 134807-06-0P 134807-20-8P 134807-28-6P 134807-29-7P 134807-30-0P 135941-95-6P 135941-97-8P 137515-66-3P 144163-45-1P 144163-71-3P 144163-72-4P 144163-97-3P 144186-53-8P 144186-54-9P 144186-59-4P 149267-56-1P 149267-64-1P 149267-65-2P 149267-73-2P 150767-02-5P 150767-04-7P 150767-05-8P 150767-08-1P 150767-09-2P 150767-10-5P 154212-60-9P 154212-61-0P 154248-99-4P 154972-22-2P 155884-24-5P 161852-61-5P 162537-10-2P 162739-34-6P 162739-35-7P 162739-36-8P 162739-37-9P 162739-44-8P 162739-48-2P 162739-49-3P 162739-50-6P 162739-51-7P 162739-52-8P 162739-54-0P 162739-55-1P 162739-56-2P

162739-57-3P 162739-58-4P 162739-59-5P 162739-60-8P 162739-61-9P  
 162739-62-0P 162739-63-1P 162739-64-2P 162739-65-3P 162739-66-4P  
 162739-67-5P 162739-68-6P 162739-69-7P 162739-70-0P 162739-71-1P  
 162739-72-2P 162739-73-3P 162739-74-4P 162739-75-5P 162739-77-7P  
 162739-78-8P 162739-80-2P 162739-81-3P 162739-82-4P 162739-83-5P  
 162739-84-6P 162739-85-7P 162739-86-8P 162739-88-0P 162739-89-1P  
 162739-90-4P 162739-91-5P 162739-92-6P 162739-93-7P 162739-94-8P  
 162739-95-9P 162739-96-0P 162739-97-1P 162739-98-2P 162739-99-3P  
 162740-00-3P 162740-01-4P 162740-02-5P 162740-03-6P 163658-33-1P  
 173772-48-0P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of retroviral protease inhibiting peptide analogs)


IT 162537-10-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of retroviral protease inhibiting peptide analogs)

RN 162537-10-2 HCAPLUS

CN L-Valine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L27 ANSWER 15 OF 24 HCAPLUS COPYRIGHT 2001 ACS

AN 1995:695866 HCAPLUS

DN 123:84005

TI Preparation of peptide analogs as retroviral protease inhibitors.

IN Kempf, Dale J.; Norbeck, Daniel W.; Sham, Hing Leung; Zhao, Chen; Sowin, Thomas J.; Reno, Daniel S.; Haight, Anthony R.; Cooper, Arthur J.

PA Abbott Laboratories, USA

SO PCT Int. Appl., 194 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A61K031-425

ICS A61K031-42; C07D277-24; C07D275-02; C07D261-08; C07D263-32

CC 34-3 (Amino Acids, Peptides, and

Proteins)


Section cross-reference(s): 1

FAN.CNT 6

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9414436	A1	19940707	WO 1993-US12326	19931216
W: AU, CA, JP, KR				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
CA 2170020	AA	19940630	CA 1993-2170020	19931216
CA 2135890	AA	19940707	CA 1993-2135890	19931216
CA 2135890	C	19960827		
AU 9459546	A1	19940719	AU 1994-59546	19931216
AU 659575	B2	19950518		
EP 674513	A1	19951004	EP 1994-905429	19931216
EP 674513	B1	19960925		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE

JP 08505844	T2	19960625	JP 1993-515323	19931216
JP 2637847	B2	19970806		
EP 727419	A2	19960821	EP 1996-106301	19931216
EP 727419	A3	19961030		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
AT 143262	E	19961015	AT 1994-905429	19931216
ES 2088839	T3	19970201	ES 1994-905429	19931216
JP 09118679	A2	19970506	JP 1996-132368	19931216
JP 10087639	A2	19980407	JP 1996-132369	19931216
EP 1090914	A2	20010411	EP 2000-124382	19931216
EP 1090914	A3	20010418		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, PT, IE				
IL 108126	A1	19950330	IL 1993-108126	19931221
AU 9514927	A1	19950615	AU 1995-14927	19950320
AU 677500	B2	19970424		
US 5539122	A	19960723	US 1995-410996	19950327
US 5552558	A	19960903	US 1995-411032	19950327
US 5696270	A	19971209	US 1995-411140	19950327
US 5580984	A	19961203	US 1995-412253	19950328
US 5679797	A	19971021	US 1995-412244	19950328
US 5583232	A	19961210	US 1995-412821	19950329
US 5597927	A	19970128	US 1995-412438	19950329
US 5674882	A	19971007	US 1995-413136	19950329
US 5583233	A	19961210	US 1995-413290	19950330
US 5625072	A	19970429	US 1995-415827	19950403
US 5591860	A	19970107	US 1995-416272	19950404
US 5597928	A	19970128	US 1995-416607	19950404
US 5608072	A	19970304	US 1995-416259	19950404
US 5565418	A	19961015	US 1995-417304	19950405
US 5659044	A	19970819	US 1995-417165	19950405
US 5659045	A	19970819	US 1995-417295	19950405
US 5616720	A	19970401	US 1995-418056	19950406
US 5635523	A	19970603	US 1995-417879	19950406
US 5892052	A	19990406	US 1995-418031	19950406
US 5541206	A	19960730	US 1995-423387	19950425
US 5846987	A	19981208	US 1997-821609	19970320
US 5886036	A	19990323	US 1997-822071	19970320
AU 9728560	A1	19971211	AU 1997-28560	19970709
AU 697681	B2	19981015		
US 6017928	A	20000125	US 1997-944351	19971006
US 6150530	A	20001121	US 1998-207881	19981208
PRAI US 1992-998114	A	19921229		
US 1993-158587	A	19931202		
US 1983-355945	B2	19830523		
US 1989-355945	B2	19890523		
US 1989-405604	B2	19890908		
US 1989-456124	B2	19891222		
US 1990-518730	A2	19900509		
US 1990-616170	B2	19901120		
US 1991-746020	B2	19910815		
US 1991-777626	B2	19911023		
CA 1993-2135890	A3	19931216		
EP 1994-905429	A3	19931216		
EP 1996-106301	A3	19931216		
JP 1994-515323	A3	19931216		
WO 1993-US12326	W	19931216		
US 1995-413136	A3	19950329		
US 1995-417879	A3	19950406		
US 1995-418031	A3	19950406		
OS MARPAT 123:84005				



AB Title compds. [I; R1 = monosubstituted thiazolyl, oxazolyl, isoxazolyl, isothiazolyl; n = 1-3; R2, R6 = H, alkyl; R3 = alkyl; R4, R41 = (substituted) Ph, thiazolyl, oxazolyl; R7 = (alkyl-substituted) thiazolyl, oxazolyl, isoxazolyl, isothiazolyl; X = H, Y = OH, or X = OH, Y = H; Z = null, O, S, CH<sub>2</sub>, NR<sub>8</sub>; R<sub>8</sub> = alkyl, cycloalkyl, OH, NH<sub>2</sub>, etc.; with provisos], were prepd. Thus, (2S,3S,5S)-5-[N-[N-[[N-methyl-N-[(2-isopropyl-4-thiazolyl)methyl]amino]carbonyl]valinyl]amino]-2-[N-[(5-thiazolyl)methoxycarbonyl]amino]-1,6-diphenyl-3-hydroxyhexane (prepn. via dimerization of Z-phenylalaninal given) inhibited HIV-13B in MT4 cells with IC<sub>50</sub> = 0.025-0.040  $\mu$ M.

ST peptide analog prepn hiv protease inhibitor; virucide heterocyclipeptide analog

IT Virucides and Virustats  
(prepn. of peptide analogs as retroviral protease inhibitors)

IT Peptides, preparation  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(prepn. of peptide analogs as retroviral protease inhibitors)

IT Virus, animal  
(human immunodeficiency, infection; prepn. of peptide analogs as retroviral protease inhibitors)

IT	155213-67-5P	165314-93-2P	165314-94-3P	165314-95-4P	165314-96-5P
	165314-97-6P	165314-98-7P	165314-99-8P	165315-00-4P	165315-01-5P
	165315-02-6P	165315-03-7P	165315-04-8P	165315-05-9P	165315-06-0P
	165315-07-1P	165315-08-2P	165315-09-3P	165315-10-6P	165315-11-7P
	165315-12-8P	165315-13-9P	165315-14-0P	165315-15-1P	165315-16-2P
	165315-17-3P	165315-18-4P	165315-19-5P	165315-20-8P	165315-21-9P
	165315-22-0P	165315-23-1P	165315-24-2P	165315-25-3P	165315-26-4P
	165315-27-5P	165315-28-6P	165315-29-7P	165315-30-0P	165315-31-1P
	165315-32-2P	165315-33-3P	165315-34-4P	165315-35-5P	165315-36-6P
	165315-37-7P	165315-38-8P			

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of peptide analogs as retroviral protease inhibitors)

IT 144114-21-6, Retropepsin

RL: BPR (Biological process); BIOL (Biological study); PROC (Process)  
(prepn. of peptide analogs as retroviral protease inhibitors)

IT 60-35-5, Acetamide, reactions 62-55-5, Thioacetamide 62-56-6, Thiourea, reactions 63-91-2, Phenylalanine, reactions 70-23-5, Ethyl bromopyruvate 74-89-5, Methylamine, reactions 75-04-7, Ethanamine, reactions 75-05-8, Acetonitrile, reactions 75-12-7, Formamide, reactions 75-44-5, Carbonic dichloride 78-81-9, Isobutylamine 79-05-0, Propionamide 88-09-5, 2-Ethylbutyric acid 105-39-5, Ethyl chloroacetate 107-10-8, 1-Aminopropane, reactions 109-89-7, reactions

110-91-8, Morpholine, reactions 122-51-0 123-39-7, N-Methylformamide  
 123-75-1, Pyrrolidine, reactions 503-74-2, 3-Methylbutyric acid  
 534-07-6, 1,3-Dichloroacetone 539-74-2, Ethyl 3-bromopropionate  
 563-83-7, Isobutyramide 638-07-3, Ethyl 4-chloroacetoacetate 867-13-0  
 870-46-2, tert-Butyl carbazate 1118-02-1, Trimethylsilyl isocyanate  
 1759-53-1, Cyclopropanecarboxylic acid 2491-20-5, Alanine methyl ester  
 hydrochloride 3400-45-1, Cyclopentanecarboxylic acid 3721-95-7,  
 Cyclobutanecarboxylic acid 5470-11-1 6065-82-3, Ethyl diethoxyacetate  
 6089-04-9 6160-65-2, Thiocarbonyl diimidazole 6306-52-1, Valine methyl  
 ester hydrochloride 6372-14-1, Z-Phenylalaninol 6921-34-2,  
 Benzylmagnesium chloride 6972-05-0, N,N-Dimethylthiourea 7204-46-8,  
 N,N-Diethylthiourea 13734-41-3 16332-06-2 16982-21-1, Ethyl  
 thiooxamate 19967-55-6 38585-74-9, 5-Thiazolemethanol 65386-28-9  
**65815-64-7** 131052-44-3 133047-44-6 154212-61-0 162739-63-1  
 162739-94-8 162740-04-7 165315-47-9 165315-59-3 165315-76-4  
 165315-78-6 165315-80-0 165315-91-3 165316-16-5 165316-37-0  
 RL: RCT (Reactant)

(prepn. of peptide analogs as retroviral protease inhibitors)


IT 115-08-2P, Methanethioamide 541-46-8P 593-75-9P 631-58-3P,  
 Propanethioamide 1114-38-1P 1503-98-6P, Cyclobutanecarboxamide  
 3217-94-5P, Cyclopentanecarboxamide 6228-73-5P, Cyclopropanecarboxamide  
 13242-92-7P 13515-65-6P, Thioisobutyramide 14294-10-1P,  
 4-Morpholinecarbothioamide 15536-75-1P 16536-95-1P 20295-34-5P,  
 Cyclopropanecarbothioamide 32955-21-8P 32955-22-9P 33142-21-1P  
 39624-97-0P 40398-36-5P, 1-Pyrrolidinecarbothioamide 42202-73-3P,  
 Cyclopentanecarbothioamide 56012-38-5P 59830-60-3P, Z-Phenylalaninal  
 79836-78-5P 98019-60-4P, 5-Isoxazolemethanol 98278-52-5P 99805-29-5P  
 104336-01-8P 110600-55-0P 111138-83-1P 118994-86-8P,  
 5-Oxazolecarboxaldehyde 126533-95-7P 126533-96-8P 126534-22-3P  
 126534-23-4P 127232-41-1P, 5-Oxazolemethanol 133047-45-7P  
 133047-46-8P 133333-27-4P 134878-06-1P 134878-07-2P 135207-10-2P  
 137649-69-5P 143747-06-2P 143838-10-2P 144141-68-4P 144163-44-0P  
 144163-85-9P 144163-96-2P 144163-97-3P 144164-10-3P 144164-11-4P  
 144186-59-4P 148682-88-6P 154212-59-6P 154212-60-9P 154248-99-4P  
 156589-97-8P, Cyclobutanecarbothioamide 156732-12-6P 156732-13-7P  
 156732-15-9P 156769-85-6P **162537-10-2P** 162739-62-0P  
 162739-68-6P 162739-69-7P 162739-70-0P 162739-71-1P 162739-72-2P  
 162739-93-7P 162739-98-2P 162739-99-3P 162740-00-3P 162740-01-4P  
 162740-02-5P 162849-92-5P 162849-93-6P 162849-94-7P 162849-95-8P  
 162849-96-9P 162990-03-6P 165315-39-9P 165315-40-2P 165315-41-3P  
 165315-42-4P 165315-43-5P 165315-44-6P 165315-45-7P 165315-46-8P  
 165315-48-0P 165315-49-1P 165315-50-4P 165315-51-5P 165315-52-6P  
 165315-53-7P 165315-54-8P 165315-55-9P 165315-56-0P 165315-57-1P  
 165315-58-2P 165315-60-6P 165315-61-7P 165315-62-8P 165315-63-9P  
 165315-64-0P 165315-65-1P 165315-66-2P 165315-67-3P 165315-68-4P  
 165315-69-5P 165315-70-8P 165315-71-9P 165315-72-0P 165315-73-1P  
 165315-74-2P 165315-75-3P 165315-77-5P 165315-79-7P 165315-81-1P  
 165315-82-2P 165315-83-3P 165315-84-4P 165315-85-5P 165315-86-6P  
 165315-87-7P 165315-88-8P 165315-89-9P 165315-90-2P 165315-92-4P  
 165315-93-5P 165315-94-6P 165315-95-7P 165315-96-8P 165315-97-9P  
 165316-03-0P 165316-04-1P 165316-05-2P 165316-06-3P 165316-07-4P  
 165316-08-5P 165316-09-6P 165316-10-9P 165316-11-0P 165316-12-1P  
 165316-13-2P 165316-14-3P 165316-15-4P 165316-17-6P 165316-18-7P  
 165316-19-8P 165316-20-1P 165316-21-2P 165316-22-3P 165316-23-4P  
 165316-24-5P 165316-25-6P 165316-26-7P 165316-27-8P 165316-28-9P  
 165316-29-0P 165316-30-3P 165316-31-4P 165316-32-5P 165316-33-6P  
 165316-34-7P 165316-35-8P 165316-36-9P 165316-38-1P 165316-39-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(prepn. of peptide analogs as retroviral protease inhibitors)


IT 65815-64-7  
 RL: RCT (Reactant)  
 (prepn. of peptide analogs as retroviral protease inhibitors)  
 RN 65815-64-7 HCAPLUS  
 CN L-Alanine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 162537-10-2P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of peptide analogs as retroviral protease inhibitors)  
 RN 162537-10-2 HCAPLUS  
 CN L-Valine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L27 ANSWER 16 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1995:522611 HCAPLUS  
 DN 122:291525  
 TI Retroviral protease inhibiting compounds  
 IN Norbeck, Daniel W.; Sham, Hing Leung; Kempf, Dale J.; Zhao, Chen  
 PA Abbott Laboratories, USA  
 SO PCT Int. Appl., 185 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 IC ICM C07D263-34  
 ICS C07D277-587; A61K031-42; A61K031-425  
 CC 34-2 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 1

FAN.CNT 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 9419332	A1	19940901	WO 1994-US1457	19940208
W: CA, JP				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
US 5461067	A	19951024	US 1994-185666	19940201
EP 683772	A1	19951129	EP 1994-908018	19940208
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				

	JP 08507061	T2	19960730	JP 1994-519025	19940208
PRAI	US 1993-23226		19930225		
	US 1994-185666		19940201		
	WO 1994-US1457		19940208		
OS	MARPAT 122:291525				
GI					

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Retroviral protease-inhibiting compds. are disclosed, specifically I [R1, R2 = H, (un)substituted alkyl, aryl, alkenyl, heterocyclyl, substituted carbonyl; Y = NHCHR4CO, NHNR4CO, Q1, Q2; Y' = COCHR3NH, CONR3NH, Q3, Q4; a, b = 0-3; c, d = 1-2; R3, R4 = H, (un)substituted alkyl, aryl, alkenyl, heterocyclyl, substituted carbonyl; R3', R4' = H, alkyl; m, n = 0-1; R5, R6 = C(T)GR7; T = O, S; G = CH2, O, S, NR8; R7 = (un)substituted alkyl or cycloalkyl, aryl, protecting group; R8 = H, alkyl, cycloalkyl] and their salts, esters, and prodrugs. For example, 5-(hydroxymethyl)thiazole (prepd. in 4 steps) reacted with 4-nitrophenyl chloroformate to give 78% of the corresponding carbonate. This reacted with 2-(tert-butoxycarbonylamino)-4S-hydroxy-5S-amino-1,6-diphenyl-2-azahehexane (prepn. given), followed by deprotection and coupling with a corresponding valine deriv., to give title compd. II. At 0.5 nM in vitro, II gave 77% inhibition of HIV-1 protease. II also gave 50% inhibition of cytopathy of MT4 cells by HIV-13B at 0.10-0.11 .mu.M, with a cellular LC50 of 17.mu.M. Examples include 69 syntheses (some prophetic), and similar biol. data for other selected I.

ST valinyl aminohydroxyazahexane prepn retroviral protease inhibitor; HIV protease inhibitor aminohydroxyazahexane prepn

IT Virucides and Virustats  
(prepn. of valinylaminohydroxyazahexane derivs. and analogs as retroviral protease inhibitors)

IT Peptides, preparation  
RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or effector, except adverse); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
(prepn. of valinylaminohydroxyazahexane derivs. and analogs as retroviral protease inhibitors)

IT Virus, animal  
(human immunodeficiency, prepn. of valinylaminohydroxyazahexane derivs. and analogs as retroviral protease inhibitors)

IT 144114-21-6, Retropepsin  
RL: BPR (Biological process); BIOL (Biological study); PROC (Process)  
(HIV; prepn. of valinylaminohydroxyazahexane derivs. and analogs as retroviral protease inhibitors)

IT 60222-90-4P, 5-Hydroxypentanal oxime 128018-44-0P  
RL: BYP (Byproduct); PREP (Preparation)  
(byproduct; prepn. of valinylaminohydroxyazahexane derivs. and analogs as retroviral protease inhibitors)

IT 115-08-2P, Thioformamide 934-53-2P, 2-Chloro-2-phenylpropane 13242-92-7P, 4-(Chloromethyl)-2-(dimethylamino)thiazole 13515-65-6P, 2-Methylpropanethioamide 14294-10-1P, 4-(Aminothiocabonyl)morpholine 15536-75-1P, 2-Methoxythioacetamide 16689-35-3P 24469-50-9P 30293-86-8P, .alpha.-Isocyanatovaline methyl ester 32939-32-5P 32955-22-9P, Ethyl thiazole-5-carboxylate 33142-21-1P, Ethyl 2-chloro-2-formylacetate 38585-74-9P, 5-(Hydroxymethyl)thiazole 39624-97-0P 40398-36-5P, 1-[Amino(thiocarbonyl)]pyrrolidine



41337-78-4P, 2-Carboethoxy-6-ethylpyridine 53370-84-6P 57699-48-6P  
 57699-55-5P 59830-60-3P, N-Benzyloxycarbonyl-L-phenylalaninal  
 65386-28-9P, 4-(Chloromethyl)-2-isopropylthiazole hydrochloride  
 98019-60-4P, 5-(Hydroxymethyl)isoxazole 99805-29-5P 100868-72-2P,  
 2-[(N-Methylamino)methyl]pyridine dihydrochloride 126533-95-7P, Ethyl  
 2-(4-morpholinyl)thiazole-4-carboxylate 126533-96-8P,  
 2-(4-Morpholinyl)-4-(hydroxymethyl)thiazole 134807-06-0P 134807-20-8P  
 134807-28-6P 134807-29-7P 134807-30-0P 137515-66-3P 144163-45-1P  
 144163-71-3P 144163-97-3P 144186-53-8P 144186-59-4P 149267-56-1P  
 149267-64-1P 149267-65-2P 149267-73-2P 150767-00-3P 150767-02-5P  
 150767-04-7P 150767-05-8P 150767-08-1P 154212-60-9P 154212-61-0P  
 154248-99-4P 155884-24-5P, Methyl 2-isopropyl-4-oxazolecarboxylate  
 161852-61-5P **162537-10-2P** 162739-61-9P 162739-62-0P  
 162739-63-1P 162739-64-2P 162739-65-3P 162739-66-4P 162739-67-5P  
 162739-71-1P 162739-76-6P 162739-77-7P 162739-78-8P 162739-79-9P  
 162739-80-2P 162739-81-3P 162739-82-4P 162739-83-5P 162739-84-6P  
 162739-85-7P 162739-86-8P 162739-93-7P 162739-94-8P,  
 2-Isopropyl-4-[(N-methylamino)methyl]oxazole 162739-95-9P 162739-96-0P  
 162739-97-1P, 2-[(N-Methylamino)methyl]-6-ethylpyridine 162739-98-2P,  
 2-(N,N-Dimethylamino)-4-(hydroxymethyl)thiazole 162739-99-3P, Ethyl  
 2-(1-pyrrolidinyl)thiazole-4-carboxylate 162740-00-3P,  
 2-(1-Pyrrolidinyl)-4-(hydroxymethyl)thiazole 162740-01-4P,  
 4-(Chloromethyl)-2-(methoxymethyl)thiazole hydrochloride 162740-02-5P,  
 2-(Methoxymethyl)-4-[(N-methylamino)methyl]thiazole 162740-03-6P,  
 4-Hydroxymethyl-2-isopropylloxazole 163658-33-1P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (intermediate; prepn. of valinylaminohydroxyazahexane derivs. and  
 analogs as retroviral protease inhibitors)  
 IT 144163-72-4P 144186-54-9P 162739-68-6P 162739-69-7P 162739-70-0P  
 162739-72-2P 162739-73-3P 162739-74-4P 162739-75-5P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (intermediate; prepn. of valinylaminohydroxyazahexane derivs. and  
 analogs as retroviral protease inhibitors)  
 IT 150767-06-9P 162739-20-0P 162739-24-4P 162739-28-8P 162739-31-3P  
 RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or  
 effector, except adverse); RCT (Reactant); SPN (Synthetic preparation);  
 THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES  
 (Uses)  
 (prepn. of valinylaminohydroxyazahexane derivs. and analogs as  
 retroviral protease inhibitors)  
 IT 162739-22-2P 162739-25-5P 162739-26-6P 162739-27-7P 162739-29-9P  
 162739-30-2P 162739-32-4P  
 RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or  
 effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic  
 use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (prepn. of valinylaminohydroxyazahexane derivs. and analogs as  
 retroviral protease inhibitors)  
 IT 150767-07-0P 150767-09-2P 162739-33-5P 162739-35-7P 162739-38-0P  
 162739-43-7P 162739-46-0P  
 RL: BAC (Biological activity or effector, except adverse); RCT (Reactant);  
 SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological  
 study); PREP (Preparation); USES (Uses)  
 (prepn. of valinylaminohydroxyazahexane derivs. and analogs as  
 retroviral protease inhibitors)  
 IT 150767-10-5P 162739-21-1P 162739-23-3P 162739-34-6P 162739-36-8P  
 162739-37-9P 162739-39-1P 162739-40-4P 162739-41-5P 162739-42-6P  
 162739-44-8P 162739-45-9P 162739-47-1P 162739-48-2P 162739-49-3P  
 162739-50-6P 162739-51-7P 162739-52-8P 162739-53-9P 162739-54-0P  
 162739-55-1P 162739-56-2P 162739-57-3P 162739-58-4P 162739-59-5P  
 162739-60-8P

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(prepn. of valinylaminohydroxyazahexane derivs. and analogs as retroviral protease inhibitors)

IT 67-64-1, Acetone, reactions 70-23-5, Ethyl bromopyruvate 74-89-5, Methylamine, reactions 75-12-7, Formamide, reactions 75-44-5, Carbonic dichloride 98-01-1, 2-Furaldehyde, reactions 98-83-9, .alpha.-Methylstyrene, reactions 100-52-7, Benzaldehyde, reactions 100-55-0, Pyridine-3-methanol 100-83-4, 3-Hydroxybenzaldehyde 105-39-5, Ethyl chloroacetate 109-94-4, Ethyl formate 110-91-8, Morpholine, reactions 122-51-0, Triethyl orthoformate 123-08-0, 4-Hydroxybenzaldehyde 123-11-5, p-Anisaldehyde, reactions 123-75-1, Pyrrolidine, reactions 459-57-4, 4-Fluorobenzaldehyde 498-60-2, 3-Furaldehyde 534-07-6, 1,3-Dichloroacetone 563-83-7, Isobutyramide 586-98-1, Pyridine-2-methanol 591-31-1, m-Anisaldehyde 688-99-3, 3-Hydroxy-5-hexene 870-46-2, tert-Butyl carbazate 872-85-5, Pyridine-4-carboxaldehyde 925-90-6, Ethylmagnesium bromide 930-45-0, (S,S)-2-Aminocyclopentanol 1121-60-4, Pyridine-2-carboxaldehyde 1779-49-3, Triphenylmethylphosphonium bromide 2043-61-0, Cyclohexanecarboxaldehyde 3731-51-9, 2-(Aminomethyl)pyridine 5470-11-1, Hydroxylamine hydrochloride 6089-04-9, 3,4,5,6-Tetrahydro-2-(2-propynyloxy)-2H-pyran 6160-65-2, Thiocarbonyl diimidazole 6306-52-1, L-Valine methyl ester hydrochloride 6372-14-1, N-Benzoyloxycarbonyl-L-phenylalaninol 6972-05-0, N,N-Dimethylthiourea 7664-41-7, Ammonia, reactions 7693-46-1, 4-Nitrophenyl chloroformate 14337-43-0, Ethyl chlorooximidoacetate 16332-06-2, 2-Methoxyacetamide 57699-57-7 69353-16-8 74111-21-0, (S,S)-2-Aminocyclohexanol 82625-45-4, 4-(2-Morpholinoethoxy)benzaldehyde 130782-46-6 135941-95-6 135941-97-8 162739-87-9 162739-88-0 162739-89-1 162739-90-4 162739-91-5 162739-92-6 162740-04-7, 4-(Chloromethyl)-2-(dimethylamino)thiazole dihydrochloride 162740-05-8, N-Isobutyrylserine methyl ester

RL: RCT (Reactant)

(reactant; prepn. of valinylaminohydroxyazahexane derivs. and analogs as retroviral protease inhibitors)


IT 162537-10-2P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
(intermediate; prepn. of valinylaminohydroxyazahexane derivs. and analogs as retroviral protease inhibitors)

RN 162537-10-2 HCAPLUS

CN L-Valine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L27 ANSWER 17 OF 24 HCAPLUS COPYRIGHT 2001 ACS

AN 1991:186008 HCAPLUS

DN 114:186008

TI Synthesis and spectroscopic properties of azaglutamine amino acid and peptide derivatives


AU Gray, C. J.; Quibell, M.; Jiang, K. L.; Baggett, N.  
 CS Sch. Chem., Univ. Birmingham, Birmingham, B15 2TT, UK  
 SO Synthesis (1991), (2), 141-6  
 CODEN: SYNTBF; ISSN: 0039-7881  
 DT Journal  
 LA English  
 CC 34-3 (Amino Acids, Peptides, and Proteins)  
 OS CASREACT 114:186008  
 AB Azaglutamines  $\text{H}_2\text{NCOCH}_2\text{CH}_2\text{N}(\text{NHBoc})\text{CO}_2\text{R}$  (Boc =  $\text{Me}_3\text{CO}_2\text{C}$ ; R = Et,  $\text{CH}_2\text{Ph}$ ) were  
 prepd. by treating  $\text{H}_2\text{NCOCH}_2\text{NHNHBOc}$  (I) with  $\text{ClCO}_2\text{R}$ . Dipeptides  
 $\text{H}_2\text{NCOCH}_2\text{CH}_2\text{N}(\text{NHBoc})\text{CO-X-OCH}_2\text{Ph}$  (X = Gly, Phe) were prepd. by treating I  
 with 2,4-( $\text{O}_2\text{N}$ ) $\text{C}_6\text{H}_3\text{O}_2\text{C-X-OCH}_2\text{Ph}$ . I was treated with  $\text{OCNCH}_2\text{CO}_2\text{Et}$  to give  
 $\text{H}_2\text{NCOCH}_2\text{CH}_2\text{N}(\text{NHBoc})\text{CONHCH}_2\text{CO}_2\text{Et}$   
 ST azaglutamine peptide; glutamine aza  
 IT Peptides, preparation  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (azaglutamine-contg., prepn. and NMR of)  
 IT 870-46-2, tert-Butyl carbazate 30189-48-1 41863-52-9  
 RL: RCT (Reactant)  
 (addn. reaction of, with acrylamide)  
 IT 79-06-1, 2-Propenamide, reactions  
 RL: RCT (Reactant)  
 (addn. reaction of, with tert-Bu carbazate)  
 IT 133382-90-8P 133382-91-9P 133382-92-0P 133382-95-3P 133382-96-4P  
 133382-99-7P 133383-00-3P 133383-04-7P 133383-05-8P 133383-06-9P  
 133383-07-0DP, derivs.  
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and NMR of)  
 IT 133382-98-6P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and reaction of, with Et isocyanatoacetate)  
 IT 1738-76-7P, Glycine benzyl ester tosylate 1738-78-9P, L-Phenylalanine  
 benzyl ester tosylate  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and reaction of, with bis(dinitrophenyl) carbonate)  
 IT 133382-89-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and reaction of, with chloroformates or Et isocyanoacetate)  
 IT 133382-97-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and reaction of, with dinitrophenoxycarbonyl amino acid esters)  
 IT **133382-93-1P 133382-94-2P**  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and reaction of, with hydrazide)  
 IT 133383-19-4P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 IT 133383-01-4P 133383-02-5P 133383-03-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn., sapon. and NMR of)  
 IT 501-53-1, Benzyl chloroformate 541-41-3, Ethyl chloroformate  
 7497-12-3, Bis(2,4-dinitrophenyl) carbonate  
 RL: RCT (Reactant)  
 (reaction of, with (tert-butoxycarbonylhydrazino)propanamide)  
 IT 2949-22-6, Ethyl isocyanato acetate  
 RL: RCT (Reactant)  
 (reaction of, with hydrazides)  
 IT **133382-93-1P 133382-94-2P**  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)

(prepn. and reaction of, with hydrazide)

RN 133382-93-1 HCAPLUS


CN L-Phenylalanine, N-[(2,4-dinitrophenoxy)carbonyl]-, phenylmethyl ester  
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 133382-94-2 HCAPLUS

CN Glycine, N-[(2,4-dinitrophenoxy)carbonyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)



L27 ANSWER 18 OF 24 HCAPLUS COPYRIGHT 2001 ACS

AN 1991:62698 HCAPLUS

DN 114:62698

TI New macrocyclic pseudopeptides containing urethane backbone linkages

AU Wu, Youling; Kohn, Joachim

CS Dep. Chem., Rutgers, State Univ., New Brunswick, NJ, USA

SO J. Am. Chem. Soc. (1991), 113(2), 687-8

CODEN: JACSAT; ISSN: 0002-7863

DT Journal

LA English

CC 34-3 (Amino Acids, Peptides, and Proteins)

OS CASREACT 114:62698

GI

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*


AB The use of urethane bonds as a new type of amide bone mimetic in the design of pseudopeptides was investigated. Urethane backbone linkages were derived from the side chain hydroxyl group of tyrosine by reacting Boc-Tyr-OHex (Boc = Me<sub>3</sub>CO<sub>2</sub>C, Hex = hexyl) with p-nitrophenyl chloroformate, followed by reaction with alanine benzyl ester to give urethane-bonded pseudodipeptide I. The resulting Tyr-Ala pseudodipeptide could be cyclized to 24- and 36-membered macrocyclic pseudopeptides II and III, resp., with cyclization yields of about 80%. The unusual ease with which the Tyr-Ala pseudodipeptide underwent cyclizations is probably a consequence of the cis conformation of the urethane bond. In a phase

transfer expt., the ion binding properties of the 24-membered macrocycle II were studied. II was an effective and selective phase transfer agent for Li<sup>+</sup> ions, solubilizing up to 0.6 molar equivalents of Li<sup>+</sup> ions in chloroform. The uptake of Na<sup>+</sup> and K<sup>+</sup> ions was only 0.025 and 0.007 molar equivalents, resp. Since the structure of II can be readily modified, the synthetic approach gives rise to a family of new macrocyclic pseudopeptides.

ST macrocyclic pseudopeptide urethane backbone linkage  
 IT Alkali metals, reactions  
 RL: RCT (Reactant)  
 (binding of, with macrocyclic pseudopeptide contg. urethane backbone linkages)  
 IT Peptides, preparation  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (cyclopseudo-, prepn. of, with urethane backbone linkages)  
 IT 573-83-1 3324-58-1 18390-55-1  
 RL: PROC (Process)  
 (binding of, with macrocyclic pseudopeptide contg. urethane backbone linkages)  
 IT 17831-01-5  
 RL: RCT (Reactant)  
 (coupling of, with tyrosine active carbonate deriv.)  
 IT 131152-83-5P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and binding of, with alkali metals)  
 IT 131152-91-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and coupling of, with alanine benzyl ester)  
 IT **131152-89-1P**  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and cyclization of)  
 IT 131152-86-8P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and esterification of, with nitrophenol)  
 IT **131152-87-9P**  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and partial deblocking of)  
 IT 131152-90-4P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and reaction of, with nitrophenyl chloroformate)  
 IT 131152-84-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 IT 131152-85-7P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn., catalytic transfer hydrogenolysis, and metal-binding properties of)  
 IT 7693-46-1, p-Nitrophenyl chloroformate  
 RL: RCT (Reactant)  
 (reaction of, with tyrosine deriv.)  
 IT 94326-61-1  
 RL: RCT (Reactant)  
 (tert-butoxycarbonylation of)  
 IT **131152-89-1P**  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and cyclization of)  
 RN 131152-89-1 HCAPLUS  
 CN L-Tyrosine, hexyl ester, ester with N-carboxy-L-alanine 1-(4-nitrophenyl) ester, mono(trifluoroacetate) (9CI) (CA INDEX NAME)


CM 1

CRN 131152-88-0  
CMF C25 H31 N3 O8  
CDES \*



CM 2

CRN 76-05-1  
CMF C2 H F3 O2




IT 131152-87-9P

RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. and partial deblocking of)

RN 131152-87-9 HCAPLUS

CN L-Tyrosine, N-[(1,1-dimethylethoxy)carbonyl]-, hexyl ester, ester with  
N-carboxy-L-alanine 1-(4-nitrophenyl) ester (9CI) (CA INDEX NAME)



L27 ANSWER 19 OF 24 HCAPLUS COPYRIGHT 2001 ACS

AN 1990:99261 HCAPLUS

DN 112:99261

TI Preparation of N-(phosphonocyclohexylhydroxypropyl) derivatives of amino  
acids and dipeptides as renin inhibitors

IN Patel, Dinesh V.

PA Squibb, E. R., and Sons, Inc., USA

SO Eur. Pat. Appl., 121 pp

CODEN: EPXXDW

DT Patent

LA English

IC ICM C07K005-06

ICS A61K037-64; C07F009-40; C07F009-65; C07F009-32; C07F009-44;  
A61K031-66

CC 34-3 (Amino Acids, Peptides, and Proteins)

Section cross-reference(s): 1

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 331105	A2	19890906	EP 1989-103489	19890228
	EP 331105	A3	19900905		
	R: AT, BE, CH, DE, ES, FR, GB, IT, LI, LU, NL, SE				
	WO 8907940	A1	19890908	WO 1989-US777	19890223
	W: DK, HU, JP				
	HU 52785	A2	19900828	HU 1989-2302	19890223
	JP 02503440	T2	19901018	JP 1989-503323	19890223
	ZA 8901594	A	19891129	ZA 1989-1594	19890301
	AU 8930999	A1	19890907	AU 1989-30999	19890302
	DK 8905469	A	19891227	DK 1989-5469	19891102
	US 5217958	A	19930608	US 1990-509398	19900412
PRAI	US 1988-163593		19880303		
	WO 1989-US777		19890223		
	US 1989-317257		19890228		

OS MARPAT 112:99261

AB Amino acid and dipeptide derivs. X-Y-(CHR5CONH)rCHR4CONHCHR3CH(OH)P(:M)(Z1 R2)ZR1 [I; M = O, S; Y = CH2, NH, O; when Y = CH2, X = N-(substituted alkyl)carbamoyl or -sulfamoyl, substituted alkanoyl, alkoxy carbonyl, alkylthio, alkanoylthio, etc.; when Y = O, X = N-(substituted alkyl)carbamoyl, substituted alkanoyl, alkoxy carbonyl, phosphono, etc.; when Y = NH, X = N-(substituted alkyl)carbamoyl, substituted alkoxy carbonyl, alkanoyl, alkyl, alkylthio, etc.; R1, R2 = H, (cyclo)alkyl, arylalkyl, (hetero)aryl; Z, Z1 = bond, (un)substituted CH2, NH, cyclic amino, heterocyclyl; R3, R5 = H, lower (halo)alkyl, arylalkyl, heterocyclalkyl, cycloalkyl, (CH2)nOH, (CH2)nNH2, (CH2)nSH, (CH2)nNHC(:NH)NH2, (CH2)nCONH2, N-substituted 5-imidazolylalkyl, etc.; R4 = any group defined for R3 and R5, N-substituted 2-imidazolyl, 4- or 2-thiazolylalkyl, 3-pyrazolylalkyl, 4- or 2-oxazolylalkyl; n = 1-5], useful as cardiovascular agents in the treatment of hypertension, congestive heart failure, renin-dependent hyperaldosteronism, myocardial infarction, etc., and as a diagnostic agent in detg. renin related disorders (no data), were prepd. I are also inhibitors of retroviral protease and thus are useful as virucides against human T-cell leukemia virus HTLV-1 and HTLV-III (no data). Thus, hydrogenation of (S)-BOC-Phe-OH (BOC = Me3CO2C) over PtO2 and amidation of the resulting (S)-BOC-NHCHQCO2H (Q = cyclohexylmethyl) with MeNHOMe.HCl in THF contg. carbodiimidazole gave (S)-BOC-NHCHQCONMeOMe. Redn. of the latter with LiAlH4 in THF/Et2O at 0.degree. to (S)-BOC-NHCHQCHO followed by addn. reaction with (MeO)2PH in DMF in the presence of KF gave a 12.7:1.0 diastereomeric mixt. of (1S)-BOC-NHCHQCH(OH)P(OMe)2 (Q unchanged). N-Deprotection of the latter with 1.2 N HCl in EtOAc and condensation with BOC-Leu-OH.H2O in DMF in the presence of hydroxybenzotriazole, Et3N, and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.HCl gave a major diastereomer (1S)-BOC-Leu-NHCHQCH(OH)P(OMe)2. Approx. 25 I were prepd.

ST amino acid phosphonocyclohexylhydroxypropyl renin inhibitor; peptide

phosphonocyclohexylhydroxypropyl prepn renin inhibitor; cardiovascular agent phosphonocyclohexylhydroxypropyl dipeptide amide; antihypertensive phosphonocyclohexylhydroxypropyl dipeptide amide

IT Aldosteronism

(treatment of, N-(phosphonohydroxyalkyl)amino acids and -dipeptides for)

IT Antihypertensives

Cardiovascular agents  
 Virucides and Virustats  
 (N-(phosphonohydroxyalkyl)amino acid and -dipeptide amides)

IT Amides  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (amino, N-(phosphonohydroxyalkyl), prepn. of, as renin inhibitors and cardiovascular agents)

IT Peptides, compounds  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (di-, amides, N-(phosphonohydroxyalkyl), prepn. of, as renin inhibitors and cardiovascular agents)

IT Heart  
 (failure, treatment of, N-(phosphonohydroxyalkyl)amino acids and -dipeptides for)

IT Heart, disease or disorder  
 (infarction, treatment of, N-(phosphonohydroxyalkyl)amino acids and -dipeptides for)

IT 3587-60-8  
 RL: RCT (Reactant)  
 (N-alkylation by, of histidine Me ester deriv.)

IT 7693-46-1, p-Nitrophenyl chloroformate  
 RL: RCT (Reactant)  
 (acylation by, of Me phenylalaninate)

IT 501-53-1, Benzyl chloroformate  
 RL: RCT (Reactant)  
 (acylation by, of aminocaproic acid)

IT 108-23-6, Isopropyl chloroformate 1070-83-3, tert-Butylacetic acid  
 RL: RCT (Reactant)  
 (acylation by, of histidinamide deriv.)

IT 24424-99-5, Di-tert-Butyl dicarbonate  
 RL: RCT (Reactant)  
 (acylation by, of histidine Me ester)

IT 645-45-4, Hydrocinnamoyl chloride  
 RL: RCT (Reactant)  
 (acylation by, of leucine)

IT 3400-45-1, Cyclopentanecarboxylic acid  
 RL: RCT (Reactant)  
 (acylation by, of phenylalanylleucinamide)

IT 60-32-2  
 RL: RCT (Reactant)  
 (acylation of, by benzyl chloroformate)

IT 7524-50-7, L-Phenylalanine methyl ester hydrochloride  
 RL: RCT (Reactant)  
 (acylation of, by cyclopentanecarboxylic acid)

IT 61-90-5, L-Leucine, reactions  
 RL: RCT (Reactant)  
 (acylation of, by hydrocinnamoyl chloride)

IT 762-04-9 868-85-9, Dimethyl phosphite  
 RL: RCT (Reactant)  
 (addn. reaction of, with aminocyclohexylpropanal deriv.)

IT 75-26-3, 2-Bromopropane 78-77-3, 1-Bromo-2-methylpropane  
 RL: RCT (Reactant)  
 (alkylation by, of [(hydroxymethoxyphosphinyl)ethyl]oxazolidinecarboxylate)

IT 25024-53-7  
 RL: RCT (Reactant)  
 (amidation of, with (aminophenyl)phosphonate deriv.)

IT 2018-66-8  
 RL: RCT (Reactant)  
 (amidation of, with (aminopropyl)phosphonate deriv.)



IT 110-91-8, Morpholine, reactions  
 RL: RCT (Reactant)  
 (amidation of, with Me [(nitrophenoxy)carbonyl]phenylalaninate)

IT 6638-79-5, O,N-Dimethylhydroxylamine hydrochloride  
 RL: RCT (Reactant)  
 (amidation of, with aminocyclohexylpropanoic acid deriv.)

IT 501-52-0, Hydrocinnamic acid  
 RL: RCT (Reactant)  
 (amidation of, with leucinamide deriv.)

IT 74-89-5, Methylamine, reactions  
 RL: RCT (Reactant)  
 (amination by, of (hydroxymethoxyphosphinyl)oxazolidinecarboxylate)

IT 13139-15-6  
 RL: RCT (Reactant)  
 (condensation of, with di-Me (aminocyclohexylhydroxypropyl)phosphonate)

IT 71-00-1, L-Histidine, reactions  
 RL: RCT (Reactant)  
 (conversion of, to Me ester)

IT 77-76-9, 2,2-Dimethoxypropane  
 RL: RCT (Reactant)  
 (cyclocondensation of, with [(cyclohexylmethyl)hydroxyethyl]carbamate deriv.)

IT 1498-40-4, Ethyl dichlorophosphine  
 RL: RCT (Reactant)  
 (esterification of, with methanol)

IT 13734-34-4  
 RL: RCT (Reactant)  
 (hydrogenation of)

IT 9015-94-5, Renin, uses and miscellaneous  
 RL: USES (Uses)  
 (inhibitors, N-(phosphonocyclohexylhydroxypropyl)amino acids and dipeptides)

IT 75-16-1, Methylmagnesium bromide  
 RL: RCT (Reactant)  
 (methylation by, of [(hydroxymethoxyphosphinyl)ethyl]oxazolidinecarboxylate)


IT 2666-93-5  
 RL: RCT (Reactant)  
 (peptide coupling of, in prepn. of renin inhibitor)

IT 1947-00-8P 2752-56-9P 15027-08-4P 20898-43-5P 22888-60-4P,  
 L-Histidine methyl ester hydrochloride 27852-48-8P 33014-68-5P  
 37736-82-6P 38155-45-2P **54601-21-7P** 64152-76-7P  
 64155-03-9P 75691-91-7P 83468-82-0P 83468-83-1P 98105-42-1P  
 111629-40-4P 114457-62-4P 114473-20-0P 115766-13-7P 120915-52-8P  
 120915-53-9P 121533-66-2P 125399-33-9P 125399-34-0P 125399-35-1P  
 125399-36-2P 125399-37-3P 125399-38-4P 125399-39-5P 125399-40-8P  
 125399-41-9P 125399-42-0P 125399-43-1P 125399-44-2P 125399-45-3P  
 125399-46-4P 125399-47-5P 125399-48-6P 125399-49-7P 125399-50-0P  
 125399-51-1P 125399-52-2P 125399-53-3P 125399-54-4P 125399-55-5P  
 125399-56-6P 125399-57-7P 125399-58-8P 125399-59-9P 125399-60-2P  
 125435-73-6P 125435-74-7P 125435-75-8P 126431-13-8P 126431-14-9P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of, as intermediate for renin inhibitor and cardiovascular agent)

IT 125399-09-9P 125399-10-2P 125399-12-4P 125399-13-5P 125399-14-6P  
 125399-15-7P 125399-16-8P 125399-17-9P 125399-19-1P 125399-20-4P  
 125399-21-5P 125399-22-6P 125399-23-7P 125399-24-8P 125399-25-9P  
 125399-26-0P 125399-27-1P 125399-28-2P 125399-29-3P 125399-30-6P  
 125399-31-7P 125399-32-8P 125435-72-5P 125472-49-3P  
 RL: SPN (Synthetic preparation); PREP (Preparation)

(prepn. of, as renin inhibitor and cardiovascular agent)  
 IT 75-77-4, Trimethylsilyl chloride, reactions  
 RL: RCT (Reactant)  
 (silylation by, of [(cyclohexylmethyl)(ethylmethoxyphosphinyl)hydroxy  
 ethyl]carbamate)  
 IT 54601-21-7P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of, as intermediate for renin inhibitor and cardiovascular  
 agent)  
 RN 54601-21-7 HCAPLUS  
 CN L-Phenylalanine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA  
 INDEX NAME)




Absolute stereochemistry.



L27 ANSWER 20 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1985:7099 HCAPLUS  
 DN 102:7099  
 TI 8-Quinoliny carbamates and their use as urinary tract antimicrobials  
 IN Paxton, Larry D.; Madison, Rita A.; Dunbar, Joseph E.  
 PA Dow Chemical Co., USA  
 SO U.S., 6 pp.  
 CODEN: USXXAM  
 DT Patent  
 LA English  
 IC A61K031-47; C07D215-34  
 NCL 424258000  
 CC 34-3 (Amino Acids, Peptides, and  
 Proteins)  
 Section cross-reference(s): 10, 27

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 4472404	A	19840918	US 1982-408292	19820816
GI					



AB Title compds. I (R1 = H, alkyl; R2 = H, NO2, halo; R3 = H, halo; R4 = alkyl) were prepd. as title agents. Thus, quinoline II was treated with Et isocyanoacetate in refluxing EtCOMe contg. Bu2Sn dilaurate to give glycinate III. III was active against Bacillus subtilus with a min. inhibitory concn. 10 ppm.

ST quinolinyl carbamate prepn urinary antimicrobial; glycine quinolinylloxycarbonyl prepn antimicrobial; bactericide quinolinyl carbamate; fungicide quinolinyl carbamate

IT Bactericides, Disinfectants, and Antiseptics  
Fungicides and Fungistats  
(quinolinyl carbamates)

IT Urinary tract  
(quinolinyl carbamates as antimicrobials for)

IT 19498-91-0P 51203-25-9P 93775-51-0P 93775-52-1P 93775-53-2P  
93775-54-3P **93775-55-4P** 93775-56-5P 93775-57-6P  
93775-58-7P 93775-59-8DP, derivs.  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(prepn. and antimicrobial activity of)

IT 93775-51-0P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. and hydrolysis and antimicrobial activity of)

IT 19642-75-2P  
RL: SPN (Synthetic preparation); PREP (Preparation)  
(prepn. of)

IT 130-16-5 521-74-4 773-76-2 826-81-3 4008-48-4  
RL: RCT (Reactant)  
(reaction of, with Et isocyanatoacetate)


IT 1943-83-5  
RL: RCT (Reactant)  
(reaction of, with chlorohydroxyquinoline)

IT 2949-22-6  
RL: RCT (Reactant)  
(reaction of, with hydroxyquinolines)


IT **93775-55-4P**  
RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)  
(prepn. and antimicrobial activity of)

RN 93775-55-4 HCAPLUS

CN Glycine, N-[[5-nitro-8-quinolinyl]oxy]carbonyl]-, ethyl ester (9CI) (CA INDEX NAME)



L27 ANSWER 21 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1977:568362 HCAPLUS  
 DN 87:168362  
 TI Preparation and properties of some .alpha.-aza-amino-acid derivatives,  
 their possible use in peptide synthesis  
 AU Gray, C. J.; Ireson, J. C.; Parker, R. C.  
 CS Dep. Chem., Univ. Birmingham, Birmingham, Engl.  
 SO Tetrahedron (1977), 33(7), 739-43  
 CODEN: TETRAB  
 DT Journal  
 LA English  
 CC 34-2 (Synthesis of **Amino Acids, Peptides,**  
**and Proteins**)  
 Section cross-reference(s): 28  
 GI



AB Azaglycine and azaphenylalanine derivs. were prepd. by acylation of  
 acylhydrazides with alkyl and aryl chloroformates. E.g., Me3CO2CNHNH2 and  
 AcNHNHCH2Ph with ClCO2Ph gave Me3CO2CNHNHCO2Ph (I) and AcNHN(CH2Ph)CO2Ph  
 (II), resp. Esters of acetyl- and benzoylazaamino acids underwent rapid  
 cyclization to oxadiazolones and were unsuitable for peptide synthesis.  
 E.g., II gave the oxadiazolone III on treatment with NaOH or NH2OH or  
 incubation at pH 7 at 37.degree. for several days. Me3CO2CNHNHCON3,  
 prepd. from I by sequential reactions with NH2NH2 and amyl nitrite was too  
 unreactive for peptide synthesis. Coupling reaction of  
 2,4-dinitrophenyloxycarbonylphenylalanine Et ester with BzNHNH2 and  
 AcNHNH2CH2Ph gave benzoylazaglycyl-L-phenylalanine Et ester and  
 acetylazaphenylalanyl-L-phenylalanine Et ester, resp.  
 ST aza amino acid prepn cyclization; hydrazide acyl acylation; aza peptide;  
 ring closure aza amino acid; oxadiazolone  
 IT Amino acids, preparation  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (aza-, prepn. of, by acylation of hydrazides)  
 IT Ring closure and formation  
 (of aza amino acids, oxadiazolones by)  
 IT Hydrazides  
 RL: RCT (Reactant)

(acyl, acylation of, aza amino acids by)

IT Peptides, preparation  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (aza-, prepn. of, by reaction of peptides with acylhydrazides)

IT 613-94-5 870-46-2 1215-52-7 7151-53-3 53370-84-6  
 RL: RCT (Reactant)  
 (acylation of)

IT 3081-24-1  
 RL: RCT (Reactant)  
 (dinitrophenyloxycarbonylation of)

IT 64512-90-9P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and azidation of)

IT 64512-92-1P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and condensation reactions of, with benzoylhydrazide and  
 acetylbenzylhydrazine)

IT 53370-82-4P 53370-85-7P 64512-88-5P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and cyclization of)

IT 64512-91-0P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and reaction of, with benzylamine)


IT 1199-02-6P 15081-44-4P 27643-12-5P 52414-76-3P 53370-83-5P  
 57699-63-5P 57699-88-4P 64512-81-8P 64512-82-9P 64512-83-0P  
 64512-84-1P 64512-85-2P 64512-86-3P 64512-87-4P 64512-93-2P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)

IT 64512-89-6P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn., deprotection, and reactions of, with ammonia and hydrazine)

IT 64512-92-1P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and condensation reactions of, with benzoylhydrazide and  
 acetylbenzylhydrazine)

RN 64512-92-1 HCAPLUS  
 CN L-Phenylalanine, N-[(2,4-dinitrophenoxy)carbonyl]-, ethyl ester (9CI) (CA  
 INDEX NAME)


Absolute stereochemistry.




L27 ANSWER 22 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
 AN 1975:4556 HCAPLUS  
 DN 82:4556  
 TI Hydrazine compounds as hetero-components in peptides. XIX. Eleodoisin  
 peptides containing the N2-methylcarbazoyl radical (azaalanine)  
 AU Niedrich, Hartmut; Oehme, Christa; Bergmann, Jutta  
 CS Zentralinst. Molekularbiol., DAW, Berlin, E. Ger.  
 SO J. Prakt. Chem. (1974), 316(5), 741-9  
 CODEN: JPCEAO

DT Journal  
 LA German  
 CC 34-3 (Synthesis of **Amino Acids, Peptides,**  
**and Pr teins**)  
 AB Acylation, sapon., and peptide coupling of 2-azaalanine peptides were  
 studied with H<sub>2</sub>NNMeCONHCHPhCO<sub>2</sub>Me (Azala-Phe-OMe) (I). 6-Azala-eledoisin  
 sequence 6-11 (II) and 5-Ala-6-Azala-eledoisin 5-11 (III) were prepd. by  
 coupling of I with Ile-Gly-Leu-Met-NH<sub>2</sub> in the presence of  
 hydroxysuccinimide and dicyclohexylcarbodiimide. II and III had only the  
 biol. activity of the 7-11 sequence, i.e. .apprx.1% of the 6-11 sequence.  
 ST azaalanine peptide eledoisin  
 IT Peptides, preparation  
 RL: PREP (Preparation)  
 (azaalanine-contg., eledoisin-related)  
 IT 3069-69-0  
 RL: RCT (Reactant)  
 (methylation of)  
 IT 40203-94-9  
 RL: RCT (Reactant)  
 (peptide coupling reaction with azaalanine derivs.)  
 IT 2577-90-4  
 RL: RCT (Reactant)  
 (peptide coupling reactions of)  
 IT 1142-20-7 2280-71-9 2304-96-3 2592-19-0 15761-38-3  
 RL: RCT (Reactant)  
 (peptide coupling reactions with azaalanine deriv.)  
 IT 54601-18-2P 54601-22-8P 54601-24-0P 54601-25-1P 54601-26-2P  
 54601-27-3P 54601-28-4P  
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. and peptide coupling reactions of)  
 IT 54601-19-3P 54601-20-6P 54601-23-9P 54601-29-5P 54601-30-8P  
 54601-31-9P 54601-32-0P 54601-33-1P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 IT 60-34-4  
 RL: RCT (Reactant)  
 (reaction of, with amino acids)  
 IT 2483-49-0 **54601-21-7**  
 RL: RCT (Reactant)  
 (reaction of, with methylhydrazine)  
 IT **54601-21-7**  
 RL: RCT (Reactant)  
 (reaction of, with methylhydrazine)  
 RN 54601-21-7 HCAPLUS  
 CN L-Phenylalanine, N-[(4-nitrophenoxy)carbonyl]-, methyl ester (9CI) (CA  
 INDEX NAME)


Absolute stereochemistry:




AN 1970:425854 HCAPLUS  
 DN 73:25854  
 TI Cyclizations via phenylthio- and phenoxy-carbonylamino intermediates. II. Formation of azasuccinic anhydrides and their polycondensation products. Pyridiniocarbonylamino intermediates  
 AU Baudet, Pierre; Otten, Cl.; Rao, D.  
 CS Lab. Chim. Org., Univ. Geneve, Geneva, Switz.  
 SO Helv. Chim. Acta (1970), 53(4), 859-69  
 CODEN: HCACAV  
 DT Journal  
 LA French  
 CC 34 (Synthesis of Amino Acids, Peptides, and Proteins)  
 AB The N-(p-nitrophenoxy-carbonyl) derivs. of glycine, DL-alanine and DL-leucine were transformed by pyridine into azasuccinic, 3-methyl-2-azasuccinic and 3-isobutyl-2-azasuccinic anhydride, resp., by way of N-carbamoylpyridinium cation intermediates. The above azasuccinic anhydrides polycondensed in the presence of pyridine yielding the corresponding polyglycine, poly-DL-alanine and poly-DL-leucine, resp. Both cyclization and polycondensation were also catalyzed by gamma.-collidine, but at a low rate. N-(p-Nitrophenoxy-carbonyl)glycine reacted with lysozyme in the presence of pyridine. Several glycine residues were introduced into the enzyme which was rendered insol. and partially inactivated.  
 ST azasuccinic anhydrides; anhydrides azasuccinic; polyamino acids; lysozyme glycy; glycylllysozyme  
 IT Pyridinium compounds  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (carbonyl derivs. of amino acids, intermediates in oxazolidinedione prepn.)  
 IT 2,5-Oxazolidinedione, derivs.  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of, from phenoxy-carbonyl amino acids and pyridine)  
 IT 1192-73-0P 2185-00-4P 4289-99-0P 21639-02-1P 25281-63-4P  
 25718-94-9P 25734-27-4P 25988-64-1P 26283-00-1P 26334-33-8P  
 26952-07-8P 27317-15-3P 27317-16-4P 27317-17-5P  
 27317-18-6P 27317-19-7P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 IT 21639-02-1P 27317-15-3P 27317-16-4P  
 27317-18-6P 27317-19-7P  
 RL: SPN (Synthetic preparation); PREP (Preparation)  
 (prepn. of)  
 RN 21639-02-1 HCAPLUS  
 CN Glycine, N-carboxy-, N-(p-nitrophenyl) ester (8CI) (CA INDEX NAME)




RN 27317-15-3 HCAPLUS  
 CN Alanine, N-carboxy-, ethyl N-(p-nitrophenyl) ester, DL- (8CI) (CA INDEX NAME)




RN 27317-16-4 HCAPLUS  
CN Alanine, N-carboxy-, N-(p-nitrophenyl) ester, DL- (8CI) (CA INDEX NAME)



RN 27317-18-6 HCAPLUS  
CN Leucine, N-carboxy-, ethyl N-(p-nitrophenyl) ester, DL- (8CI) (CA INDEX NAME)



RN 27317-19-7 HCAPLUS  
CN Leucine, N-carboxy-, N-(p-nitrophenyl) ester, DL- (8CI) (CA INDEX NAME)



L27 ANSWER 24 OF 24 HCAPLUS COPYRIGHT 2001 ACS  
AN 1967:18854 HCAPLUS  
DN 66:18854  
TI Peptide syntheses. XXXII. .gamma.-Hydroxyisocaproyl and 3-nitrophenoxycarbonyl moieties as protecting groups  
AU Wieland, Theodor; Lamperstorfer, Ch.; Birr, Christian  
CS Univ. Frankfurt/M., Frankfurt/M., Ger.  
SO Makromol. Chem. (1966), 92, 277-86  
CODEN: MACEAK  
DT Journal  
LA German  
CC 34 (Synthesis of Amino Acids, Peptides, and Proteins)



AB cf. CA 64, 14272g. The title protecting groups were found to react readily with various amino acids, and to be removed quant. by CF<sub>3</sub>CO<sub>2</sub>H treatment and irradiation with uv light ( $\lambda$  >290 m.m.), resp. Thus, a soln. of 15 g. glycine (I) in 100 ml. 2N NaOH was evapd. to dryness in vacuo and the residue dried in a desiccator. The resulting glycine Na salt (9.7 g.) was pulverized and dissolved in 100 g. molten imidazole (II) on a boiling water bath. Isocapro lactone (16 g.) prepd. according to Stevens and Tarbell (CA 50, 937e) was added, the mixt. was heated 1 hr., cooled, and washed several times with Me<sub>2</sub>CO to give 43% N-( $\gamma$ -hydroxyisocaproyl)glycine (III) Na salt, mixed with a little I Na salt. The crude salt (4.2 g.) was dissolved in 10 ml. H<sub>2</sub>O, cooled in ice, acidified to pH 2.6 with 0.1N H<sub>2</sub>SO<sub>4</sub>, satd. with NaCl, and extd. with five 10 ml. vols. AcEt. The org. exts. were evapd. to a vol. of 20 ml. and treated with petr. ether to give 1.6 g. III, m. 106.degree. (AcOEt-petr. ether). In a similar fashion, N-( $\gamma$ -hydroxy-isocaproyl)-L-leucine (IV), which sintered at 190.degree. (decompn.), was prepd. in 10% yield. A mixt. of 600 mg. III and L-leucine Me ester hydrochloride was converted by the anhydride method of Determann, et al. (CA 57, 9948c) to N-( $\gamma$ -hydroxyisocaproyl)-glycyl-L-leucine Me ester in 36% yield. In a similar manner, 390 mg. IV and L-tyrosine benzyl ester tosylate were converted to 50% N-( $\gamma$ -hydroxyisocaproyl)-L-leucyl-L-tyrosine benzyl ester (V), m. 154-5.degree.. Total hydrolysis of V gave equal amts. leucine (VI) and tyrosine. The  $\gamma$ -hydroxyisocaproyl group was found most difficult to remove in the case of V, 10 mg. of which was dissolved in 0.2 ml. 50% CF<sub>3</sub>CO<sub>2</sub>H soln., kept 12 hrs. at room temp., and evapd. in a desiccator over concd. H<sub>2</sub>SO<sub>4</sub> and KOH. The residue was shown by electrophoresis to be L-leucyl-L-tyrosine benzyl ester. For the prepn. of 3-nitrophenoxycarbonyl derivs., 38 g. COCl<sub>2</sub> was dissolved in 160 ml. ice-cold abs. C<sub>6</sub>H<sub>6</sub>. The soln. was stirred, treated dropwise during 2 hrs. with a soln. of 30 g. m-O<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>OH (VII) and 36.5 g. PhNMe<sub>2</sub> in AcOEt at 5-10.degree., and stirred 12 hrs. at room temp. The mixt. was extd. 3 times with 100-ml. vols. N HCl, washed with 100 ml. H<sub>2</sub>O, and the org. phase was evapd. to dryness in vacuo to give 74% 3-O<sub>2</sub>NC<sub>6</sub>H<sub>4</sub>O<sub>2</sub>CCl (VIII), b13 150-1.degree.. A mixt. of 6.5 g. VI, 4 g. MgO, 100 ml. H<sub>2</sub>O, and 30 ml. Et<sub>2</sub>O was cooled in ice, stirred vigorously, and treated during 30 min. with a mixt. of 10 g. VIII dild. with an equal vol. Et<sub>2</sub>O. The mixt. was acidified with 30 ml. concd. H<sub>2</sub>SO<sub>4</sub> and extd. with three 30-ml. vols. AcOEt. The combined org. exts. were washed 3 times with 20-ml. vols. 2N HCl and 5 times with 20-ml. vols. H<sub>2</sub>O, dried, and evapd. to dryness in vacuo at 30.degree. to give 62% 3-nitrophenoxycarbonyl-L-leucine, mixed with some VII. In a similar fashion, L-phenylalanine (IX) was converted to 52% 3-nitrophenoxycarbonyl deriv. (X), m. 118.degree. (AcOEt-petr. ether). A soln. of 3.3 g. X and 0.89 g. NH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>Me in 50 ml. abs. tetrahydrofuran (XI) was cooled to -15.degree., stirred, and treated with 0.92 ml. POCl<sub>3</sub>, followed at once by 2.8 ml. Et<sub>3</sub>N. The mixt. was stirred 1 hr. at -15.degree., treated with 20 ml. H<sub>2</sub>O, and concd. in vacuo. The residue was extd. with three 10-ml. vols. AcOEt, which exts. were combined and washed with 10 ml. H<sub>2</sub>O, 10 ml. 5% aq. NaHCO<sub>3</sub>, and three 10-ml. vols. H<sub>2</sub>O, dried, and evapd. to dryness in vacuo at 30.degree. to give 32.5 g. 3-nitrophenoxycarbonyl-L-phenylalanylglycine Me ester, m. 162.degree. (AcOEt-petr. ether and MeOH-H<sub>2</sub>O), still mixed with VII. Et<sub>3</sub>N (1.43 ml.) and then 0.95 ml. ClCO<sub>2</sub>Et was added with shaking to a soln. of 3.3 g. X in 20 ml. XI, and the mixt. cooled to -15.degree.. After 8 min., the mixt. was warmed to 0.degree., treated with a soln. of 2.7 g. L-alanylglycine benzyl ester hydrochloride in 30 ml. XI and 20 ml. H<sub>2</sub>O (mixed just before addn. with 1.43 ml. Et<sub>3</sub>N), and removed from the cooling bath and shaken until there was a strong evolution of CO<sub>2</sub>. The soln. was evapd. in vacuo at 40.degree., and the residue mixed with 60 ml. AcOEt, which phase was washed with two 10-ml. vols. each of N HCl and H<sub>2</sub>O, two vols. 5% aq. NaHCO<sub>3</sub>, and five 10-ml. vols. H<sub>2</sub>O, dried, and evapd. in vacuo at

30.degree. to give 75% 3-nitrophenoxy-carbonyl-L-phenylalanyl-glycyl-L-alanine benzyl ester, m. 180-1.degree. (MeOH-H<sub>2</sub>O). The photolysis of X was performed by irradiating a soln. of 0.98 g. X in 200 ml. 25% aq. XI with a water-cooled high-pressure-Hg lamp (Hanau Q 81). Paper electrophoresis showed hydrolysis to be complete after 4 hrs., and 89% IX was recovered. Expts. to cleave the 3-nitrophenoxy-carbonyl group with aq. Et<sub>3</sub>N led to hydantoin formation. 18 references.

ST NITROPHENOXYCARBONYL PEPTIDE PROTECTIVE GROUP; HYDROXYISOCAPROYL PEPTIDE PROTECTING GROUPS; PEPTIDE PROTECTING GROUPS HYDROXYISOCAPROYL; NITROPHENOXYCARBONYL PEPTIDE PROTECTIVE GROUP; PEPTIDE PROTECTING GROUPS HYDROXYISOCAPROYL; HYDROXYISOCAPROYL PEPTIDE PROTECTING GROUPS

IT Peptides, preparation

RL: PREP (Preparation)

(4-hydroxy-4-methylvaleryl group and (m-nitrophenoxy)carbonyl group as protective groups for amino group in)

IT Amino group

(4-hydroxy-4-methylvaleryl group and (m-nitrophenoxy)carbonyl group as protective groups for amino group in peptide prepn.)

IT (m-Nitrophenoxy)carbonyl group

4-Hydroxy-4-methylvaleryl group

(as protective group for amino group in peptide prepn.)

IT 14235-02-0P 14235-03-1P 14235-04-2P 14235-05-3P 14235-06-4P

14235-07-5P 14235-13-3P 14317-60-3P

RL: SPN (Synthetic preparation); PREP (Preparation)

(prepn. of)

IT 14235-06-4P 14235-13-3P


RL: SPN (Synthetic preparation); PREP (Preparation)

(prepn. of)

RN 14235-06-4 HCAPLUS

CN Alanine, N-carboxy-3-phenyl-, N-(m-nitrophenyl) ester, L- (8CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 14235-13-3 HCAPLUS

CN Leucine, N-carboxy-, N-(m-nitrophenyl) ester, L- (8CI) (CA INDEX NAME)

Absolute stereochemistry.

